

# IODIS

**Hand Sanitizing**—in food handling and meat cutting operations. Rinse and immerse hands in a 1 oz. IODIS per 5 gals. of water solution (25 ppm titratable iodine)

**Sanitizing Poultry Drinking Water** - Add 1 ounce of IODIS per 10 gallons of drinking water.

**Shoe Bath Sanitizer**—To prevent tracking harmful organisms into animal areas, scrape shoes and place in a 3 oz. per gallon solution of IODIS for 30 seconds prior to entering area.

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS & DOMESTIC ANIMALS  
DANGER**

Corrosive. Causes eye damage and skin irritation. Harmful if swallowed. Do not get in eyes, on skin or on clothes. Wear goggles or face shield and rubber gloves when handling. Wash thoroughly with soap and water after handling. Avoid contamination of food and foodstuffs. Avoid breathing spray mist. Remove contaminated clothing and wash before reuse.

**ENVIRONMENTAL HAZARDS:** This product is toxic to fish. "Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance contact your State Water Board or Regional Office of the EPA."

**PHYSICAL AND CHEMICAL HAZARDS:** Do not use, pour, spill or store near heat or open flame.

**STORAGE AND DISPOSAL**

- Storage:** Store only in tightly closed, original container in a secure area inaccessible to children. Do not contaminate water, food, or feed by storage or disposal.
- Pesticide Disposal:** Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the hazardous waste representative at the nearest EPA Regional Office for guidance.
- Container Disposal:** Do not reuse empty container. Triple rinse empty container with water. Return metal drums to reconditioner or puncture and dispose of in a sanitary landfill or by other procedures approved by state and local authorities. Plastic containers may be disposed of in a sanitary landfill, incinerated or, if allowed by local authorities, by burning. If burning stay out of smoke.
- General:** Consult Federal, State or local disposal authorities for approved alternative procedures.



Zephyr Cove, NV 89448

Manufactured by

**PRESERVE INTERNATIONAL** ZEPHYR COVE, NV 89448, USA Telephone (800) 995-1607

**Active Ingredient**

Iodine\* .....1.75%

**Inert Ingredients** .....98.25%

\* from Nonylphenoxypoly (ethyleneoxy) ethanol-iodine complex

**KEEP OUT OF REACH OF CHILDREN  
DANGER**

**STATEMENT OF PRACTICAL TREATMENT**

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse.

If swallowed, drink promptly large quantities of water. Avoid alcohol. Call a physician immediately.

**NOTE TO PHYSICIAN:** Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

**SEE LEFT PANEL FOR ADDITIONAL  
PRECAUTIONARY STATEMENTS**

EPA Reg. No. 66171-10

EPA Est. No. 66171-TN-001

**Net Contents**

**NFPA HAZARD CLASSIFICATION**

HEALTH	FIRE	REACTIVITY	SPECIFIC
2	0	0	Acid

**DOT SHIPPING NAME**

**CORROSIVE LIQUIDS, N.O.S.,**  
(Contains Phosphoric Acid),  
8, UN1760, PGII

**DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

For disinfection remove gross filth or heavy soil. Apply IODIS with a cloth, mop or mechanical coarse spray device. When applied with a mechanical coarse spray device, surface must be sprayed until thoroughly wetted. Treated surfaces must remain wet for 10 minutes. Fresh solution should be prepared daily or more often if the solution becomes diluted or soiled.

IODIS is a proven "one-step" cleaner-disinfectant-sanitizer-deodorizer in the presence of moderate amounts of organic soil and 400 ppm hard water.

**Disinfection—Hospitals, Veterinary Clinics, Other Health Care Institutions, Institutions, Industry, Schools, Farm Premises and Poultry Houses:**


Remove gross filth or heavy soil. Apply 1/2 oz. per gallon IODIS to walls, floors and other hard (inanimate) non-porous surfaces such as tables, chairs, countertops, tile, porcelain and bed frames. Allow surfaces to remain wet for at least 10 minutes. For heavily soiled areas, a preliminary cleaning is required.

For farms and poultry use, prior to disinfection, all poultry, other animals and their feeds must be removed from the premises. This includes emptying all troughs, racks and other feeding and water appliances. Removed all litter and droppings from floors, walls and other surfaces occupied or traversed by poultry or other animals.

After 10 minutes disinfection, ventilate buildings, coop and other closed spaces. Do not house poultry, or other animals or employ equipment until treatment has been absorbed, set or dried.

**Sanitizing—Meat and food processing equipment and utensils:** Remove heavy soils by flushing, scraping, or mopping. Wash with 1 oz. per gallon of IODIS (or other good cleaner). Rinse with clean water and sanitize surfaces with 1 oz. IODIS per 5 gallons of water (25 ppm titratable iodine). Surface should remain in contact with solution for 1-2 minutes. Drain well and air dry. No potable water rinse is required.

**Sanitizing Food Grade Egg Shells :** To sanitize clean shell eggs intended for food or food products, spray with a solution of 1 ounce of product in 5 gallons of water (providing 25 ppm active). The solution must be equal to or warmer than the eggs, but not to exceed 130°F. Wet eggs thoroughly and allow to drain. Eggs that have been freshly washed may be sanitized with this compound only if the eggs are rinsed prior to application of the compound. A subsequent potable water rinse is not required. Eggs must be reasonably dry before casing or breaking. The solution must not be reused for sanitizing eggs.



66171-10  
3/7/2000  
Page 121