



U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Antimicrobials Division (7510C)
1200 Pennsylvania Avenue NW
Washington, D.C. 20460

EPA Reg. Number:
66171-102

Date of Issuance:
DEC 4 2013

Term of Issuance:
Conditional

Name of Pesticide Product:
Iodis 1.75

NOTICE OF PESTICIDE:
 Registration
 Reregistration

(under FIFRA, as amended)

Name and Address of Registrant (include ZIP Code):

Preserve International
P. O. Box 17003
Reno, NV 89511

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.


This product (OPP Decision No. D-477701) is conditionally registered in accordance with FIFRA sec 3(c)(7)(A) provided that you:

1. Submit and/or cite all data required for registration of your product under FIFRA sec. 3(c)(7)(A) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for re-registration of your product under FIFRA section 4.
2. Submit three (3) copies of your final printed labeling before distributing or selling the product bearing the revised labeling.
If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records. Should you have any questions regarding this letter, please contact me at (703) 308-8062 or by e-mail at Fuller.Demsom@epa.gov. When submitting information or data in response to this letter, a copy of this letter should accompany the submission to facilitate processing.

Enclosure: EPA stamped label.

Signature of Approving Official:


Demson Fuller
EPA Product Manager 32
Regulatory Management Branch II
Antimicrobials Division (7510P)

Date:

DEC 4 2013

IODIS 1.75

Disinfectant-Cleaner-Sanitizer-Deodorizer
With 5% Organic Iodine and 400 ppm Hard Water Tolerance

For Hospital, Institutional, Industrial, School, Agriculture, and Dairy
Active ingredients.....1.75%
Inert ingredients.....98.25%
*from Nonylphenoxypoly (ethylenoxy) ethanol-iodine complex

KEEP OUT OF REACH OF CHILDREN
DANGER

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

FIRST AID

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes; remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye; call a poison control center or doctor for treatment advice.
IF ON SKIN OR CLOTHING: Take off contaminated clothing; rinse skin immediately with plenty of water for 15-20 minutes; call a poison control center or doctor for treatment advice.
IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible; call a poison control center or doctor for further treatment advice.
IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice; have person sip a glass of water if able to swallow; do not induce vomiting unless told to do so by the poison control center or doctor; do not give anything by mouth to an unconscious person.
HOT LINE NUMBER: Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1 (800) 255-3924 for emergency medical treatment information. For information on this pesticide (including health concerns, medical emergencies, or pesticide incidents), call the National Pesticide Telecommunications Network at 1 (800) 858-7378.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

SANITIZING GLASSES, DISHWARE, POTS, PANS, AND UTENSILS IN FOOD SERVICE OPERATIONS:

1. Pre-rinse or pre-scrape equipment and utensils. If needed, pre-soak to remove gross food particles and soil.
2. Wash equipment and utensils in lukewarm water that contains adequate amount of a good detergent or cleaner. Keep this water clean by changing it frequently.
3. Rinse equipment and utensils free of wash water prior to sanitization.
4. Sanitize equipment and utensils by immersing them in a solution of 1 ounce Iodis 1.75 per 5 gallons of water (providing 25ppm Iodis available iodine). Allow 1-2 minutes contact time (or longer if required by local sanitation codes). Fresh solution should be prepared if the solution becomes diluted or soiled. Use an approved iodine test kit to check for proper concentrations.
5. After required contact time of 1-2 minutes or more, allow equipment and utensils to drain and then air dry. Do not rinse before use.

SANITIZING PREVIOUSLY CLEANED FOOD CONTACT SURFACES:


Sanitize previously cleaned and rinsed hard, non-porous surfaces such as glass, metal, plastic and porcelain with a solution of 1 ounce Iodis 1.75 per 5 gallons water. Immerse equipment with sanitizing solution and allow 1-2 minutes of contact time. Drain solution from equipment and does not require a final rinse with potable water in accordance with Federal Food Additive Regulation 178.1010.

SANITIZING POULTRY AND LIVESTOCK DRINKING WATER:

Add 1 ounce of Iodis 1.75 per 10 gallons of drinking water.

NET CONTENTS: _____ Batch No.: _____
EPA Reg. No. 66171- EPA Est. No. 66171-TN-001, 66171-CA-001 (First letter of batch number indicates establishment)

www.preserveinternational.com



Manufactured By:

PRESERVE INTERNATIONAL

Reno, NV 89511 • (800) 995-1607 • www.preserveinternational.com

FOR SANITIZING DAIRY EQUIPMENT AND UTENSILS: 1. Wash the equipment to be sanitized using a Presolve International manual cleaner. Be sure to follow label directions. Rinse and allow to drain. 2. Prepare a solution by mixing 1 ounce of Iodis 1.75 sanitizer for each 5 gallons of water, or add 1/2 ounce of sanitizer per 10 quarts of water. 3. Immerse equipment: milking claws, liners, this will provide 25ppm Iodis available iodine. 4. For equipment that can't be immersed (bulk milk cups, into solution making sure that all surfaces are in contact with the solution. A contact time of 1-2 minutes is required. 5. For equipment that can't be immersed (bulk milk cups, receiver groups, etc.) coarse spray a 25ppm Iodis solution onto all pre-cleaned surfaces for a sparkling, spot free shine. 6. Allow solution to air dry. 7. Iodis 1.75 solution if sanitizing solution fades noticeably or becomes diluted or soiled. 8. Repeat this is its own indicator of germicidal activity. The amber or yellow color shows the presence of the active ingredient, Iodine.

SHOE BATH SANITIZER:

To prevent tracking harmful organisms into animal areas, scrape shoes and place in a 3 oz. per gallon solution of Iodis 1.75 for 30 seconds prior to entering area.

DEODORIZING:

Iodis 1.75 destroys many odors as it sanitizes and disinfects. For general deodorant applications (garbage pails, refuse containers, etc.), swab surface to be treated with a solution containing 1/2 ounce Iodis 1.75 per gallon of water. Allow treated surfaces to drain and air dry.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS & DOMESTIC ANIMALS DANGER

DANGER Corrosive. Causes eye damage and skin irritation. Harmful if swallowed. Do not get in eyes, on skin or on clothes. Wear goggles or face shield and rubber gloves when handling. Wash thoroughly with soap and water after handling. Avoid contamination of food and foodstuffs. Avoid breathing spray mist. Remove contaminated clothing and wash before reuse.
ENVIRONMENTAL HAZARDS: This product is toxic to fish and aquatic organisms. "Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans or other waters unless in accordance with the requirements of a National Pollution Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance contact your State Water Board or Regional Office of the EPA."
PHYSICAL AND CHEMICAL HAZARDS: Do not use, pour, spill or store near heat or open flame.

STORAGE AND DISPOSAL

Store in tightly closed, original container in a secure area inaccessible to children. Do not contaminate water, food, or feed by storage or disposal. Pesticide Disposal: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinseate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the hazardous waste representative at the nearest EPA Regional Office for guidance.
Container Handling: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available or offer for reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

(For greater than 5 gallons) Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the container over onto its other end and tip it back and forth several times. Repeat this procedure two more times. (For 5 gallons or less) Triple rinse (or equivalent) promptly after emptying. Triple rinse as follows: Empty the container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinseate into application equipment or a mix tank or store rinseate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.
General: Consult Federal, State or local disposal authorities for approved alternative procedures.

DOT SHIPPING NAME
UN 3264, Corrosive Liquid, n.o.s.
(Contains Phosphoric Acid), 8, PG III

Concentrated broad spectrum biocidal for use as a one-step cleaner-disinfectant and no rinse sanitizer in: • Veterinary Clinics • Poultry and Livestock Drinking Water • Sanitizing Commercial Eggs • Egg Processing Plants • Dairy & Poultry Farms • Meat and Poultry Plants • Food Processing Plants • Kennels, Pet Shops & Zoos • Hand Sanitizing in Food Plants • Animal Handling Facilities.
Used as directed, Iodis 1.75 is effective in hard water (up to 400ppm as CaCO₃) and in 5% organic serum load against the following viruses, bacteria, and pathogenic fungi: Polio 1, Vaccinia, Herpes Simplex Type 1, Influenza A2/Durham 305, Canine Parvovirus; bactericidal against *Salmonella choleraesuis*, *Staphylococcus aureus*, *Pseudomonas aeruginosa*, and *E. coli*; fungicidal against *Trichophyton mentagrophytes* (the athlete's foot fungus). On pre-cleaned contact surfaces, Iodis 1.75 is effective against *Salmonella choleraesuis*, *Listeria monocytogenes* and *Escherichia coli*, O157:H7, Hemorrhagic.

DIRECTIONS FOR USE:

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. For disinfection remove gross filth or heavy soil. Apply Iodis 1.75 with a cloth, mop or mechanical coarse spray device. When applied with a mechanical coarse spray device, surface must be sprayed until thoroughly wetted. Treated surfaces must remain wet for 10 minutes. Fresh solution should be prepared daily or more often if the solution becomes diluted or soiled.

Disinfection: Veterinary Clinics, Fish Hatcheries, Schools, Farm Premises and Poultry Houses:

Remove gross filth or heavy soil. Apply 1/2 oz. per gallon Iodis 1.75 to walls, floors and other hard (non-porous) surfaces such as tables, chairs, countertops, tile, porcelain and bed frames. Allow surfaces to remain wet for at least 10 minutes. For heavily soiled areas, a preliminary cleaning is required.

For farms and poultry use, prior to disinfection, all poultry, other animals and their feeds must be removed from the premises. This includes emptying all troughs, racks and other feeding and water appliances. Remove all litter and droppings from floors, walls and other surfaces occupied or traversed by poultry or other animals.

After 10 minutes disinfection, ventilate buildings, coop and other closed spaces. Do not use poultry, or other animals or employ equipment until treatment has been absorbed, set or dried.

SANITIZING FOOD PROCESSING EQUIPMENT AND UTENSILS:

Remove heavy soils by flushing, scraping, or mopping. Wash with 1 oz. per gallon of Iodis 1.75 (or other good cleaner). Rinse with clean water and sanitize surfaces with 1 oz. Iodis 1.75 per 5 gallons of water (25ppm Iodis available iodine). Surface should remain in contact with solution for 1-2 minutes. Drain well and air dry. No potable water rinse is required.

SANITIZING FOOD GRADE EGG SHELLS:

To sanitize clean shell eggs intended for food or food products, spray with a solution of 1 ounce of product in 5 gallons of water (providing 25 ppm active). The solution must be equal to or warmer than the eggs, but not to exceed 130°F. Wet eggs thoroughly and allow to drain. Eggs that have been freshly washed may be sanitized with this compound only if the eggs are rinsed prior to application of the compound. A subsequent potable water rinse is not required. Eggs must be reasonably dry before casing or breaking. The solution must not be reused for sanitizing eggs.


SANITIZING MILKING EQUIPMENT AND BULK TANKS:

Before and after each milking, flush milking equipment, utensils, and bulk tanks with a solution of 1 ounce Iodis 1.75 per 5 gallons of water (providing 25ppm Iodis available iodine). Allow a 1-2 minute contact time. Drain thoroughly and air dry before use.

REMOVAL OF MILKSTONE AND HARD WATER DEPOSITS:

Yellow color may appear on equipment after the usual sanitizing procedure. This may indicate the presence of milkstone (or other deposits) which can be removed by applying a solution containing equal parts of Iodis 1.75 and white vinegar. Wash thoroughly with brush and then rinse with potable water. Daily use of Iodis 1.75 will help prevent the formation of milkstone and hard water deposits.

December 4, 2013
Under the Federal Insecticide, Fungicide, and Pesticide Act as amended, for this pesticide, registered under EPA Reg. No. 66171-1008



© 2013