

10/12/2006

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

WASHINGTON, D.C. 20460

OFFICE OF PREVENTION, PESTICIDES AND TOXIC SUBSTANCES

OCT 1 2 2006

Ms. Jane Rothwell Makhteshim Agan of North America, Inc. 4515 Falls of Neuse Road, Suite 300 Raleigh, NC 27609

Dear Ms. Rothwell:

Subject: Amendment - add supplemental label permitting use on caneberries in Ohio

Diazinon AG500

EPA Reg. No. 66222-9

Your submission dated October 3, 2006

The amendment referred to above, submitted in connection with registration under FIFRA section 3(c)(7(a), is acceptable and a stamped copy of the supplemental label is enclosed for your records.

Sincerely,

George LaRocca. Product Manager 13 Insecticide Branch Registration Division (7505P)



ACCEPTED

OCT 1 2 2006

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 6622279

SUPPLEMENTAL LABELING

RESTRICTED USE PESTICIDE DUE TO AVIAN AND AQUATIC TOXICITY

FOR RETAIL SALE TO AND USE ONLY BY CERTIFIED APPLICATORS OR PERSONS UNDER THEIR DIRECT SUPERVISION AND ONLY FOR THOSE USES COVERED BY THE CERTIFIED APPLICATOR'S CERTIFICATION.

DIAZINON AG500

Insecticide
EPA REG NO. 66222-9
FOR CONTROL OF CERTAIN INSECTS ON CANEBERRIES

ACTIVE INGREDIENT: Diazinon:		% By Wt.
O,O-diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphoroth	hioate	48.0%
INERT INGREDIENTS*	,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,	52.0%
	•	TOTAL 100.0%

Contains 4 lbs. Diazinon per gallon *Contains xylene range aromatic solvent

KEEP OUT OF REACH OF CHILDREN CAUTION

DIRECTIONS FOR USE

It is violation of federal law to use this product in a manner inconsistent with its labeling.

This labeling must be in the possession of the user at the time of pesticide application. All applicable directions, restrictions and precautions on the EPA registered label are to be followed.

CROP	PEST	RATE	COMMENTS
CANEBERRIES (Blackberries, Boysenberries, Dewberries, Loganberries, Raspberries) (California, Ohio, Oregon, and Washington Only) (7)	Raspberry Fruitworm	1 qt. per acre	Apply when insects occur. Apply in a minimum of 100 gals. of water per acre (200 gals. per acre maximum). Make application when blossom buds separate.
	Raspberry Crown Borer (Raspberry Root Borer)	2 qts. per acre	Apply a single application in a minimum of 100 gals. of water per acre as a drench to the crown and lower canes in the spring before buds break.
	 Do not apply more than 2 qts. of product per acre per application. Make a maximum of one application per year regardless of the target pest. The REI is 5 days 		