

ACCEPTED

MAY 15 2002

SUPPLEMENTAL LABELING

**RESTRICTED USE PESTICIDE
DUE TO AVIAN AND AQUATIC TOXICITY**

FOR RETAIL SALE TO AND USE ONLY BY CERTIFIED APPLICATORS OR PERSONS UNDER THEIR DIRECT SUPERVISION AND ONLY FOR THOSE USES COVERED BY THE CERTIFIED APPLICATOR'S CERTIFICATION.

DIAZOL® 50W
A Wettable Powder Insecticide
EPA REG NO. 66222-10

FOR USE ON FIGS IN CALIFORNIA, WATERCRESS IN HAWAII, AND FILBERTS IN OREGON AND WASHINGTON

ACTIVE INGREDIENT:	% BY WT.
Diazinon [O O-diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate]	50%
INERT INGREDIENTS:	50%
	TOTAL 100%

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

DIRECTIONS FOR USE

It is violation of federal law to use this product in a manner inconsistent with its labeling. This labeling must be in the possession of the user at the time of pesticide application. All applicable directions, restrictions, and precautions on the EPA-registered label are to be followed.

Crop	Pest	Rate	Comments
FIGS (California Only)	Vinegar flies- <i>drosophila</i> spp., Dried fruit beetles	Apply 1 lb. per 100 gals of water not to exceed 200 gals. of water per application	Apply as infestations occur. Repeat application after 7 days if necessary.
	<ul style="list-style-type: none"> Do not apply more than 2 pounds of product per acre per application and no more than 6 pounds of product per acre per year. To avoid illegal residues, allow a minimum of 5 days between the last application and harvest. 		
FILBERTS (HAZELNUTS) (Oregon and Washington Only)	Filbert Leafrollers, Aphids	Apply 2-4 lbs. in 250-400 gals. of water per acre	Apply as a thorough cover spray. Repeat application after 14 days if necessary.
	<ul style="list-style-type: none"> Do not apply more than 4 pounds of product per acre per application and no more than 12 pounds per year. To avoid illegal residues, allow a minimum of 21 days between the last application and harvest. 		
WATERCRESS (Hawaii Only)	Cyclamen Mites	Apply 1 pound per 100 gals. of water per acre.	Apply as a thorough cover spray. Spray should be directed into the crowns and growing tips of plants. If infestation is heavy, make a second application within 10 days after the first application.
	<ul style="list-style-type: none"> Do not apply more than 1 pound of product per acre per application and no more than 2 pounds of product per acre per season. To avoid illegal residues; allow a minimum of 5 days between the last application and harvest. 		