



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

APR 13 2009

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

Anne Stout
Makteshim Agan of North America, Inc.
4515 Falls of Neuse Rd., Ste. 300
Raleigh, NC 27609

Subject: Change Primary Brand Name
MANA Cotoran 4L
EPA Reg. No. 66222-32
Application Dated April 7, 2009

Dear Ms. Stout:

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is acceptable, provided you make the following changes:

1. Change the heading from "Inert Ingredients" to "Other Ingredients".
2. On page 4 under Marestail (Horseweed), correct the typo from "Ignite" to "Ignite".
3. On page 6 under Specific Weed Problems, correct the typo from "Staple LS" to "Staple LX".
4. Revise the Pesticide Disposal text to read "Wastes resulting from the use of this product **must** be disposed of on site or at an approved waste disposal facility."
5. Revise the Refillable Container text to read "Refill this container with ~~diazinon~~ **fluometuron** only."

The amended label supercedes all previously accepted labels. A stamped copy of the label is enclosed for your records. Submit one copy of final printed labeling incorporating the above changes before you release this product for shipment.

Sincerely,

A handwritten signature in black ink, appearing to read "Jim Tompkins".

Jim Tompkins
Product Manager 25
Herbicide Branch
Registration Division (7505P)

2/9

MANA COTORAN® 4L HERBICIDE

For Preemergence and Postemergence Weed Control in Cotton
Not for use in Sumpter County, Georgia and in the counties of Colbert, Lawrence, Limestone,
Madison, and Tuscaloosa in Alabama.

ACTIVE INGREDIENT	% BY WT.
Fluometuron: 1,1-dimethyl-3-(a,a,a-trifluoro-m-tolyl) urea	41.7%
INERT INGREDIENTS	58.3%
TOTAL	100.0%

MANA Cotoran 4L Contains 4 Pounds Active Ingredient Per Gallon

ACCEPTED
with COMMENTS
in EPA Letter Dated
APR 13 2009

KEEP OUT OF REACH OF CHILDREN CAUTION

Manufactured for:
Makhteshim Agan of North America, Inc.
4515 Falls of Neuse Rd., Suite 300
Raleigh, NC 27609

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

66222-32
EPA Est. No. _____

EPA Reg. No. 66222-32

NET CONTENTS: _____ GALLON(S)

FIRST AID	
IF IN EYES	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 to 20 minutes • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
IF ON SKIN OR CLOTHING	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 to 20 minutes. • Call a poison control center or doctor for treatment advice.
IF SWALLOWED	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice • Have person sip a glass of water if able to swallow • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
IF INHALED	<ul style="list-style-type: none"> • Move person to fresh air • If person is not breathing, call 911 or ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact PROSAR at 1-877-250-9291 for emergency medical treatment information.	
NOTE TO PHYSICIAN: If MANA Cotoran 4L is ingested, lavage stomach or induce vomiting. A slurry of activated charcoal in water can be left in the stomach. Give a saline laxative and supportive therapy.	
See label for Additional Precautions and Directions for Use.	

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS (AND DOMESTIC ANIMALS) CAUTION

Harmful if swallowed, inhaled or absorbed through the skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Avoid breathing spray mist.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical resistant to this product are polyethylene or polyvinyl chloride. If you want more options, follow the instructions for Category A on an EPA chemical-resistance category selection chart.

Mixers and loaders supporting aerial application must wear:

- coveralls over long-sleeved shirt and long pants,
- chemical-resistant gloves,
- chemical-resistant footwear and socks, and
- a chemical-resistant apron

All other mixers, loaders, applicators, flaggers and other handlers must wear:

- long-sleeved shirt,
- long pants
- shoes and socks, and
- chemical-resistant gloves (except applicators and flaggers)

See engineering controls for additional requirements and options.

Follow manufacturer's instructions for cleaning / maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Discard clothing or other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

Engineering Controls Statements:

When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

Enclosed cockpits for Aerial Applicators: Pilots must use an enclosed cockpit that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(6)].

Important: When reduced PPE is worn because a closed system is being used, handlers must be provided all PPE specified above for 'Applicators and other handlers' and have such PPE immediately available for use in an emergency, such as spill or equipment breakdown.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing / PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This chemical, fluometuron is known to leach through soil into ground water under certain conditions, including karst geology areas, as a result of registered uses. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground water contamination.

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters or rinsate.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its label.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation. Do not apply this product through any type of irrigation system.

Not for use in Sumpter County, Georgia and in the counties of Colbert, Lawrence, Limestone, Madison, and Tuscaloosa in Alabama.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours.

Exception: If product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls worn over short-sleeved shirt and short pants
- Chemical-resistant gloves made of any waterproof material
- Chemical-resistant footwear plus socks
- Protective eyewear

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECUATIONS ON THIS LABEL MAY RESULT IN CROP INJURY, POOR WEED CONTROL, AND/OR ILLEGAL RESIDUES.

GENERAL INFORMATION

MANA Cotoran 4L is effective for both annual grass and annual broadleaf weed control in cotton.

Weed Control

MANA Cotoran 4L applied according to directions will selectively control annual weeds in cotton, such as:

Annual Grass Weeds		
Barnyardgrass	Goosegrass	Fall panicum
Foxtail	Crowfootgrass	Signalgrass (Brachiaria)
Crabgrass	Ryegrass	
Annual Broadleaf Weeds		
Buttonweed	Lambsquarters	Ragweed
Cocklebur	Morningglory	Sesbania
Florida Pusley	Pigweed	Sicklepod
Groundcherry (Wright)	Prickly Sida (teaweed)	Smartweed
Jimsonweed	Purslane	Tumbleweed

MANA Cotoran 4L at rates up to 4 pints per acre in cotton is not effective for the control of established perennial weeds, such as johnsongrass, from rhizomes.

COTTON

MANA Cotoran 4L may be applied as a preplant or preemergence treatment to the surface of the soil, or lightly incorporated immediately after planting, or as a directed, semi-directed, or over-the-top postemergence application. For preemergence treatment only, MANA Cotoran 4L may be tank-mixed with most liquid nitrogen fertilizers. MANA Cotoran 4L may be combined with other herbicides in cotton as directed in the section **COMBINATIONS**.

PRECAUTIONS

The use of MANA Cotoran 4L preplant or preemergence in conjunction with a soil applied organophosphate insecticide or other systemic insecticide at planting may result in injury to cotton. In addition, MANA Cotoran 4L applied to cotton with any application of Command® (clomazone) herbicide may result in crop injury. To the extent consistent with applicable law, MANA will not be held responsible for losses or damage resulting from such uses.

The second application of MANA Cotoran 4L cannot be made earlier than 20 days following the initial application.

Rainfall that is sufficient to germinate seeds or for growing crops is adequate to activate the herbicidal activity of MANA Cotoran 4L; where dry weather conditions prevail, the herbicidal activity of MANA Cotoran 4L may be delayed or reduced. Cultivate if weeds develop.

Do not make more than two applications of MANA Cotoran 4L to the same crops or field in any 1 year. Do not plant crops other than cotton after the last application of MANA Cotoran 4L, without following the plantback intervals, or injury may result.

Plantback intervals for the following crops in areas other than AZ and NM* are as follows:

- 3 months for wheat
- 8 months for field corn, sweet corn, and peanuts
- 9 months for rice, grain, sorghum and soybeans
- 12 months for all other crops not listed above

*See statement below for AZ and NM plantback restrictions.

Do not apply within 60 days of harvest. Do not feed foliage from treated cotton plants to livestock.

APPLICATION SOIL RESTRICTIONS

Broadcast Rates per Pints Per Acre of MANA Cotoran 4L and Active Ingredient Equivalent

Soil Texture	Maximum One-time	Annual Maximum Number	Annual Maximum
--------------	------------------	-----------------------	----------------

	Application Rate	of Applications Per Year	Application Rate
Sand, loamy sand, and sandy loam soils	2 pts (1 lb ai/A)	2	4 pts (2 lb ai/A)
Loam, silt loam, silt, sandy clay loam, silty clay loam, and clay loam soils	3.2 pts. (1.6 lb ai/A)	2	6 pts (3 lb ai/A)
Sandy clay, silty clay, and clay soils	4 pts (2 lb ai/A)	2	6 pts (3 lb ai/A)

Mandatory Interval between Applications: There must be a minimum of 20 days between the first and second application of MANA Cotoran 4L.

Other State and Local Requirements:

Applicators must follow all state and local pesticide drift requirements regarding application of products containing the active ingredient fluometuron. Where states have more stringent regulations, they must be observed. Before applying MANA Cotoran 4L, refer to **Application Soil Restrictions Table** in the PRECAUTIONS section for specific application rate soil restrictions.

West TX only: Do not use MANA Cotoran 4L on sandy, loamy sand, or fine sandy loam soils. Do not use MANA Cotoran 4L on cotton planted in furrows.

AR, LA and MS only: Use 1.6 pints of MANA Cotoran 4L per acre on sandy loam soils low in organic matter.

AZ and NM only: Cotton, corn and grain sorghum can be planted the next spring. Do not plant treated areas to other crops until 1 year after last application. Do not use on sandy or coarse-textured soil of less than 1% organic matter.

Preplant Treatment (except AZ and NM)

MANA Cotoran 4L may be applied for burndown of existing annual weeds and residual control of weeds prior to planting cotton. Apply MANA Cotoran 4L at a rate of 2 to 4 pints per acre from 15 to 75 days prior to anticipated planting. On light silt and sandy soils, low in organic matter, use the lower rate. Apply before weeds germinate or before weed seedlings are more than 2 inches tall. If weeds are emerged prior to application, the addition of a surfactant is recommended. When emerged weeds are taller than 2 inches or weeds not listed on the MANA Cotoran 4L label are present, MANA Cotoran 4L may be tank mixed with other products registered for preplant applications in cotton. Consult tank mix product labels for specific rate instructions and precautions. Early preplant applications may require additional weed control before planting.

Before applying MANA Cotoran 4L refer to **Application Soil Restrictions Table** in the PRECAUTIONS section for specific application rate soil restrictions.

Preemergence applications of herbicides with a similar mode of action (such as diuron) following a preplant application of MANA Cotoran 4L may result in cotton injury. This is most critical where applications of MANA Cotoran 4L are made less than 30 days preplant, on coarse textured soils, and on soils low in organic matter.

Marestail (Horseweed): For improved residual control of marestail, apply a preplant burndown tank mix of glyphosate plus Clarity® plus MANA Cotoran 4L or Ignite® plus MANA Cotoran 4L.

Preplant Incorporated Use (AZ and NM)

MANA Cotoran 4L may be applied preplant as a broadcast spray at 3.2 pints per acre after the beds for preplant irrigation have been formed. Apply before the beds are dragged off (using a bed shaper set to remove 1 to 2 inches from beds) in preparation for planting. Furrow irrigation should precede the application and planting should follow as soon as possible. Plant into untreated soil after drag-off and return treated soil to the beds when furrows are reformed. Do not allow furrow irrigation water to pond over the beds.

Before applying MANA Cotoran 4L refer to **Application Soil Restrictions Table** in the PRECAUTIONS section for specific application rate soil restrictions.

Preemergence Treatment

Apply MANA Cotoran 4L at a rate of 2 to 4 pints per acre. On light silt and sandy soils, low in organic matter, use the lower rate.

Before applying MANA Cotoran 4L refer to **Application Soil Restrictions Table** in the PRECAUTIONS section for specific application rate soil restrictions.

Plantback intervals must be as follows:

- 3 months for wheat
- 8 months for field corn, sweet corn, and peanuts
- 9 months for rice, grain, sorghum and soybeans
- 12 months for all other crops not listed above

Postemergence Treatments

MANA Cotoran 4L may be applied postemergence to cotton, but preemergence to weeds, or postemergence to both cotton and weeds. MANA Cotoran 4L should be applied at rates of 2 to 4 pints per acre. Apply as a directed, semi-directed, or over-the-top spray. MANA Cotoran 4L may be applied to cotton plants at any stage of growth from 3 inches high to lay-by. Use the higher rate when applying postemergence to weeds; weeds should be 2 inches high or less. Use of a surfactant that is approved for use on food and feed crops and is compatible with MANA Cotoran 4L when applied to cotton, may enhance the postemergence herbicidal activity of Cotoran 4L. The surfactant rate should not exceed 0.5% of the carrier volume (1 pint per 25 gallons of water).

Before applying MANA Cotoran 4L refer to **Application Soil Restrictions Table** in the PRECAUTIONS section for specific application rate soil restrictions.

Combinations / Tank Mix Partners

Parallel™ PCS:

Use in a tank mixture with MANA Cotoran 4L preemergence in cotton for control of all weeds as listed on both labels. This combination will also control spotted spurge, hyssop spurge, nodding spurge, and prostrate spurge. Apply to the soil surface at planting or after planting, but before weeds or crop emerge, using the appropriate rates from the following table.

MANA Cotoran 4L + Parallel PCS – Cotton (AR, LA, MS, Eastern OK, TN, Bootheel of MO, Gulf coast, Rio Grande Valley, and Eastern TX)

Soil Texture	Broadcast Rates Per Acre		
	MANA Cotoran 4L	Parallel PCS	
		Area 1**	Area 2***
Sand, Loamy sand	DO NOT USE		
Sandy loam*	2 pts	.75-1.0 pts	.85 – 1.0 pts
Loam, Silt loam, Silt	2 – 3.2 pts	1.0 – 1.33 pts	1.0 – 1.33 pts
Fine Soil	3.2 – 4 pts.	1.0 – 1.33 pts	1.33 pts

*Use on sandy loam soils only in eastern OK, Gulf Coast, Rio Grande Valley and eastern TX.

** Area 1 = AR, LA, MS, Bootheel of MO and TN

*** Area 2 = Eastern OK, Gulf Coast, Rio Grande Valley, and Eastern TX

Precautions:

- (1) Do not apply MANA Cotoran 4L + Parallel PCS on sand or loamy sand soils or in areas where water is likely to "pond" over the bed, or crop injury may occur.
- (2) Do not apply broadcast to furrow-planted cotton.
- (3) The use of MANA Cotoran 4L preplant or preemergence in conjunction with a soil applied organophosphate insecticide or other systemic insecticide at planting may result in injury to cotton. In addition, MANA Cotoran 4L applied to cotton with any application of Command® (clomazone) herbicide may result in crop injury. To the extent consistent with applicable law, MANA will not be held responsible for losses or damage resulting from such uses.
- (4) Do not use on Taloka silt loam, or crop injury may occur.

Before applying MANA Cotoran 4L refer to **Application Soil Restrictions Table** in the PRECAUTIONS section for specific application rate soil restrictions.

Refer to the Parallel PCS label for further instructions, precautions, and limitations.

Note: Do not feed treated forage to livestock, or graze treated areas.

Rotational Crops: Refer to the crop rotation instructions for MANA Cotoran 4L alone on this label and for Parallel PCS alone on the Parallel PCS label.

Triflurex® HFP:

Use in a tank mixture with MANA Cotoran 4L (all areas except AZ) or before an overlay with MANA Cotoran 4L in cotton for control of all weeds listed on both labels.

For the **tank mixture**, follow soil preparation and incorporation procedures for Triflurex HFP. Apply MANA Cotoran 4L + Triflurex HFP tank mix at rates specified below in a minimum of 15 gallons of water per acre.

MANA Cotoran 4L + Triflurex HFP Tank Mixture – Cotton (All areas except AZ)

Soil Texture	Broadcast Rates Per Acre	
	MANA Cotoran 4L	Triflurex HFP
Sand, Loamy sand, Sandy loam	2 pts	1 pt
Loam, Silty clay loam*, Silt loam, Silt, Sandy clay loam,* Clay loam	3.2 pts	1.5 pts

7/9

Clay, clay loam, Silty clay loam*, Silty clay, Sandy clay, Sandy clay loam*	4 pts	2 pts
---	-------	-------

*Silty clay loam and sandy clay loam are transitional soils. If these soils are predominantly sand or silt, they are usually classified as medium-textured soils. If predominantly clay, they are classified as fine-textured soils.

For the overlay, apply MANA Cotoran 4L as a preemergence soil surface treatment as specified on this label at 2 to 4 pints per acre after a preplant soil incorporated treatment of Triflurex HFP. Apply Triflurex HFP as specified on its label at 1 to 2.5 pints per acre.

Precautions: Do not use the tank mix in AZ. Do not apply the tank mix of MANA Cotoran 4L + Triflurex HFP in liquid fertilizer. Refer to the Triflurex HFP label for direction, precautions, and limitations for use of that product, as well as the directions, precautions, and limitations on this MANA Cotoran 4L label.

Before applying MANA Cotoran 4L refer to **Application Soil Restrictions Table** in the PRECAUTIONS section for specific application rate soil restrictions.

Staple® LX:

A program of Staple LX at 1.3 – 2.1 fl. Oz / A plus MANA Cotoran 4L at 2 to 4 pints per acre applied preemergence followed by Staple LX early postemergence can be used for improved control of bristly starbur, coffee senna, common ragweed, Florida beggarweed, hemp sesbania, jimsonweed, ladythumb, smartweed, lambsquarter, annual morningglory (cypressvine, entireleaf, ivyleaf, pitted, purple, red/scarlet, sharppod / cotton, smallflower) Pennsylvania smartweed, pigweed (redroot, smooth, spiny), prickly sida, spotted spurge, spurred anoda, velvetleaf and for suppression of palmer pigweed. On light silt and sandy soils, low in organic matter, use the lower rate. Before applying MANA Cotoran 4L refer to **Application Soil Restrictions Table** in the PRECAUTIONS section for specific application rate soil restrictions.

Specific Weed Problems (Sickepod, Wild Poinsettia, Yellow Nutsedge): For improved control of the above weeds and other labeled weeds that often occur in high populations and/or have multiple seasonal flushes, a program of Staple LS at 1.7-2.1 oz/A plus MANA Cotoran 4L at 2-4 pints per acre applied preemergence followed by a postemergence application of Staple LX alone or in combination with MSMA or DSMA is recommended. On light silt and sandy soils, low in organic matter, use the lower rate.

Before applying MANA Cotoran 4L refer to **Application Soil Restrictions Table** in the PRECAUTIONS section for specific application rate soil restrictions.

Prowl®:

Apply MANA Cotoran 4L as a preemergence surface treatment as specified on this label at 2 to 4 pints per acre after a preplant treatment of Prowl for control of all weeds listed on both labels. Refer to the Prowl label for directions, precautions, and limitations for use of that product.

Before applying MANA Cotoran 4L refer to **Application Soil Restrictions Table** in the PRECAUTIONS section for specific application rate soil restrictions.

DSMA or MSMA:

MANA Cotoran 4L may be tank-mixed with DSMA or MSMA and applied as a directed spray to cotton from 3 inches high to first bloom. Use the broadcast rate of 2 to 4 pints of MANA Cotoran 4L per acre and the necessary amount of DSMA or MSMA plus 0.5% (1 pint / 25 gallons) of a surfactant that is approved by FDA for use on food and feed crops and is compatible with MANA Cotoran 4L + DSMA or MANA Cotoran 4L + MSMA when applied to cotton. When using this mixture, do not allow spray to contact cotton foliage. Do not graze treated areas or feed treated forage or gin trash to livestock. Adhere strictly to the dosage, precautions, and limitations of the labels of the products.

Before applying MANA Cotoran 4L refer to **Application Soil Restrictions Table** in the PRECAUTIONS section for specific application rate soil restrictions.

APPLICATION INSTRUCTIONS

Mixing Instructions

- (1) Fill spray tank ½ to ¾ full with clean water or nitrogen solution.
- (2) Start agitation.
- (3) Pour MANA Cotoran 4L directly from the container in to the partially filled spray tank.
- (4) If MANA Cotoran 4L is to be tank-mixed with another herbicide, add that product after the MANA Cotoran 4L is thoroughly mixed in the partially filled spray tank.
- (5) Add the rest of the water or nitrogen solution.
- (6) Provide agitation during mixing and application to maintain a uniform suspension.

To determine compatibility, add 2 teaspoons of MANA Cotoran 4L to 1 pint of of nitrogen solution (equals 4 pints in 25 gallons; if using other application rates, adjust compatibility test amounts proportionally). Stir or shake thoroughly. Let stand 5 minutes. If it remains mixed or can be remixed readily, the mixture is compatible and can be used. If MANA

Cotoran 4L settles out when mixed with a nitrogen solution, a suspensibility agent, such as Unite[®], Compex[®] or equivalent at 1 to 3 pints per 100 gallons of fertilizer may improve compatibility. Add 1/8 to 3/8 teaspoon of Unite to the compatibility test to determine its effectiveness.

Equipment

Thoroughly clean sprayer prior to use. Do not use a sprayer contaminated with 2,4-D or other materials, as crop damage or sprayer clogging may result.

All application must be properly maintained and calibrated using appropriate carriers or surrogates. Use properly calibrated conventional spray equipment that will apply spray uniformly and has hydraulic or mechanical agitation, except in Arizona, where only mechanical agitators are recommended. Screens in nozzles, in suction, and in-line strainers should be no finer than 50-mesh. Use a pump with capacity to maintain 35-40 psi at the nozzles. If hydraulic agitation is used, the pump should also provide sufficient agitation in the tank to keep the mixture in suspension. Avoid leaving spray mixture in tank without constant agitation. If bypass agitation is used, it should terminate at the bottom of the tank to minimize foaming. Wash sprayer thoroughly with clean water immediately after use.

Apply in a minimum of 10 gallons per acre for preplant incorporated or preemergence applications, or apply in a minimum of 20 gallons per acre for postemergence application.

Aerial Application

Apply MANA Cotoran 4L in a minimum of 5 gallons of water per acre. Avoid applications under conditions where uniform coverage cannot be contained or where extensive spray drift may occur. Flagmen and loaders should avoid inhalation of spray mist and prolonged contact with skin.

Band Treatment

For band treatment, calculate the amount of MANA Cotoran 4L needed by the formula:

$$\frac{\text{Band width (in)}}{\text{Row width (in)}} \times \text{broadcast rate per acre} = \text{amount needed per acre of field}$$

SPRAY DRIFT MANAGEMENT

A variety of factors including weather conditions (e.g., wind direction, wind speed, temperature, relative humidity) and method of application can influence pesticide drift. The applicator must evaluate all factors and make appropriate adjustments when applying this product.

Wind Speed: Do not apply at wind speeds greater than 15 mph.

Droplet Size: Apply as a medium or coarser spray (ASAE Standard 572).

Temperature Inversions: If applying at wind speeds less than 3 mph, the applicator must determine if a) conditions of temperature inversion exist, or b) stable atmospheric conditions exist at or below nozzle height. Do not make applications into areas of temperature inversions or stable atmospheric conditions.

Other State and Local Requirements: Applicators must follow all state and local pesticide drift requirements regarding application of fluometuron. Where states have more stringent regulations, they must be observed.

Equipment: All application equipment must be properly maintained and calibrated using appropriate carriers or surrogates.

Additional requirements for aerial applications:

1. The boom length must not exceed 75% of the wingspan or 90% of the rotor blade diameter.
2. Release spray at the lowest height consistent with efficacy and flight safety. Do not release spray at a height greater than 10 feet above the crop canopy unless a greater height is required for aircraft safety.
3. When applications are made with a crosswind, the swath must be displaced downwind. The applicator must compensate for this displacement at the up and downwind edge of the application area by adjusting the path of the aircraft upwind.

Additional requirement for groundboom application:

1. Do not apply with a nozzle height greater than 4 feet above the crop canopy.

9/9

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal.

PESTICIDE STORAGE: Keep container closed when not in use. Store product in original container only, away from food or feed.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL:

Nonrefillable Container (five gallons or less): Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then puncture and dispose of in a sanitary landfill, incinerate or burn, if allowed by state and local authorities. Stay out of smoke from burning container.

Nonrefillable Container (greater than five gallons): Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then puncture and dispose of in a sanitary landfill, incinerate or burn, if allowed by state and local authorities. Stay out of smoke from burning container.

Refillable Container: Refillable container. Refill this container with diazinon only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

FOR 24-HOUR EMERGENCY ASSISTANCE (SPILL, LEAK, OR FIRE), CALL INFOTRAC AT 1-800-535-5053.

LIMITATION OF WARRANTY AND LIABILITY

Read the entire directions for use, conditions of warranties and limitations of liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following **CONDITIONS, DISCLAIMER OF WARRANTIES and LIMITATIONS OF LIABILITY.**

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Makhteshim Agan of North America, Inc. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: To the extent consistent with applicable law, Makhteshim Agan of North America, Inc. makes no other warranties, express or implied, of merchantability or of fitness for a particular purpose or otherwise, that extend beyond the statements made on this label. No agent of Makhteshim Agan of North America, Inc. is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. To the extent consistent with applicable law, Makhteshim Agan of North America, Inc. disclaims any liability whatsoever for special, incidental or consequential damages resulting from the use or handling of this product.

LIMITATIONS OF LIABILITY: To the extent consistent with applicable law, the exclusive remedy of the user or buyer for any and all losses, injuries or damages resulting from the use or handling of this product, whether in contract, warranty, tort, negligence, strict liability or otherwise, shall not exceed the purchase price paid or at Makhteshim Agan of North America, Inc.'s election, the replacement of product.

Cotoran is a registered trademark of Agan Chemical Manufacturers, Ltd.

Command is a registered trademark of FMC Corporation

Compex is a registered trademark of KALO Agricultural Chemicals, Inc.

Clarity and Prowl are registered trademarks of BASF

Ignite is a registered trademark of Bayer

Parallel is a trademark of MANA for metolachlor

Staple is a registered trademark of E. I. duPont de Nemours & Co.

Triflurex is a registered trademark of Agan Chemical Manufacturers, Ltd. for trifluralin

Unite is a registered trademark of HACO, Inc.