

COTORAN[®] ACCU-PAK[®]

HERBICIDE

FOR PREEMERGENCE AND POSTEMERGENCE WEED CONTROL IN COTTON

ACTIVE INGREDIENT:	% BY WT.
Fluometuron: 1,1-dimethyl-3-(a,a,a-trifluoro-m-tolyl) urea	85.0%
INERT INGREDIENTS:	15.0%
	TOTAL 100.0%

Cotoran Accu-Pak is a water dispersible granule.

KEEP OUT OF REACH OF CHILDREN

WARNING-AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

This outer protective container contains Cotoran Accu-Pak in an inner water-soluble bag. Entire inner bag and contents dissolve in water. After opening outer container, immediately dump entire unopened inner bag into the partially filled sprayer or mix tank. Do not handle the soluble bag or expose it to moisture, since this may cause rupturing.

FIRST AID

IF IN EYES:

- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.

IF ON SKIN OR CLOTHING:

- Call a poison control center or doctor for treatment advice.
- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes.

IF SWALLOWED:

- Call a poison control center or doctor for treatment advice.
- Call a poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to by a poison control center or doctor.
- Do not give anything by mouth to an unconscious person.

IF INHALED:

- Move person to fresh air.
- If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.
- Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact PROSAR at 1-877-250-9291 for emergency medical treatment information.

NOTE TO PHYSICIAN: If Cotoran Accu-Pak is ingested, lavage stomach or induce vomiting. A slurry of activated charcoal in water can be left in the stomach. Give a saline laxative and supportive therapy.


NET CONTENTS: ___ POUND(S)

EPA Reg. No. 66222-34
EPA Est. No. 11603-IS-01



MAKHTESHIM
A G A N
NORTH AMERICA

Makhteshim-Agan of North America, Inc.
551 Fifth Avenue, Suite 1100
New York, NY 10176



**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
WARNING-AVISO**

Causes eye irritation. Do not get in eyes. Harmful if swallowed, inhaled, or absorbed through the skin. This product may cause skin rash after repeated exposure. Avoid contact with skin or clothing. Avoid breathing dust or spray mist.

ENVIRONMENTAL HAZARDS

This chemical is known to leach through soil into ground water under certain conditions as a result of registered uses. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground water contamination. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash waters.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks
- Protective eyewear

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROL STATEMENTS: When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

Exception: If the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Waterproof gloves
- Shoes plus socks
- Protective eyewear

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN CROP INJURY, POOR WEED CONTROL, AND/OR ILLEGAL RESIDUES.

CHEMIGATION STATEMENT

Do not apply this product through any type of irrigation system.

GENERAL INFORMATION

Cotoran Accu-Pak is effective for both annual grass and annual broadleaf weed control in cotton.

WEED CONTROL

Cotoran Accu-Pak applied according to directions will selectively control annual weeds in cotton, such as:

ANNUAL GRASS WEEDS

- | | |
|---------------|----------------------------------|
| Barnyardgrass | Foxtail |
| Crabgrass | Goosegrass |
| Crowfootgrass | Ryegrass |
| Fall panicum | Signalgrass (<i>Bracharia</i>) |

ANNUAL BROADLEAF WEEDS

- | | |
|-----------------------|------------------------------|
| Spurred anoda | Prickly sida (teaweed) |
| Cocklebur | Purslane |
| Florida pusley | Ragweed |
| Groundcherry (Wright) | Sesbania |
| Jimsonweed | Sicklepod |
| Lambsquarters | Smartweed |
| Morningglory | Russian thistle (Tumbleweed) |
| Pigweed | |

Cotoran Accu-Pak at rates up to 2.4 lbs. per acre in cotton is not effective for the control of established perennial weeds, such as johnsongrass from rhizomes.

**APPLICATION PROCEDURES
MIXING INSTRUCTIONS**

Prepare no more spray mixture than is needed for the immediate operation. Thoroughly clean spray equipment before using this product. **Vigorous agitation is necessary to maintain tank mix uniformity. Maintain a maximum of agitation throughout the spraying operation.** Flush spray tank thoroughly with water daily after use and dispose of pesticide rinsate by application to an already treated area.

Add 2/3 of the required amount of water to the spray or mixing tank. With the agitator running, drop the required number of **unopened** soluble bags of Cotoran Accu-Pak into the tank. If boron or fertilizer is to be added to the spray tank, do not add it until the water-soluble package is completely dissolved. Continue agitation while adding the remainder of water or nitrogen solution and during application to maintain a uniform suspension.

If Cotoran Accu-Pak settles out when mixed with a nitrogen solution, a suspensibility agent, such as Unite[®], Compex[®], or equivalent, at 1-3 pts. per 100 gals. fertilizer, may improve compatibility.

EQUIPMENT

Thoroughly clean sprayer prior to use. Do not use a sprayer contaminated with 2,4-D or other materials as crop damage or sprayer clogging may result.

Use properly calibrated conventional spray equipment that will apply spray uniformly and has hydraulic or mechanical agitation, except in Arizona where only mechanical agitators are recommended. Screens in nozzles, in suction, and in-line strainers should be no finer than 50-mesh. Use a pump with capacity to maintain 35-40 psi at the nozzles. If hydraulic agitation is used, the pump should also provide sufficient agitation in the tank to keep the mixture in suspension. Avoid leaving spray mixture in tank without constant agitation. If bypass agitation is used, it should terminate at the bottom of the tank to minimize foaming. Wash sprayer thoroughly with clean water immediately after use.

Apply in a minimum of 10 gals. per acre for preplant incorporated or preemergence applications, or apply in a minimum of 20 gals. per acre for postemergence application.

BAND TREATMENT

For band treatment, calculate the amount of Cotoran Accu-Pak needed by the formula:

<u>Band width in inches</u>	X	Broadcast rate per	=	Amount needed per
Row width in inches		acre		acre of field

IMPORTANT: Do not allow Cotoran Accu-Pak to become wet or overheated during storage as this may lead to the soluble bag rupturing or to chemical changes which will impair the herbicidal effectiveness of Cotoran Accu-Pak and which may also generate flammable vapors.

COTTON

Cotoran Accu-Pak may be applied as a preemergence treatment to the surface of the soil, or lightly incorporated immediately after planting, or as a directed, semi-directed, or over-the-top postemergence application. For preemergence treatment only, Cotoran Accu-Pak may be tank-mixed with most liquid nitrogen fertilizers. Cotoran Accu-Pak may be combined with other herbicides in cotton as directed in the **COMBINATIONS** section.

Cotoran Accu-Pak is most effective when used as follows:

- 1st Application:** Preemergence broadcast. See **PREEMERGENCE TREATMENT** below.
- 2nd Application:** Postemergence, directed, semi-directed, or over-the-top when cotton is 3-6 inches high. See **POSTEMERGENCE TREATMENT** below.
- 3rd Application:** If needed, postemergence, directed, or semi-directed, at lay-by (last normal cultivation), using no more than 1.2 lbs. per acre.

PREEMERGENCE TREATMENT

Apply Cotoran Accu-Pak at a rate of 1.2-2.4 lbs. per acre. On light silt and sandy soils low in organic matter, use the lower rate. Cotoran Accu-Pak may be applied to the surface of the soil or lightly incorporated into the soil after planting cotton.

PREPLANT INCORPORATED USE (ARIZONA AND NEW MEXICO)

Cotoran Accu-Pak may be applied preplant as a broadcast spray at 1.9 lbs. per acre after the beds for preplant irrigation have been formed. Apply before the beds are dragged off (using a bed shaper set to remove 1-2 inches from beds) in preparation for planting. Furrow irrigation should precede the application and planting should follow as soon as possible. Plant into untreated soil after drag-off and return treated soil to the beds when furrows are reformed. Do not allow furrow irrigation water to pond over the beds.

POSTEMERGENCE TREATMENTS

Cotoran Accu-Pak may be applied postemergence to cotton but preemergence to weeds, or postemergence to both cotton and weeds. Cotoran Accu-Pak should be applied at rates of 1.2-2.4 lbs. per acre. Apply as a directed, semi-directed, or over-the-top spray. Cotoran Accu-Pak may be applied to cotton plants at any stage of growth from 3 inches high to lay-by. Use the higher rate when applying postemergence to weeds; weeds should be 2 inches high or less. Use of a surfactant that is approved by FDA for use on food and feed crops and is compatible with Cotoran Accu-Pak when applied to cotton may enhance the postemergence herbicidal activity of Cotoran Accu-Pak. The surfactant rate should not exceed 0.5% of the carrier volume (1 pt. per 25 gals. of water).

PRECAUTION: Rainfall that is sufficient to germinate seeds or for growing crops is adequate to activate the herbicidal activity of Cotoran Accu-Pak; where dry weather conditions prevail, the herbicidal activity of Cotoran Accu-Pak may be delayed or reduced. Cultivate if weeds develop.

Do not make more than three applications of Cotoran Accu-Pak to the same crops or field in any one year. Do not plant crops other than cotton or sugarcane within 6 months of the last application of Cotoran Accu-Pak or injury may result.

The use of Cotoran Accu-Pak (any application method) in conjunction with a systemic insecticide at planting may result in injury to cotton. In addition, Cotoran Accu-Pak applied to cotton with any application of Command® (clomazone) herbicide may result in crop injury. Makhteshim-Agan will not be held responsible for losses or damage resulting from such uses.

West Texas only: Do not use Cotoran Accu-Pak on sandy, loamy sand, or fine sandy loam soils. Do not use Cotoran Accu-Pak on cotton planted in furrows.

Arkansas, Louisiana, and Mississippi only: Use 1.0 lb. of Cotoran Accu-Pak per acre on sandy loam soils low in organic matter.

Arizona and New Mexico: Cotton, corn, and grain sorghum can be planted the next spring. Do not plant treated areas to other crops until one year after last application. Do not use on sandy or coarse-textured soil of less than 1% organic matter.

Note: Do not apply within 60 days of harvest. Do not feed foliage from treated cotton plants or gin trash to livestock.

COMBINATIONS

DUAL® 8E: Use in a tank mixture with Cotoran Accu-Pak preemergence in cotton for control of all weeds as listed on both labels. This combination will also control spotted spurge, hyssop spurge, nodding spurge, and prostrate spurge. Apply to the soil surface at planting or after planting but before weeds or crop emerge using the appropriate rates from the table below.

Note: If incompatibility occurs, add X-77® at 0.5% volume/volume spray (4 pts./100 gals.) to help overcome this condition.

COTORAN ACCU-PAK + DUAL 8E- COTTON (AR, LA, MS, EASTERN OK, TN, BOOTHEEL OF MO, AND GULF COAST, RIO GRANDE VALLEY, AND EASTERN TX)

SOIL TEXTURE	BROADCAST RATES PER ACRE	
	COTORAN ACCU-PAK	DUAL 8E
Sand, loamy sand	DO NOT USE	
Sandy loam*	1.2 lbs.	1.25-1.5 pts.
Loam, Silt, Silt loam	1.2-1.9 lbs.	1.5-2 pts.
Silty clay loam, Sandy clay loam, Silty clay, Sandy clay, Clay loam, Clay	1.9-2.4 lbs.	2 pts.

*Use on sandy loam soils only in eastern OK and Gulf Coast, Rio Grande Valley, and eastern TX.

PRECAUTIONS:

1. Do not apply Cotoran Accu-Pak + Dual 8E on sand or loamy sand soils or in areas where water is likely to "pond" over the bed or crop injury may occur.
2. Do not apply broadcast to furrow-planted cotton.
3. The use of Cotoran Accu-Pak (any application method) in conjunction with a systemic insecticide at planting may result in injury to cotton. In addition, Cotoran Accu-Pak applied to cotton with any application of Command (clomozone) herbicide may result in crop injury. Makhteshim-Agan will not be held responsible for losses or damage resulting from such uses.
4. Do not use on Taloka silt loam or crop injury may occur.

Refer to the Dual 8E label for further instructions, precautions, and limitations.

Note: Do not feed treated forage or gin trash to livestock or graze treated areas.

Rotational Crops: Refer to the crop rotation instructions for Cotoran Accu-Pak alone on this label and for Dual 8E alone on the Dual 8E label.

TREFLAN® EC: Apply Cotoran Accu-Pak as a preplant incorporated or preemergence soil surface treatment as recommended on this label after a preplant soil incorporated treatment of Treflan EC. Refer to the Treflan EC label for directions, precautions, and limitations for use of that product, as well as the directions, precautions, and limitations on this Cotoran Accu-Pak label.

PROWL®: Apply Cotoran Accu-Pak as a preplant incorporated or preemergence soil surface treatment as recommended on this Cotoran Accu-Pak label after a preplant soil-incorporated treatment of Prowl. Refer to the Prowl label for directions, precautions, and limitations for use of that product.

ZORIAL RAPID 80: Apply Cotoran Accu-Pak according to directions on this Cotoran Accu-Pak label as a preemergence soil surface treatment following a preplant incorporated application of Zorial Rapid 80 or tank mix with Zorial Rapid 80 and apply as a preemergence treatment. Refer to the Zorial Rapid 80 label for directions for use, precautions, and limitations for use of that product.

DSMA OR MSMA: Cotoran Accu-Pak may be tank-mixed with DSMA or MSMA and applied as a directed spray to cotton from three inches high to first bloom. Use the broadcast rate of 1.2-2.4 lbs. Cotoran Accu-Pak per acre and the necessary amount of DSMA or MSMA plus 0.5% (1 pt. per 25 gals.) of a surfactant that is approved by FDA for use on food and feed crops and is compatible with Cotoran Accu-Pak + DSMA or Cotoran Accu-Pak + MSMA when applied to cotton. When using this mixture, do not allow spray to contact cotton foliage. Do not graze treated areas or feed treated forage or gin trash to livestock. Adhere strictly to the dosage, precautions, and limitations on the labels of the products.

STORAGE AND DISPOSAL

Store in a cool, dry place. Do not store this product under wet conditions. Handle outer container carefully when stored at temperatures less than 50°F to avoid breakage of inner soluble bag.

PESTICIDE DISPOSAL: Do not contaminate water, food, or feed by storage or disposal. Pesticide wastes may be toxic. Improper disposal of unused pesticide, spray mixture, or rinse water is a violation of federal law. If these wastes cannot be used according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance in proper disposal methods.

CONTAINER DISPOSAL: Do not reuse outer container. Dispose of empty outer container in a sanitary landfill, or by incineration, or by open burning, if allowed by state and local authorities. If burned, keep out of smoke.

For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup procedures and disposal of wastes. In the event of a major spill, fire, or other emergency, call INFOTRAC at 1-800-535-5053 day or night.

WARRANTY STATEMENT

MAKHTESHIM-AGAN OF NORTH AMERICA warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on such label only when used in accordance with directions under normal use conditions. It is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of MAKHTESHIM-AGAN OF NORTH AMERICA. In no case shall MAKHTESHIM-AGAN OF NORTH AMERICA be liable for consequential, special, or indirect damages resulting from the use or handling of this product. All such risks shall be assumed by the Buyer. In addition to the foregoing, no purchaser of this product (other than an end user) shall be entitled to any reimbursement for any loss suffered as a result of any suspension or cancellation of the registration for this product by the U.S. Environmental Protection Agency. Except as expressly provided herein, MAKHTESHIM-AGAN OF NORTH AMERICA makes no warranties, guarantees, or representations of any kind, either expressed or implied, or by usage of trade, statutory or otherwise, with regard to the product sold, including, but not limited to merchantability, fitness for a particular purpose, use or eligibility of the product for any particular trade usage. The exclusive remedy of any buyer or user of this product for any and all losses, injuries, or damages resulting from or in any way arising from the use, handling, or application of this product, whether in contract, warranty, tort, negligence, strict liability, or otherwise, shall be damages not exceeding the purchase price paid for this product or, at MAKHTESHIM-AGAN OF NORTH AMERICA's election, the replacement of this product.

Cotoran® is a registered trademark of Agan Chemical Manufacturers, Ltd.
Accu-Pak is a registered trademark.