







UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, D.C. 20460

OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

MAY 2 0 2010

Bert Volger Makhteshim Agan of North America, Inc. 4515 Falls of Neuse Rd., Suite 300 Raleigh, NC 27609

Dear Mr. Volger:

Subject:

Supplemental Label to RIMON® 0.83EC Insecticide for

Dry Bean Use in California EPA Registration No. 66222-35 Submission Date: May 13, 2010

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable. A stamped copy is enclosed for your records. This supplemental label will expire 18 months from the stamped date of February 12, 2010, on the RIMON® 0.83EC Insecticide master label. Please submit one (1) final printed copy for the above mentioned label. If you have any questions regarding this label, please contact Jennifer Gaines at (703) 305-5967 or via e-mail at gaines.jennifer@epa.gov.

Sincerely yours,

John Hebert

Product Manager (07)

Insecticide-Rodenticide Branch Registration Division (7505P)

SUPPLEMENTAL LABEL



RIMON 0.83EC Insecticide

EPA REG. NO. 66222-35

For Use on Beans (dry)

ACCEPTED MAY 2 0 2010

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide Registered under EPA Reg. No. (da222-35

This supplemental label expires on August 12, 2011 and must not be used or distributed after this date.

Active Ingredient: (% by weight)

Novaluron: 1-[3-chloro-4-(1,1,2-trifluoro-2-

*Contains 0.83 lbs. novaluron per gallon.

KEEP OUT OF REACH OF CHILDREN WARNING • AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.

(If you do not understand the label, find someone to explain it to you in detail.)

DIRECTIONS FOR USE Beans (dry)

INSECTS CONTROLLED BY RIMON 0.83 IN BEANS (DRY):

Target Pests	Rates (Fl. Ozs. / A)	Application Instructions
Armyworms Loopers Webworms	6 to 12	Apply when the majority of the target pest population is at egg hatch to early instars.
Bean leaf beetle Cucumber beetle Mexican bean beetle	9 to 12	Apply when the majority of the target pest population is at egg hatch to early instars.
Lygus	12	Apply when plant bugs appear and oviposition is initiated.
Thrips Whiteflies	12	Apply when the majority of the target pest population is at egg hatch to early instars. Do not apply more than two applications against whiteflies or thrips per season.

Use higher rates and higher spray volumes when larvae are large or foliage canopy is tall or dense.

Repeat applications as needed to protect new foliage growth, blooms and fruit, but not less than 7 days apart.

The use of surfactants/adjuvants (including non-ionic surfactants) on these crops is prohibited.

Do not apply more than 36 fl. oz. per acre per season.

Do not apply within 1 day of harvest.

THIS LABEL IS IN ADDITION TO THE STANDARD FEDERAL LABEL PRINTED ON THE CONTAINER

Follow all applicable directions, restrictions, and precautions on this supplemental label and the main EPA registered label. It is a violation of federal law to use this product in a manner inconsistent with its labeling. This label and the federal label for this product must be in the possession of the user at the time of pesticide application.

RIMON® is a Registered Trademark of Makhteshim Chemical Works, Ltd. Manufactured for:
Makhteshim Agan of North America, Inc.
4515 Falls of Neuse Rd., Suite 300
Raleigh, NC 27609
919-256-9300

^{*}U.S. Patent No. 5,142,064; 4,980,376; 5,089,525