

66222-35

9/5/2013

1/3

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, D.C. 20460



OFFICE OF CHEMICAL SAFETY  
AND POLLUTION PREVENTION

Mr. Jonathan Janis  
Makhteshim Agan of North America  
3120 Highwoods Blvd., Suite 100  
Raleigh, NC 27604

SEP 05 2013

Dear Mr. Janis:

Subject: Submission of supplemental label to add soybean and peanuts  
RIMON<sup>®</sup> 0.83EC Insecticide  
EPA Reg. No. 66222-35  
Your original submission dated: July 19, 2013

The labeling referred to above submitted in connection with the Federal Insecticide, Fungicide and Rodenticide Act, as amended is acceptable. A stamped copy is inclosed for your records. Please submit one (1) final printed copy for the above mentioned label before releasing the product for shipment. If you have any questions regarding this label, please contact Jennifer Gaines at (703) 305-5967 or [gaines.jennifer@epa.gov](mailto:gaines.jennifer@epa.gov).

Regards,

A handwritten signature in black ink, appearing to read "Reuben Baris".

Reuben Baris, Acting PM 07  
Insecticide-Rodenticide Branch  
Registration Division (7505P)

**ACCEPTED**  
**SEP 05 2013**  
 Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide Registered under EPA Reg. No. 66222-35

**Rimon 0.83EC Insecticide**  
 EPA REG. NO. 66222-35

**SUPPLEMENTAL LABELING**

This label expires on July 3, 2016 and must not be distributed or used after that date.

READ THE ENTIRE LABEL FOR RIMON 0.83EC INSECTICIDE BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label booklet for RIMON 0.83EC Insecticide and this supplement.

<b>ACTIVE INGREDIENT:</b>	<b>% BY WT.</b>
Novaluron:	
1-[3-chloro-4-(1,1,2-trifluoro-2-trifluoromethoxyethoxy)phenyl]-3-(2,6-difluorobenzoyl)urea*	9.3%
<b>OTHER INGREDIENTS:</b>	<b>90.7%</b>
<b>TOTAL:</b>	<b>100.0%</b>

\* Contains 0.83 lbs. Novaluron per gallon.

**USE ON PEANUTS AND SOYBEANS**

**KEEP OUT OF REACH OF CHILDREN**  
**WARNING - AVISO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.  
(If you do not understand the label, find someone to explain it to you in detail.)

**DIRECTIONS FOR USE**

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- The labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Rimon 0.83EC Insecticide before applying.
- Use of Rimon 0.83EC Insecticide according to this labeling is subject to the use precautions and limitations imposed by the label affixed to the container for Rimon 0.83EC Insecticide.

**PEANUTS\***

Target Pests	Rate (Fl. Oz / A)	Application Instructions
Green cloverworm Mexican bean beetle Velvet bean caterpillar	6-8	Make applications when larvae are small (< 0.5 inches) to give greater control and minimum insect damage to leaves. Repeat application if damaging numbers reappear. Use higher rates and higher spray volumes when the target pest population is 2X or more above state threshold level, or foliage canopy is tall, or dense and larvae are present in the lower part of the canopy, or if greater residual control is desired.
Amyworms, including: Beet armyworm Fall armyworm Southern armyworm Yellow-striped armyworm Lesser cornstalk borer Soybean looper (suppression) Thrips (suppression)	6-12	Apply at egg-hatch stage or when first signs of feeding occur. Use higher rates and higher spray volumes when larvae are more than 1/4 inch long, the target pest population is 2X or more above state threshold level, or foliage canopy is tall or dense and larvae are present in the lower part of the canopy. Repeat application if damaging numbers reappear to protect new growth.
Grasshoppers (nymphs only)	9-12	For best results, apply when the majority of infesting grasshoppers are in the early nymphal stages of development. RIMON 0.83EC Insecticide is not effective in controlling grasshoppers once they reach the adult stage. If a large influx from neighboring fields should occur, a tank mix with a knockdown insecticide may be necessary to reduce the population to minimize extensive foliage feeding.

**Peanuts continued**

Target Pests	Rate (Fl. Oz / A)	Application Instructions
<p><u>Aerial Application:</u> Apply in sufficient water (3 to 10 gallons per acre) to achieve uniform coverage of foliage.</p> <p><u>Ground Application:</u> Apply in 9 to 35 gallons of water per acre to give uniform coverage.</p> <p>Do not exceed 36 fl oz per acre per season.</p> <p>Reapplication on a 7 to 14 day interval may be required (refer to Grasshopper Application Instructions for more information).</p> <p>Do not harvest within 28 days of application.</p> <p>Do not feed treated peanut hay or vines to livestock.</p> <p>* Not registered for use in California.</p>		

**SOYBEANS\***

Target Pests	Rate (Fl. Oz / A)	Application Instructions
Green cloverworm Mexican bean beetle Saltmarsh caterpillar Velvet bean caterpillar	6-10	Make applications when larvae are small (< 0.5 inches) to give greater control and minimum insect damage to leaves. Repeat application if damaging numbers reappear. Use higher rates and higher spray volumes when the target pest population is 2X or more above state threshold level, or foliage canopy is tall, or dense and larvae are present in the lower part of the canopy, or if greater residual control is desired. RIMON 0.83EC Insecticide may be applied at the lower rate (6 fl. oz.) to prevent velvet bean caterpillar build-up when the vegetative growth of soybeans is completed and as pod formation begins. Consult local Extension Service regarding infestation levels requiring treatment.
Beet armyworm Cabbage looper Corn earworm Fall armyworm Soybean looper (suppression) Stinkbugs Tobacco budworm	6-12	Apply at egg-hatch stage or when first signs of feeding occur. Use higher rates and higher spray volumes when larvae are more than ¼ inch long, the target pest population is 2X or more above state threshold level, or foliage canopy is tall or dense and larvae are present in the lower part of the canopy. Repeat application if damaging numbers reappear to protect new growth.
Grasshoppers (nymphs only)	9-12	For best results, apply when the majority of infesting grasshoppers are in the early nymphal stages of development. RIMON 0.83EC Insecticide is not effective in controlling grasshoppers once they reach the adult stage. If a large influx from neighboring fields should occur, a tank mix with a knockdown insecticide may be necessary to reduce the population to minimize extensive foliage feeding.
<p><u>Aerial Application:</u> Apply in sufficient water (3 to 10 gallons per acre) to achieve uniform coverage of foliage.</p> <p><u>Ground Application:</u> Apply in 9 to 35 gallons of water per acre to give uniform coverage.</p> <p>Do not exceed 36 fl oz per acre per season.</p> <p>Reapplication on a 10 to 14 day interval may be required.</p> <p>Do not harvest within 30 days of application.</p> <p>Do not feed treated soybean forage to livestock.</p> <p>* Not registered for use in California.</p>		

Read the "LIMITATION OF WARRANTY AND LIABILITY" in the label booklet for RIMON 0.83EC Insecticide before using.

071813  
070313



3120 Highwoods Blvd, Suite 100, Raleigh NC 27604  
T: 866.406.6262 • F: 919.256.9308