

66222-42

10/22/2010

1/3



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF
CHEMICAL SAFETY AND
POLLUTION PREVENTION

Christie Hitchcock
Regulatory Specialist
Makhteshim Agan of North America, Inc.
4515 Falls of Neuse Road, Suite 300
Raleigh, NC 27609

OCT 22 2010

Subject: Bumper® 41.8EC
EPA Reg. No. 66222-42
Your amendment dated September 3, 2010
EPA Decision Number 440330

Dear Ms Hitchcock:

The supplemental label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act as amended is acceptable. These new uses must be added to the master label by amendment at the next printing or within 18 months, whichever comes first.

One copy of the label stamped "Accepted" is enclosed for your records. This label supersedes all labels previously accepted for this product. Please submit one copy of the final printed label before the product is released for shipment.

If you have any questions, please contact Robert Westin by phone at (703) 305-5721 or via email at westin.robert@epa.gov.

Sincerely,

A handwritten signature in black ink, which appears to read "Shaja Boyner", is written over the typed name.

Shaja Boyner
Product Manager (20)
Fungicide Branch
Registration Division (7504P)


Enclosure



Supplemental Labeling

BUMPER® 41.8 EC

(PROPICONAZOLE) FUNGICIDE
EPA Reg. No. 66222-42



SUPPLEMENTAL DIRECTIONS FOR USE ON CORIANDER (CILANTRO), GARDEN BEETS AND PARSLEY

DIRECTIONS FOR USE

- It is a violation of Federal law to use this product inconsistent with its labeling.
- This supplemental labeling must be in the possession of the user at the time of pesticide application.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.
- This new use can be found on the EPA stamped label for Bumper® 41.8 EC (66222-42) dated XX-XX-XX.

CROP USE DIRECTIONS AGRICULTURAL USES

CROP	PESTS CONTROLLED	RATE OF APPLICATION PER ACRE	APPLICATION INSTRUCTIONS
GARDEN BEETS	Leaf Spot (<i>Cereospora beticola</i>) Powdery Mildew (<i>Erysiphe polygoni</i>)	3-4 fl oz	Begin applications at first sign of disease and repeat on a 14 day interval. Make no more than 2 consecutive applications before rotating to another registered fungicide with a different mode of action. If disease levels continue to increase, immediately switch to a fungicide with a different mode of action. Applications are most effective when applied and allowed to dry before a rainfall. Use sufficient water volume to provide thorough coverage. Bumper® 41.8 can be applied by ground, air, or chemigation.
	1) Do not apply within 14 days of harvest 2) Do not apply more than 16 fl oz per acre per season. 3) Do not apply more than 0.45 lb a.i. propiconazole per acre per season.		
MINT	Rust (<i>Puccinia menthae</i>)	4 fl oz	Apply specified rate of Bumper 41.8 EC in a minimum of 20 gals of water per acre using ground application. Begin applications when plants are 2 to 4 inches high or when conditions become favorable for disease development. Make a second application 14 days after the first application.
	• Do not apply within 30 days of harvest. • Do not exceed 8 fl oz of Bumper 41.8EC per acre per season. • Do not apply more than 0.22 lb a.i. propiconazole per acre per season.		



M A N A

Supplemental Labeling

PARSLEY Fresh and Dried Leaves	Leaf Spot (<i>Cercospora</i> spp.)	3-4 fl oz	Begin applications at first sign of disease and repeat on a 14 day interval. Make no more than 2 consecutive applications before rotating to another registered fungicide with a different mode of action. If disease levels continue to increase, immediately switch to a fungicide with a different mode of action. Applications are most effective when applied and allowed to dry before a rainfall. Use sufficient water volume to provide thorough coverage. Bumper 41.8EC can be applied by air, ground, or chemigation equipment.
CILANTRO (Coriander), Leaves	Leaf Spot (<i>Alternaria</i> spp.)		
	Powdery Mildew (<i>Erysiphe</i> spp.)		
	<ul style="list-style-type: none"> Do not apply more than 16 fl oz per acre per season of Bumper 41.8EC. Do not apply more than 0.45 lb a.i. propiconazole per acre per season. Do not apply within 14 days of harvest. 		

Bumper is a registered trademark of Makhteshim Agan of North America, Inc.

Bumper 41.8 EC (66222-42)(to EPA 09-03-10)