



ACCEPTED

DEC - 5 2003

Under the Foderal Insecticide, Functoide, and Redenticide Act, as amended, for the pesticide registered under RPA Res. No. 26222-63

SUPPLEMENTAL LABELING THIONEX® 3EC

INSECTICIDE

For Agricultural or Commercial Use Only

Not for use or storage in or around residential sites-see DIRECTIONS FOR USE/GENERAL

INSTRUCTIONS for prohibited areas

EPA REG. NO. 66222-63

*Contains Xylene Range Aromatic Solvent.

Contains Aylene Range Afornatic Solvent.

Contains 3 pounds of endosulfan per gallon.

KEEP OUT OF REACH OF CHILDREN



SEE OTHER PANELS FOR ADDITIONAL PRECAUTIONARY INFORMATION.

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Do not apply this product through any type of irrigation equipment.

COMMERCIALLY GROWN ORNAMENTALS Grown Outdoors in Nurseries ONLY

Ornamental Trees and Shrubs

	Officiality	IIICI IICC	is and onlines
CROP	AP	TE OF PLICA- FION	METHOD OF APPLICATION
LEATHER- LEAF FERN (LEATHER HOLLY FERN) (Except California)	10 tea	qt. per 0 gals. (1¼ spoons er gal.)	Begin treatment when first larval feeding is observed in the mid-vein area at the base of the leaflets. Repeat at intervals of 2 to 3 weeks as necessary.
	 Food utensils such as teaspoons must not be used for food purposes after use in measuring pesticides. Do not exceed a maximum of 3 lbs. active ingredient (i.e., 4 qts.) per acre per year. Do not use in the greenhouse. 		