

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

WASHINGTON, D.C. 20460

JUL 2 0 2006

OFFICE OF PREVENTION, PESTICIDES AND TOXIC SUBSTANCES

Ms. Jane Rothwell Registration Specialist Makhteshim Agan of North America, Inc. 4514 Falls of Neuse Rd., Suite 300 Raleigh,NC 27609

Dear Ms. Rothwell:

SUBJECT: Prodiamine 65 WG Herbicide

EPA Registration No. 66222-89

Application and Letter Dated July 10, 2006, Request To Amend Registration by the Addition of Supplemental Labeling for Use Sites: Irrigation, Drainage Ditches,

Spreading Grounds, Channels, Canals and Similar Sites; and on

Levees of Wastewater Treatment Facilities

Your application to amend the registration of the subject pesticide product by adding supplemental labeling for the use sites: irrigation, drainage ditches, spreading grounds, channels, canals and similar sites; and on levees of wastewater treatment facilities has been reviewed and found acceptable under the federal Insecticide, Fungicide and Rodenticide Act, as amended (FIFRA), provided that you:

- 1. Submit one (1) copy of the final printed supplement labeling prior to your shipment under the enclosed stamped labeling.
- 2. Incorporate the subject supplemental labeling into your master label for Prodiamine 65 WG Herbicide at its next printing.

If these conditions are not complied with, this registration will be subject to cancellation in accordance with FIFRA, section 6(e). Your release for shipment under this labeling constitutes acceptance of these conditions. A stamped copy of supplemental labeling is enclosed for your records.

Sincerely yours,

Joanne I. Miller Product Manager (23) Herbicide Branch

Registration Division (7505P)

Enclosure

JUL 20 2006

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.

SUPPLEMENTAL LABELING

PRODIAMINE 65 WG

EPA Reg. No. 66222-89

SUPPLEMENTAL LABELING FOR CONTROL OF CERTAIN WEEDS ON DRAINAGE DITCHES CALIFORNIA AND ARIZONA ONLY

ACTIVE INGREDIENT: % BY WT. Prodiamine (CAS No. 29091-21-2) 65.0% INERT INGREDIENTS: 35.0%

KEEP OUT OF REACH OF CHILDREN CAUTION

IF IN EYES:

· Hold eye open and rinse slowly and gently with water for 15-20 minutes.

· Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.

FIRST AID

Call a poison control center or doctor for treatment advice.

IF ON SKIN OR

CLOTHING:

Take off contaminated clothing.

• Rinse skin immediately with plenty of water for 15 - 20 minutes.

· Call a poison control center or doctor for treatment advice.

IF INHALED:

· Move person to fresh air.

 If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.

 Call a poison control center or doctor for further treatment advice. Call a poison control center or doctor immediately for treatment advice.

SWALLOWED:

Have person sip a glass of water if able to swallow.

Do not induce vomiting unless told to do so by a poison control center or doctor.

Do not give anything by mouth to an unconscious person.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. FOR 24-HOUR CHEMICAL EMERGENCY INFORMATION (SPILL, LEAK, FIRE, OR EXPOSURE), CALL INFOTRAC at 1-800-535-5053.

FOR 24-HOUR EMERGENCY MEDICAL INFORMATION, CALL PROSAR at 1-877-250-9291.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if inhaled or absorbed through the skin. Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

DIRECTIONS FOR USE

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR PEST CONTROL, CROP INJURY, OR ILLEGAL RESIDUES.

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

- This supplemental labeling must be in the possession of the user at the time of pesticide application.
- This supplemental label contains new or supplemental instructions for use of this product which do not appear on the EPA-registered package label. Follow all instructions on this label and the label on the container carefully.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on this label and the EPA registered label attached to the container.

Page 1 of 2

Makhteshim Agan of North America, Inc. 4515 Falls of Neuse Road, Suite 300, Raleigh NC 27609 T: 919.256.9305 • F: 919.256.9308

FOR CONTROL OF CERTAIN WEEDS ON DRAINAGE DITCHES CALIFORNIA AND ARIZONA ONLY

Irrigation, Drainage Ditches, Spreading Grounds, Channels, Canals and Similar Sites:

Apply 1.0 to 2.3 lbs. per acre for preemergent control of shallow germinating annual grasses and small seeded broadleaf weeds. Prodiamine 65 WG inhibits growth and development of newly germinated weed seeds. It will not control established vegetation. Higher rates of application provide for a longer control period. Sequential applications are allowed so long as the total amount of product applied does not exceed the maximum annual application of 2.3 lbs. per acre. Apply only when water is not present. For irrigation ditches, apply when ditch is not in use. To minimize movement of Prodiamine 65 WG with irrigation water and to activate product, it is essential that the herbicide be fixed in the soil by moisture. Apply before expected seasonal rainfall (if possible when soil in the ditch is still moist). Following treatment, if rainfall has not totaled at least one-half inch within 14 to 21 days, fill ditch with water and allow to stand for 48 hours; drain off any remaining water before using ditch.

Wastewater Treatment Facilities (Levees)

Apply 1.0 to 2.3 lbs. per acre for preemergent control of shallow germinating annual grasses and small seeded broadleaf weeds. Use the higher rate of application for longer control periods. Sequential applications are allowed so long as the total amount of product applied does not exceed the maximum annual application of 2.3 lbs. per acre. Applications may be made down to the water level. Do not apply to the water. Care must be taken not to allow overspray of the water or drift into the water.