

66222-99

2/26/2009

1 of 20



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

FEB 26 2009

Mrs. Jane Rothwell
Registration Specialist
Makhteshim Agan of North America, Inc.
4515 Falls of Neuse Rd., Suite 300
Raleigh, NC 27609

RE: Notification for Alternate Marketing Language specific to the State of California
EPA Registration Number: 66222-99
Date of Submission: January 10, 2009

Dear Mrs. Rothwell:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 dated January 10, 2009, for the product Fanfare 2EC. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the actions requested fall within the scope of PRN 98-10. The label submitted with the application has been stamped "Notification" and will be placed in our records.

If you have any questions, please me directly at 703-305-6249 or Joyce Edwards of my staff at 703-308-5479.

Sincerely,

A handwritten signature in black ink, appearing to read "Linda", is located below the word "Sincerely,".

Linda Arrington
Notifications & Minor Formulations Team Leader
Registration Division (7505P)
Office of Pesticide Programs

20920

Please read instructions on reverse before completing form.

Form Approved, OMB No. 2070-0080, Approval expires 2-28-95



United States
Environmental Protection Agency
Washington, DC 20460

☐ Registration
☐ Amendment
☒ Other

OPP Identifier Number

Application for Pesticide - Section I

1. Company/Product Number 66222-99	2. EPA Product Manager Bewander Alexander	3. Proposed Classification <input type="checkbox"/> None <input checked="" type="checkbox"/> Restricted
4. Company/Product (Name) Fanfare 2EC	PM# 13	
5. Name and Address of Applicant (Include ZIP Code) Makhteshim Agan of North America, Inc. 4515 Falls of Neuse Rd., Suite 300 Raleigh, NC 27609 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ FEB 26 2009 Product Name _____	

Section - II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

Notification to EPA for alternate marketing language on label in accordance with PR Notice 98-10. This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula for this product. I understand that it is a violation of 18 USC Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under Sections 12 and 14 of FIFRA.

Section - III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No		<input type="checkbox"/> Metal <input type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____	
* Certification must be submitted		If "Yes" Unit Packaging wgt.	No. per container	If "Yes" Package wgt.	No. per container
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container		5. Location of Label Directions <input type="checkbox"/>	
6. Manner in Which Label is Affixed to Product <input type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled		<input type="checkbox"/> Other _____			

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name Jane Rothwell	Title Registration Specialist	Telephone No. (Include Area Code) 901-861-4400
2. Signature 		FEDERAL APPLICATION Received (Stamped)
3. Title Registration Specialist		
4. Typed Name Jane Rothwell		
5. Date 1-10-09		



January 10, 2009

Document Processing Desk (NOTIF)
 Registration Division (7505P)
 Office of Pesticide Programs
 U.S. Environmental Protection Agency
 Ariel Rios Building
 1200 Pennsylvania Ave, NW
 Washington DC 20460

ATTN: Bewanda Alexander for George LaRocca (13)
 Phone: 703-305-7460

Re: Fanfare 2EC, EPA Reg. No. 66222-99
 Notification of additional marketing language in accordance with PR Notice 98-10 §N

Dear Ms. Alexander:

As allowed in PR Notice 98-10, Maktheshim Agan of North America Inc. is submitting a notification to add alternate marketing language specific to the state of California. We are filing the notification as we need it in order to register the product in California and other states.

In support of this submission, the following documents are attached:

- Application for Pesticide Registration (8570-1)
- One copy of final printed labeling (one annotated)

This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula for this product. I understand that it is a violation of 18 USC Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under Sections 12 and 14 of FIFRA.

Thank you for your help with this request. Please contact me should you have any questions at (901)-861-4400 or by email at janer@manainc.com.

Sincerely,

Jane Rothwell
 Registration Specialist

Enclosures

40 of 20

RESTRICTED USE PESTICIDE

Toxic to fish and aquatic organisms

For retail sale to and use only by certified applicators or persons under their direct supervision and only for the uses covered by the certified applicator's certification.

FANFARE[®] 2EC
INSECTICIDE/MITICIDE

Contains bifenthrin, the active ingredient used in Capture 2EC.
Fanfare 2EC is not manufactured or distributed by FMC Corporation.

ACTIVE INGREDIENT:

% BY WT.

Bifenthrin: (2 methyl[1,1'-biphenyl]-3-yl)methyl 3-(2-chloro-3,3,3-trifluoro-1-propenyl)-2,2-dimethyl-cyclopropanecarboxylate* 25.1%

OTHER INGREDIENTS:** 74.9%

TOTAL 100.0%

*Cis isomers 97% minimum, trans isomers 3% maximum.

**Contains petroleum distillates.

This product contains 2 pounds active ingredient per gallon.

EPA Reg. No. 66222-99

EPA Est. No. **7429-CA-002, 5905-CA-001, 11678-IS-001, 11678-ISR-002**

Manufactured for:

NOTIFICATION Makhteshim Agan of North America, Inc.
4515 Falls of Neuse Rd., Suite 300
Raleigh, NC 27609

FEB 26 2009

NET CONTENTS: _____ Gallons

KEEP OUT OF REACH OF CHILDREN
WARNING-AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail).

FIRST AID

- | | |
|--------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| IF SWALLOWED: | <ul style="list-style-type: none">• Immediately call a poison control center or doctor.• Do not induce vomiting unless told to do so by a poison control center or doctor.• Do not give any liquid to the person.• Do not give anything by mouth to an unconscious person. |
| IF IN EYES: | <ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice. |
| IF ON SKIN OR CLOTHING: | <ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice. |
| IF INHALED: | <ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth if possible.• Call a poison control center or doctor for further treatment advice. |

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-877-250-9291 for emergency medical treatment information.

NOTE TO PHYSICIAN: This product is a pyrethroid. If large amounts have been ingested, the stomach and intestines should be evacuated. Treatment is symptomatic and supportive. Digestible fats, oils, or alcohol may increase absorption and should be avoided. This product contains a petroleum distillate; vomiting may cause aspiration pneumonia.

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
WARNING

May be fatal if swallowed. Causes substantial but temporary eye injury. Do not get in eyes or on clothing. Wear protective eyewear (goggles, face shield, or safety glasses.) Harmful if absorbed through skin. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category E on an EPA chemical resistance category selection chart.

Handlers who may be exposed to the dilute through application or other tasks must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves, such as barrier laminate, nitrile rubber, neoprene rubber, or Viton
- Shoes plus socks

Handlers who may be exposed to the concentrate through mixing, loading, application, or other tasks must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves, such as barrier laminate, nitrile rubber, neoprene rubber, or Viton
- Shoes plus socks
- Protective eyewear

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to fish and aquatic invertebrates. Use with care when applying in areas adjacent to any body of water. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not make applications when weather conditions favor drift from treated areas. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwaters.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are actively visiting the treatment area.

The use of bifenthrin is prohibited in areas that may result in exposure of endangered species to bifenthrin. Prior to use in a particular county contact the local extension service for procedures and precautions to use to protect endangered species.

PHYSICAL/CHEMICAL HAZARDS

COMBUSTIBLE. Do not use or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves, such as barrier laminate, nitrile rubber, neoprene rubber, or Viton
- Shoes plus socks

CHEMIGATION USE DIRECTIONS

Apply this product only through sprinkler including center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set, or hand move irrigation systems. Do not apply this product through any other type of irrigation system.

Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system.

Crop injury, lack of effectiveness, or illegal residues in the crop can result from non-uniform distribution of treated water. Contact your State Agricultural Extension Service specialists, equipment manufacturers, or other experts for consultation on the suitability of the equipment set up to obtain effective control of the target insect pests.

A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise. Failure to cease application during a mechanical stoppage may result in undesirable residues to adjacent areas.

The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.

60920

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.

The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Do not apply when wind speed favors drift beyond the area intended for treatment.

For sprinkler irrigation, meter Fanfare 2EC at a continuous uniform rate during the entire irrigation period. To ensure accurate application over the treated area, apply in sufficient volume of water or other diluent. If non-emulsified oil is used as the diluent, use 1 to 2 pints per acre. Continuous agitation of the pesticide supply tank for the duration of the application period is recommended. When chemigation systems are used, 0.5 inch per acre of irrigation water is suggested except that for Low Energy Precision Application (LEPA) irrigation, a minimum of 0.75 inch of water per acre is suggested.

RESISTANCE MANAGEMENT

Fanfare 2EC contains a Group 3 Insecticide. With repeated use of Group 3 insecticides as the primary method of control in the same field or in successive years, insect/mite populations can develop resistant biotypes. If this occurs, insect/mite biotypes with acquired resistance to Group 3 insecticides may eventually dominate the insect/mite population. This may result in partial or total loss of control of those species by Fanfare 2EC or other Group 3 insecticides.

To delay development of insecticide resistance, the following practices are recommended:

- Base insecticide applications on comprehensive IPM programs. This program should include an insect management program that includes cultural and biological control where possible.
- Use good resistance management strategies established for the use area. This may include the use of insecticide rotations or tank mixes with other Groups of insecticide and miticides in an IPM program.
- Always apply Fanfare 2EC at the recommended rates and according to label directions. Do not use less than recommended label rates alone or in tank mixtures unless directed otherwise in supplemental labeling supplied by Makhteshim Agon of North America, Inc. (MANA).
- Monitor treated populations in the field for loss of control. If poor performance cannot be attributed to improper application or extreme weather conditions, a resistant strain of insect may be present. Immediately consult your local MANA representative or agricultural advisor for the best alternative method of control for your area.
- Do not treat seedling plants grown for transplant in greenhouse, shade houses, or field plots.
- Consult your local extension specialist, certified crop advisor, and/or manufacturer for insecticide resistance management and/or IPM recommendations for the specific site and resistant pest problems.

ROTATIONAL CROPS

Crops for which bifenthrin tolerances exist may be rotated at any time. All other crops may be rotated 30 days following the final application of Fanfare 2EC.

MIXING INSTRUCTIONS

The spray tank must be clean, thoroughly rinsed, and decontaminated before adding either Fanfare 2EC alone or with tank mix combinations (see **Fanfare 2EC in Tank Mixtures** section below). If water is used as the carrier, use clean water.

For aerial applications made on brassicas (see **CROPS** section of the label below for full list of approved brassicas), canola, crambe, rapeseed, foliar applications on corn, cucurbits (see **CROPS** section of the label below for full list of approved cucurbits), eggplant, grapes, head lettuce, and succulent peas and beans (see **CROPS** section of the label below for full list of approved succulent peas and beans), 1 to 2 quarts of emulsified oil may be substituted for 1 to 2 quarts of water in the finished spray. For aerial applications made on cotton, 1 quart of emulsified oil may be substituted for one quart of water in the finished spray. Thorough coverage is essential to achieve control.

Fanfare 2EC Used Alone: When Fanfare 2EC is used alone, add the recommended amount to the spray tank when the tank is half filled with water or other carrier, then add the rest of the water or other carrier (as permitted on this label). Provide sufficient agitation during mixing and application to maintain a uniform emulsion.

Fanfare 2EC with Fertilizer: Fill the spray tank approximately one half full with water and/or liquid fertilizer, add the proper amount of Fanfare 2EC, then add the rest of the water and/or fertilizer. Provide sufficient agitation during mixing and application to maintain a uniform spray mixture.

A jar compatibility test should be performed with the appropriate ratio of Fanfare 2EC and fertilizer to ensure the mixture will stay in solution. Maintain constant agitation during mixing and application.

Fanfare 2EC in Tank Mixtures: If a tank mixture is used, it is recommended that a compatibility test be done before actual tank mixing. A jar test for physical compatibility is recommended for untried mixtures using proper ratios and mixing sequences of all ingredients to be included in the mixture. Once compatibility is confirmed for the tank mix, fill the tank half full with water or other carrier. Start and continue agitation throughout mixing following conventional mixing order practices. Fanfare 2EC may be applied in tank mixtures with other products approved for use on registered crops. Observe all restrictions and precautions which appear on the labels of these products.

California Closed System: Special Equipment: The registration of Fanfare 2EC in California requires that the product be used in a closed system that meets the criteria for closed systems as established by the California Department of Pesticide Regulation (CDPR). The criteria and a list of suppliers meeting the criteria are available from CDPR.

APPLICATION INSTRUCTIONS AND RECOMMENDATIONS

The rate of Fanfare 2EC applied will vary according to pest pressure and timing of application. Use lower rates under light to moderate infestations and higher rates under heavy insect pressure and for mite control. Arid climates generally require higher rates.

Unless otherwise specified for a specific crop, apply when pest population reaches economic (damaging) threshold and repeat as necessary to maintain control. Thorough coverage is essential to achieve control.

In the **COMMENTS** section of the label for each crop, the recommended application rate when applied by ground and/or air is listed as an amount of spray per acre. In all cases, this refers to finished spray per acre.

SPRAY DRIFT REQUIREMENTS

Wind Direction and Speed

Only apply this product if the wind direction favors on-target deposition. Do not apply when the wind velocity exceeds 15 mph.

Temperature Inversion

Do not make aerial or ground applications into temperature inversions. Inversions are characterized by stable air and increasing temperatures with height above the ground. Mist or fog may indicate the presence of an inversion in humid areas. The applicator may detect the presence of an inversion by producing smoke and observing a smoke layer near the ground surface.

Droplet Size

Use only Medium or coarser spray nozzles (for ground and non-ULV aerial application) according to ASAE (S572) definition for standard nozzles. In conditions of low humidity and high temperatures, applicators should use a coarser droplet size.

Additional Requirements for Ground Applications

Wind speed must be measured adjacent to the application site on the upwind side, immediately prior to application.

For ground boom applications, apply using a nozzle height of no more than 4 feet above the ground or crop canopy.

For airblast applications, turn off outward pointing nozzles at row ends and when spraying the outer two rows. To minimize spray loss over the top in orchard applications, spray must be directed into the canopy.

Additional Requirements for Aerial Applications

The spray boom should be mounted on the aircraft as to minimize drift caused by wingtip or rotor vortices. The minimum practical boom length should be used and must not exceed 75% of the wing span or rotor diameter.

Flight speed and nozzle orientation must be considered in determining droplet size.

Spray must be released at the lowest height consistent with pest control and flight safety. Do not release spray at a height greater than 10 feet above the crop canopy unless a greater height is required for aircraft safety.

When applications are made with a cross-wind, the swath will be displaced downwind. The applicator must compensate for this displacement at the downwind edge of the application area by adjusting the path of the aircraft upwind.

BUFFER ZONES

Vegetative Buffer Strip

Construct and maintain a minimum 10-foot-wide vegetative filter strip of grass or other permanent vegetation between the field edge and down gradient aquatic habitat (such as, but not limited to, lakes; reservoirs; rivers; permanent streams; marshes or natural ponds; estuaries; and commercial fish farm ponds).

Only apply products containing (*name of pyrethroid*) onto fields where a maintained vegetative buffer strip of at least 10 feet exists between the field and down gradient aquatic habitat.

For guidance, refer to the following publication for information on constructing and maintaining effective buffers:

Conservation Buffers to Reduce Pesticide Losses. Natural Resources Conservation Services. USDA, NRCS. 2000. Fort Worth, Texas. 21 pp.

<http://www.in.csusda/v/technical/agronom/newconbuf.pdf>

In New York State, this product may not be applied within 100 feet (using ground equipment) to 300 feet (using aerial equipment) of coastal marshes or streams that drain into coastal marshes.

Buffer Zone for Ground Application (groundboom, overhead chemigation, or airblast)

Do not apply within 25 feet of aquatic habitats (such as, but not limited to, lakes, reservoirs, rivers, streams, marshes, natural ponds, estuaries, and commercial fish ponds).

Buffer Zone for ULV Aerial Application

Do not apply within 450 feet of aquatic habitats (such as, but not limited to, lakes, reservoirs, rivers, streams, marshes, natural ponds, estuaries, and commercial fish ponds).

Buffer Zone for Non-ULV Aerial Application

Do not apply within 150 feet of aquatic habitats (such as, but not limited to, lakes, reservoirs, rivers, streams, marshes, natural ponds, estuaries, and commercial fish ponds).

PREHARVEST INTERVAL

The required days to wait between the last application and harvest are given in () after each crop name.

**CROPS
ARTICHOKE (5)**

PEST	DOSAGE		COMMENTS
	LB AI/A	FL OZ/A	
Artichoke Plume Moth, Cribrate Weevil	0.10	6.4	Ground application: Apply in water in a minimum of 75 gallons per acre as a full cover spray. Air application: Apply in water in a minimum of 10 gallons per acre.
<ul style="list-style-type: none"> Do not apply more than 0.5 lb. active ingredient (32 ounces formulated) per acre per season. Repeat applications if needed to maintain control, but do not make applications less than 15 days apart. 			

BRASSICAS (7)

Head and Stem Brassica Vegetables including Broccoli, Chinese Broccoli (gailon, white flowering broccoli), Brussels sprouts, Cauliflower, Cavalo broccoli, Kohlrabi, Cabbage, Chinese Cabbage (napa), Chinese Mustard Cabbage (gai choy)

Cavalo broccolo, Kohlrabi, Cabbage, Chinese Cabbage (napa), Chinese mustard Cabbage (gai choy)			
PEST	DOSAGE		COMMENTS
	LB AI/A	FL OZ/A	
Aphids, Armyworms, Corn Earworm, Crickets, Cucumber Beetle, Cutworms, Diamondback Moth, Flea Beetle, Ground Beetles, Imported Cabbageworm, Leafhoppers, Loopers, Saltmarsh Caterpillar, Stink Bugs, Thrips, Tobacco Budworm, \ Whitefly, Wireworm (adults)	0.033-0.10	2.1-6.4	Ground application: Apply in water in a minimum of 10 gallons per acre. Air application: Apply in water in a minimum of 2 gallons per acre. Emulsified oil may be substituted for water. See section entitled MIXING INSTRUCTIONS for details on the amount of oil to use in the spray tank in lieu of water.
Banks Grass Mite, Carmine Mite, Lygus spp., Pacific Spider Mite, Twospotted Spider Mite	0.08-0.10	5.12-6.4	
<ul style="list-style-type: none">Do not apply more than 0.5 lb. active ingredient (32 ounces formulated) per acre per season.Do not make more than 5 applications after bloom.Repeat applications if needed to maintain control, but do not make applications less than 7 days apart.			

CANEBERRIES (3)

Caneberries including Blackberries, Bingleberries, Dewberries, Loganberries, Lowberries, Marionberries, Olallieberries, Raspberries, Youngberries

PEST	DOSAGE		COMMENTS
	LB AI/A	FL OZ/A	
Leafrollers, Orange Tortrix, Root Weevils	0.05-0.10	3.2-6.4	Ground application: Apply in water in a minimum of 50 gallons per acre. Air application: Apply in water in a minimum of 10 gallons per acre. A total of two applications may be made. The first should be made pre-bloom and the second should be made post-bloom.
Spider Mites	0.10	6.4	
<ul style="list-style-type: none">Do not apply more than 0.2 lb. active ingredient (12.8 ounces formulated) per acre per season.			

CANOLA, CRAMBE, RAPESEED (35)

PEST	DOSAGE		COMMENTS
	LB AI/A	FL OZ/A	
Aphids, Armyworms, Cutworms, Diamondback Moth, Flea Beetles, Flea Hoppers, Grasshoppers, Loopers, Lygus Bugs, Other Lepidopterous Larvae, Plant Bugs, Seedpod Weevil, Stink Bugs, Thrips, Whitefly	0.033-0.04	2.1-2.6	Ground application: Apply in water in a minimum of 10 gallons per acre. Air application: Apply in water in a minimum of 2 gallons per acre. Emulsified oil may be substituted for water. See section entitled MIXING INSTRUCTIONS for details on the amount of oil to use in the spray tank in lieu of water.
<ul style="list-style-type: none"> Do not apply more than 0.08 lb. active ingredient (5.12 ounces formulated) per acre per season. Repeat applications if needed to maintain control, but do not make applications less than 14 days apart. 			

90220

CHRISTMAS TREES (For Use Only in Washington and Oregon)

PEST	DOSAGE		COMMENTS
	LB/AI/A	FL. OZ/A	
Root Weevil, Spruce Spider Mite	0.06-0.1	3.9-6.4	<p>Ground application: Apply in water in a minimum of 20 gallons per acre.</p> <p>Air application: Apply in water in a minimum of 5 gallons per acre.</p> <p>Fanfare is generally not phytotoxic to Christmas trees. However, applications should be made to a small representative group of plants to ensure that a particular variety grown under current conditions is not unusually sensitive to Fanfare 2EC.</p> <p>Maintain a minimum of 21 days between applications.</p>
<ul style="list-style-type: none"> Do not apply more than 0.1 lb. active ingredient (6.4 ounces formulated) per acre per treatment. Do not make more than 3 applications in a crop year. Do not make applications through irrigation systems. 			

CILANTRO, CORIANDER (3)

PEST	DOSAGE		COMMENTS
	LB AI/A	FL OZ/A	
Aphids, Beet Armyworm, Cabbage Looper, Cutworm, Flea Beetle, Grasshoppers, Leafminer, Saltmarsh Caterpillar, Spotted Cucumber Beetle, Thrips, Whitefly	0.033-0.10	2.1-6.4	<p>Ground application: Apply in water in a minimum of 10 gallons per acre.</p> <p>Air application: Apply in water in a minimum of 2 gallons per acre.</p> <p>Apply in sufficient water to obtain thorough coverage.</p>
Two Spotted Spider Mite	0.08-0.10	5.12-6.4	
<ul style="list-style-type: none"> Do not apply more than 0.5 lb. active ingredient (32 ounces formulated) per acre per season. Do not make applications less than 7 days apart. 			

CITRUS (Except Florida)(1)*

PEST	DOSAGE		COMMENTS
	LB AI/A	FL OZ/A	
Asian Cockroach, Diaprepes Root Weevil (<i>Diaprepes abbreviatus</i>), Fire ants	0.25-0.50	16-32	<p>Ground application: Apply in water in a minimum of 30 gallons per acre.</p> <p>Use a hand-gun or shielded sprayer to apply to individual citrus trees if they are not planted in solid rows.</p> <p>Diaprepes root weevil emergence generally occurs in the spring, but weather conditions can prompt a second emergence in the fall. In areas where only a spring emergence is expected, use 32 ounces of Fanfare 2EC. In areas where a second emergence is expected, use 16 ounces of Fanfare 2EC in the early season and 16 ounces of Fanfare 2EC later in the season.</p> <p>If the length of control of Fanfare 2EC is not sufficient to cover the emergence of the root weevil, use other pest control measures as recommended by State Agricultural Extension Specialists or other local experts.</p> <p>*Use in California not permitted unless accompanied by a supplemental label.</p>
<ul style="list-style-type: none"> Do not apply through irrigation systems. Do not allow any application of Fanfare 2EC to contact fruit or foliage. Do not apply more than 0.5 lb. active ingredient (32 ounces formulated) per acre per year. Do not apply by air. 			

CITRUS (Florida only)(1)

PEST	DOSAGE		COMMENTS
	LB AI/A	FL OZ/A	
Blue Green Citrus Root Weevil (<i>Pachnaeus opalus</i>), Brown Leaf Notcher (<i>Epicaerus mexicanus</i>), Diaprepes Root Weevil (<i>Diaprepes abbreviatus</i>), Little Leaf Notcher (<i>Artipus floridanus</i>), Southern Blue Green Citrus Root Weevil (<i>Pachnaeus litus</i>)	0.25-0.50	16-32	<p>Ground application: Apply in water in a minimum of 40 gallons per acre.</p> <p>Greater spray volumes increase uniformity of coverage. Also coverage uniformity may be aided by using a pre and post irrigation application.</p> <p>Use a hand-gun or shielded sprayer to apply to individual citrus trees if they are not planted in solid rows.</p> <p>All citrus root weevils have a similar life cycle. They have three immature stages: egg, larva, and pupa. Adult weevils emerge from the soil and lay eggs on host plants above ground, the larvae enter the soil to feed on roots, and the pupae and teneral adult stages are spent below ground. Adults emerge beneath citrus trees throughout the year; it is at this time that Fanfare 2EC applications should be timed. Peak adult emergence varies within and among species and by region. Peak emergence for the blue-green root weevil is normally April and May. Diaprepes adult emergence from the soil appears to be triggered by the onset of regular rainfall events and can have two emergence peaks, in mid-May to mid-July and/or late-August to mid-October. The second peak is variable and may relate to host plant availability. Little leaf notcher has three generations per year. Although there is considerable overlap of generations, adults appear most abundant in April/May, July/August, and October/November.</p> <p>For best control of emerging root weevils, apply Fanfare 2EC to the soil beneath the citrus trees from the trunk to the drip line of the tree.</p> <p>Fanfare 2EC protects citrus tree roots from citrus root weevils by forming a barrier which provides contact activity on neonate larvae when they fall to the ground shortly after hatching from eggs which were oviposited in the citrus tree foliage.</p> <p>Once application is made, be careful not to disturb the treated soil.</p> <p>In areas where only a spring emergence is expected, use 32 ounces of Fanfare 2EC. In areas where a second emergence is expected, use 16 ounces of Fanfare 2EC in the early season and 16 ounces of Fanfare 2EC later in the season.</p> <p>If the length of control of Fanfare 2EC is not sufficient to cover the emergence of the root weevil, use other pest control measures as recommended by State Agricultural Extension Specialists or other local experts.</p>
Asian Cockroach, Fireant	0.1-0.25	6.4-16	

- Do not apply through irrigation systems.
- Do not allow any application of Fanfare 2EC to contact fruit or foliage.
- Do not apply more than 0.5 lb. active ingredient (32 ounces formulated) per acre per year.
- Do not apply by air.

CONIFER SEED ORCHARDS

(For Use Only in Alabama, Arkansas, Florida, Georgia, Louisiana, Mississippi, Oklahoma, South Carolina, Tennessee, Texas, Virginia)

PEST	DOSAGE		REMARKS
	LB/AI/A	FL. OZ/A	
Cone Worms, Seed Bugs, Seed Worms	0.1-0.2	6.4-12.8	<p>Ground application: Apply in water in a minimum of 100-500 gallons per acre.</p> <p>Air application: Apply in water in a minimum of 10 gallons per acre or 0.5 gallon refined vegetable oil per acre.</p> <p>Apply in sufficient water to obtain thorough coverage.</p> <p>Begin applications 7 days after peak pollen flight and continue on 30 day intervals up to a maximum of 0.6 lb. active per acre per season.</p>
<ul style="list-style-type: none">Do not make more than six applications per season or apply more than 0.6 lb. active ingredient (38.4 ounces formulated) per acre per season.			

CORN: FIELD CORN (GRAIN AND SILAGE), POPCORN, FIELD CORN GROWN FOR SEED (AT PLANTING)(30)

CORN: FIELD CORN (GRAIN AND SILAGE), POP CORN, FIELD CORN GROWN FOR SEED (AT PLANTING/30)			
PEST	DOSAGE		COMMENTS
	LB. AI	FL. OZ	
Corn Rootworm Larvae (Northern, Southern, Western)	.0046 pound active per 1,000 linear feet of row	0.30 fluid ounces per 1,000 linear feet of row	Ground application: Apply in water in a minimum of 3 gallons per acre. For use on corn at planting, apply in a 5 inch to 7 inch T-band over the open seed furrow. The spray nozzle should be centered over the row behind the planter shoe in front of the press wheel. In-furrow pop-up fertilizers may be used alone or in tank mixtures with Fanfare 2EC. See the section entitled MIXING INSTRUCTIONS, Fanfare 2EC with Fertilizer for additional instructions and precautions when mixing with fertilizers.
Army Cutworm, Cutworm Species, Grubs, Seedcorn Beetle, Seedcorn Maggot, True Armyworm or Armyworm Species, Wireworms	.0023 to .0046 pound active per 1,000 linear feet of row	0.15 to 0.30 fluid ounces per 1,000 linear feet of row	
<ul style="list-style-type: none">• Do not apply to soil where there is greater than 30% cover of crop residue remaining.• Do not graze livestock in treated area or cut treated crops for feed within 30 days of treatment.• Do not apply more than 0.1 lb. active ingredient (6.4 ounces formulated) per acre per season as an at-planting application.			

Row spacings (inches) ¹	40	38	36	30
Fanfare 2EC (pounds ai per acre)	0.060	0.064	0.069	0.080
Fanfare 2EC (formulated ounces per acre)	3.9	4.1	4.4	5.12

¹ Use this table to determine the Fanfare 2EC needs per acre.

CORN: FIELD CORN (GRAIN AND SILAGE), POPCORN, FIELD CORN GROWN FOR SEED (PRE & PPI)

PEST	DOSAGE		COMMENTS
	LB AI/A	FL OZ/A	
Armyworm spp., Black Cutworm, Seedcorn Maggot, Stalkborer, White Grub, Wireworm	0.047 to 0.062 Pre-Plant Incorporated (PPI)	3-4 Preplant Incorporated (PPI)	Ground application: Apply in water in a minimum of 3 gallons per acre. Use the recommended dosage as a preplant incorporated treatment either alone or in tank mix combination with registered preplant incorporated herbicides. Incorporate Fanfare 2EC to the intended planting depth but no deeper than 3 inches. The 3 to 4 oz. rate must be applied as PPI and can be tank mixed and applied with PPI herbicides. The 2.56 oz rate may be applied PRE and can be tank mixed and applied with PRE herbicides.
Armyworm spp, Black Cutworm, Stalkborer	0.040 Pre-Emergence (PRE)	2.56 Preemergence (PRE)	

CORN: FIELD CORN (GRAIN AND SILAGE), POPCORN, FIELD CORN GROWN FOR SEED (FOLIAR)(30)

PEST	DOSAGE		COMMENTS
	LB AI/A	FL OZ/A	
Aphids, Army Cutworm, Beet Armyworm, Cereal Leaf Beetle, Chinch Bug, Common Stalk Borer, Corn Earworm, Corn Rootworm Adult, Cucumber Beetle Adult, Cutworm Species, European Corn Borer, Fall Armyworm, Flea Beetle, Grasshoppers, Greenbug, Japanese Beetle Adult, Sap Beetle, Southern Armyworm, Southern Corn Leaf Beetle, Southwestern Corn Borer, Stinkbugs, Tarnished Plant Bug, True Armyworm or Armyworm Species, Webworms, Western Bean Cutworm, Yellowstriped Armyworm	0.033-0.10	2.1-6.4	<p>Ground application: Apply in water in a minimum of 10 gallons per acre except see specific comment below for TX, NM, OK, and AZ mite control.</p> <p>Air application: Apply in water in a minimum of 2 to 5 gallons per acre except see specific comment below for TX, NM, OK, and AZ mite control. In all states, insect control will be improved by increasing the finished spray per acre to 5 gallons. In Texas, New Mexico, Oklahoma, and Arizona, use a minimum of 10 gallons of water per acre by ground and 5 gallons of water per acre by air when making applications to control mites. Emulsified oil may be substituted for water. See section entitled MIXING INSTRUCTIONS for details on the amount of oil to use in the spray tank in lieu of water.</p> <p>Applications of Fanfare 2EC should be made as necessary to maintain control being careful not exceed reapplication intervals or maximum dosage rates specified in this section.</p> <p>For pests which attack the ear, apply just before silking.</p> <p>For corn borer control, make application just before or at egg hatch.</p> <p>For mite control, apply when colonies first form prior to leaf damage and before they disperse into the canopy (for Banks Grass Mite, before dispersal into the upper 2/3 of the plant). Use higher rates of Fanfare 2EC when pest pressure is severe or crop is under stress from drought and/or heat. When these conditions exist, tank mixtures with dimethoate have shown good control.</p>
Banks Grass Mite, Carmine Mite, Twospotted Spider Mite	0.08-0.10	5.12-6.4	
<ul style="list-style-type: none">Do not apply more than 0.3 lb. active ingredient (19.2 ounces formulated) per acre per season including pre and PPI, at-planting, plus foliar applications.Do not graze livestock in treated areas or cut treated crops for feed within 30 days of the last application.Use of ultra low volume (ULV) application on corn is prohibited.Do not make aerial or ground applications to corn if heavy rainfall is imminent.Use of Fanfare 2EC on corn is prohibited in all coastal counties.			

CORN: SWEET CORN, SWEET CORN GROWN FOR SEED (AT PLANTING)(30)

PEST	DOSAGE		COMMENTS
	LB AI	FL OZ	
Corn Rootworm Larvae (Northern, Southern, Western)	.0046 pound active per 1,000 linear feet of row	0.30 fluid ounces per 1,000 linear feet of row	Ground application: Apply in water in a minimum of 3 gallons per acre. For use on corn at planting, apply in a 5 inch to 7 inch T-band over the open seed furrow. The spray nozzle should be centered over the row behind the planter shoe in front of the press wheel. In-furrow pop-up fertilizers may be used alone or in tank mixtures with Fanfare 2EC. See the section entitled MIXING INSTRUCTIONS, Fanfare 2EC with Fertilizer for additional instructions and precautions when mixing with fertilizers.
Amy Cutworm, Cutworm Species, Grubs, Seedcorn Beetle, Seedcorn Maggot, True Armyworm or Armyworm species, Wireworms	.0023 to .0046 pound active per 1,000 linear feet of row	0.15 to 0.30 fluid ounces per 1,000 linear feet of row	
<ul style="list-style-type: none">Do not apply to soil where there is greater than 30% cover of crop residue remaining.Do not graze livestock in treated area or cut treated crops for feed within 30 days of treatment.Do not apply more than 0.1 lb. active ingredient (6.4 ounces formulated) per acre per season as an at plant application.			

Row spacings (inches) ¹	40	38	36	30
Fanfare 2EC (pounds ai per acre)	0.060	0.064	0.069	0.080
Fanfare 2EC (formulated ounces per acre)	3.9	4.1	4.4	5.12

¹ Use this table to determine the Fanfare 2EC needs per acre.

CORN: SWEET CORN, SWEET CORN GROWN FOR SEED (FOLIAR)(1)

PEST	DOSAGE		COMMENTS
	LB AI	FL OZ	
Aphids, Army Cutworm, Beet Armyworm, Cereal Leaf Beetle, Chinch Bug, Common Stalk Borer, Corn Earworm, Corn Rootworm Adult, Cucumber Beetle Adult, Cutworm Species, European Corn Borer, Fall Armyworm, Flea Beetle, Grasshoppers, Greenbugs, Japanese Beetle Adult, Sap Beetle, Southern Armyworm, Southern Corn Leaf Beetle, Southwestern Corn Borer, Stinkbugs, Tarnished Plant Bug, True Armyworm or Armyworm Species, Webworms, Western Bean Cutworm, Yellowstriped Armyworm	0.033-0.10	2.1-6.4	<p>Ground application: Apply in water in a minimum of 10 gallons per acre.</p> <p>Air application: Apply in water in a minimum of 2 gallons per acre.</p> <p>Emulsified oil may be substituted for water. See section entitled MIXING INSTRUCTIONS for details on the amount of oil to use in the spray tank in lieu of water.</p> <p>Applications of Fanfare 2EC should be made as necessary to maintain control being careful not exceed reapplication intervals or maximum dosage rates specified in this section.</p> <p>For pests which attack the ear, apply just before silking.</p> <p>For corn borer control, make application just before or at egg hatch.</p> <p>For mite control, apply when colonies first form prior to leaf damage and before they disperse into the canopy (for Banks Grass Mite, before dispersal into the upper 2/3 of the plant).</p> <p>Use higher rates of Fanfare 2EC when pest pressure is severe or crop is under stress from drought and/or heat. When these conditions exist, tank mixtures with dimethoate have shown acceptable control.</p>
Banks Grass Mite, Carmine Mite, Twospotted Spider Mite	0.08-0.10	5.12-6.4	
<ul style="list-style-type: none">• Do not apply more than 0.2 lb. active ingredient (12.8 ounces formulated) per acre per season.• Do not graze livestock in treated areas or cut treated crops for feed within 1 day of last application.• Use of ultra low volume (ULV) application on corn is prohibited.• Do not make aerial or ground applications to corn if heavy rainfall is imminent.• Use of Fanfare 2EC on corn is prohibited in all coastal counties.			

COTTON (14)

PEST	DOSAGE		COMMENTS
	LB. A/A	FL. OZ/A	
European Corn Borer, Soybean (Banded) Thrips, Tobacco Thrips	0.02-0.10	1.3-6.4	Ground application: Apply in water in a minimum of 5 gallons per acre. Air application: Apply in water in a minimum of 1 gallon per acre. Emulsified oil may be substituted for water. See section entitled MIXING INSTRUCTIONS for details on the amount of oil to use in the spray tank in lieu of water. ULV application: Apply in a minimum of 1 quart per acre using refined vegetable oil with aircraft calibrated to give adequate coverage. Applications of Fanfare 2EC should be made as necessary to maintain control being careful not exceed reapplication intervals or maximum dosage rates specified in this section. To Control Boll Weevil: Apply Fanfare 2EC at 3 to 4 day intervals until pest populations are reduced below economic threshold levels. To Control Mites and Aphids: Apply when pests first appear. Repeat as necessary to maintain control without exceeding maximum application rates and reapplication intervals. Higher rates should be used when an economic threshold has been established.
Boll Weevil, Bollworm, Cabbage Looper, Cotton Aphid, Cotton Fleahopper, Cotton Leafperforator, Cutworms, Fall Armyworm, Plant Bugs, Saltmarsh Caterpillar, Southern Garden Leafhopper, Stink Bugs, Tobacco Budworm, Whitefly, Yellow Striped Armyworm	0.04-0.10	2.6-6.4	
Beet Armyworm, Carmine Spider Mite, Lygus Spp. , Pink Bollworm, Twospotted Spider Mite	0.06-0.10	3.8-6.4	

- Do not apply more than 0.5 lb. active ingredient (32 ounces formulated) per acre per season.
- Do not graze livestock in treated areas or cut treated crops for feed.
- Do not make more than 10 synthetic pyrethroid applications (of one product or combination of products) to a cotton crop in one growing season. Synthetic pyrethroid products include Ambush®, Ammo®, Asana® XL, Baythroid®, Capture®, Danitol®, Karate®, Mustang®, and Scout X-TRA®.

CUCURBITS (3)

Chayote (fruit), Chinese waxgourd (Chinese preserving melon), Citron melon, Cucumber, Gherkin, Edible Gourd [(includes hyotan, cucuzza), *Luffa* spp. (includes hechima, Chinese okra), *Momordica* spp. (includes balsam apple, balsam pear, bitter melon, Chinese cucumber)], Muskmelon (hybrids and/or cultivars of *Cucumis melo*) (includes true cantaloupe, cantaloupe, casaba, crenshaw melon, golden pershaw melon, honeydew melon, honey balls, mango melon, Persian melon, pineapple melon, Santa Claus melon, and snake melon), Pumpkin (*Cucurbita* spp.), Squash, summer (includes crookneck squash, scallop squash, straightneck squash, vegetable marrow, zucchini), Squash, winter (includes butternut squash, calabaza, hubbard squash (*C. mixta*; *C. pepo*)(includes acorn squash, spaghetti squash)), Watermelon (includes hybrids and/or varieties of *Citrullus* spp.)

PEST	DOSAGE		COMMENTS
	LB A/A	FL OZ/A	
Aphids, Armyworms, Cabbage Looper, Corn Earworm, Cucumber Beetles, Cutworms, Grasshoppers, Leafhoppers, Melonworms, Pickleworms, Plant Bugs, Rindworms, Squash Bugs, Squash Vine Borer, Stink Bugs, Tobacco Budworm	0.04-0.10	2.6-6.4	Ground application: Apply in water in a minimum of 20 gallons per acre. Air application: Apply in water in a minimum of 5 gallons per acre. Emulsified oil may be substituted for water. See section entitled MIXING INSTRUCTIONS for details on the amount of oil to use in the spray tank in lieu of water.
Banks Grass Mite, Carmine Mite, Lygus spp., Twospotted Spider Mite, Whitefly	0.08-0.10	5.12-6.4	
<ul style="list-style-type: none">Do not apply more than 0.3 lb. active ingredient (19.2 ounces formulated) per acre per season.Do not make more than two applications after bloom.Repeat applications if needed to maintain control, but do not make applications less than 7 days apart.			

DRIED BEANS AND PEAS (14)**Dried cultivars of:****Bean** (*Lupinus* spp.): Grain Lupin, Sweet Lupin, White Lupin, White Sweet Lupin**Bean** (*Phaseolus* spp.): Field bean, Kidney Bean, Lima Bean (dry), Navy Bean, Pinto Bean, Tepary Bean**Bean** (*Vigna* spp.): Adzuki Bean, Blackeyed Pea, Catjang, Cowpea, Crowder Pea, Moth Bean, Mung Bean, Rice Bean, Southern Pea, Urd Bean**Broad Bean (dry):** Chickpea; Guar; Lablab Bean; Lentil; Pea (*Pisum*), Field pea; Pigeon Pea

PEST	DOSAGE		COMMENTS
	LB AI/A	FL OZ/A	
Banks Grass Mite, Twospotted Spider Mite, Carmine Mite, Lygus spp.	0.08-0.10	5.12-6.4	Ground application: Apply in water in a minimum of 10 gallons per acre. Air application: Apply in water in a minimum of 2 gallons per acre. Emulsified oil may be substituted for water. See section entitled MIXING INSTRUCTIONS for details on the amount of oil to use in the spray tank in lieu of water. Thorough coverage is essential to achieve control.
Aster Leafhopper, Flea Beetle, Grasshoppers, Leafhoppers	0.025-0.10	1.6-6.4	
Aphids, Beet Armyworm, Fall Armyworm, Southern Armyworm, Yellowstriped Armyworm, Bean Leaf Beetle, Cucumber Beetles, Japanese Beetle, Adult Sap Beetle, Plant Bug, Stink Bugs, Tarnished Plant Bug, Alfalfa Caterpillar, Cloverworm, European Corn Bore*, Cutworms, Western Bean Cutworm, Corn Earworm, Loopers, Corn Rootworm Adults, Thrips, Webworms, Pea Weevil, Pea Leaf Weevil, Whitefly, Imported Cabbageworm, Saltmarsh Caterpillar, Tobacco Budworm, Leafminer	0.33-0.10	2.1-6.4	
<ul style="list-style-type: none">Do not apply more than 0.2 lb. active ingredient (12.8 ounces formulated) to beans per acre per season.Do not make applications less than 7 days apart.			

FRUITING VEGETABLES (7)

Eggplant, Groundcherry, Pepino, Pepper (bell and non-bell)

PEST	DOSAGE		COMMENTS
	LB AI/A	FL OZ/A	
Armyworms (including Beet Armyworm, Fall Armyworm, Southern Yellowstriped Armyworm), Cabbage Loopers, Colorado Potato Beetle, Corn Earworm, Cucumber Beetles, European Corn Borer, Flea Beetles, Leafminers, Loopers, Pepper Weevil, Plant Bugs, Stink Bugs, Thrips, Tomato Hornworm, Tomato Pinworm, Vegetable Leafminer, Whitefly	0.033-0.10	2.1-6.4	Ground application: Apply in water in a minimum of 10 gallons per acre. Air application: Apply in water in a minimum of 2 gallons per acre. Emulsified oil may be substituted for water. See section entitled MIXING INSTRUCTIONS for details on the amount of oil to use in the spray tank in lieu of water.
Banks Grass Mite, Broad Mite, Carmine Mite, Lygus spp., Pacific Spider Mite, Twospotted Spider Mite	0.08-0.10	5.12-6.4	
<ul style="list-style-type: none">To maintain a proper spray interval, do not make applications less than 7 days apart.Do not apply more than 0.2 lb. active ingredient (12.8 ounces formulated) per acre per season.			

GRAPES (30)*

PEST	DOSAGE		COMMENTS
	LB AI/A	FL OZ/A	
Eastern Grape Leafhopper, Variegated Leafhopper, Western Grape Leafhopper	0.05-0.10	3.2-6.4	Ground application: Apply in water in a minimum of 25 gallons per acre. Air application: Apply in water in a minimum of 10 gallons per acre. Emulsified oil may be substituted for water. See section entitled MIXING INSTRUCTIONS for details on the amount of oil to use in the spray tank in lieu of water. When pest pressure is moderate to severe, use the higher rate. *Use in California not permitted unless accompanied by a supplemental label.
Black Vine Weevil, Glassywinged Sharpshooter, Twospotted Spider Mite	0.10	6.4	
• Do not apply more than 0.1 lb. active ingredient (6.4 ounces formulated) per acre per season.			

HOPS (14)

PEST	DOSAGE		COMMENTS
	LB AI/A	FL OZ/A	
Aphids, Armyworms, Cutworms, Leafrollers, Loopers	0.06-0.10	3.8-6.4	Ground application: Apply in water in a minimum of 100-150 gallons per acre in early season; 200-250 gallons per acre late season. Air application: Apply in water in a minimum of 10 gallons per acre. Make a directed spray up the vine 3 feet and the soil surface 1.5 to 2 feet on either side of the plant to control root weevil.
Root Weevils	0.05-0.10	3.2-6.4	
Twospotted Spider Mite	0.10	6.4	
<ul style="list-style-type: none">Do not apply more than 0.1 lb. active ingredient (6.4 ounces formulated) per acre per application.Do not apply more than 0.3 lb. active ingredient (19.2 ounces formulated) per acre per season.To maintain a proper spray interval, do not make applications less than 21 days apart.The use of ultra low volume (ULV) application on hops is prohibited.			

LEAFY BRASSICAS (7)

Broccoli Raab, Bok Choy, Kale, Mizuna, Mustard Greens, Mustard Spinach, Rape Greens, Turnip Greens*

PEST	DOSAGE		COMMENTS
	LB AI/A	FL OZ/A	
Aphids, Armyworms, Corn Earworm, Crickets, Cucumber Beetles, Cutworms, Diamondback Moth, Flea Beetles, Grasshoppers, Ground Beetles, Imported Cabbageworm, Japanese Beetle (adult), Leafhoppers, Loopers, Saltmarsh Caterpillar, Stink Bugs, Thrips, Tobacco Budworm, Whitefly, Wireworm (adults)	0.033-0.10	2.1-6.4	Ground application: Apply in water in a minimum of 10 gallons per acre. Air application: Apply in water in a minimum of 2 gallons per acre. Emulsified oil may be substituted for water. See section entitled MIXING INSTRUCTIONS for details on the amount of oil to use in the spray tank in lieu of water. Thorough coverage is essential to achieve control. *This use not permitted in California unless allowed by an approved supplemental label.
Banks Grass Mite, Carmine Mite, Lygus spp., Pacific Spider Mite, Twospotted Spider Mite	0.08-0.10	5.12-6.4	
<ul style="list-style-type: none">Do not apply more than 0.40 lb. active ingredient (25.6 ounces formulated) per acre per season.Repeat applications if needed to maintain control, but do not make applications less than 7 days apart.			

LETTUCE, HEAD (7)

PEST	DOSAGE		COMMENTS
	LB AI/A	FL OZ/A	
Aphids, Armyworms, Corn Earworm, Cucumber Beetles, Cutworms, Diamondback Moth, Flea Beetle, Imported Cabbageworm, Leafhoppers, Loopers, Salt Marsh Caterpillar, Stink Bug spp., Tobacco Budworm, Whitefly	0.033-0.10	2.1-6.4	Ground application: Apply in water in a minimum of 15 gallons per acre. Air application: Apply in water in a minimum of 5 gallons per acre. Emulsified oil may be substituted for water. See section entitled MIXING INSTRUCTIONS for details on the amount of oil to use in the spray tank in lieu of water.
Carmine Mite, Lygus spp., Twospotted Spider Mite	0.08-0.10	5.12-6.4	
<ul style="list-style-type: none">• To maintain a proper spray interval, do not make applications less than 7 days apart.• Do not apply more than 0.5 lb. active ingredient (32 ounces formulated) per acre per season.			

MAYHAW (30)

PEST	DOSAGE		COMMENTS
	LB AI/A	FL OZ/A	
Plum Curculio	0.08-0.10	5.12-6.4	Ground application: Apply in water in a minimum of 28 gallons of finished spray per acre. Air application: Apply in water in a minimum of 2 gallons per acre. Apply in sufficient water to obtain uniform coverage as needed.
<ul style="list-style-type: none"> Do not apply more than 0.2 pound active ingredient (12.8 ounces formulated) per acre per season. To maintain a proper spray interval, do not make applications less than 7 days apart. 			

OKRA (7) *

PEST	DOSAGE		COMMENTS
	LB AI/A	FL OZ/A	
Aphids, Armyworms, Corn Earworm, Cucumber Beetles, Cutworms, European Corn Borer, Flea Beetles, Japanese Beetle (Adult), Leafminers, Loopers, Stink Bugs, Thrips, Whitefly	0.033-0.10	2.1-6.4	Ground application: Apply in water in a minimum of 10 gallons of finished spray per acre. Air application: Apply in water in a minimum of 2 gallons per acre. Apply in sufficient water to obtain uniform coverage as needed.
Broad Mite, Carmine Mite, Lygus spp., Two Spotted Spider Mite	0.08-0.10	5.12-6.4	
<ul style="list-style-type: none">• To maintain a proper spray interval, do not make applications less than 7 days apart.• Do not apply more than 0.20 lb. active ingredient (12.8 ounces formulated) per acre per season.• *Not registered for use in California unless accompanied by a supplemental label.			

PEANUT (14)

PEST	DOSAGE		COMMENTS
	LB AI/A	FL OZ/A	
Beet Armyworm, Corn Earworm, Cutworm species, Fall Armyworm, Grasshoppers, Green Cloverworm, Leafhoppers, Lesser Cornstalk Borer, Loopers, Rednecked Peanut Worm, Southern Armyworm, Southern Corn Rootworm, Stink Bugs, Threecornered Alfalfa Hopper, Velvetbean Caterpillar, Yellowstriped Armyworm	0.033-0.1	2.1-6.4	Ground application: Apply in water in a minimum of 10 gallons of finished spray per acre. Air application: Apply in water in a minimum of 2 gallons per acre. Apply in sufficient water to obtain uniform coverage as needed.
Aphids, Spider Mites, Thrips, Whitefly	0.06-0.1	5.12-6.4	
<ul style="list-style-type: none">Do not apply more than 0.5 pound active ingredient (32 ounces formulated) per acre per season.To maintain a proper spray interval, do not make applications less than 14 days apart.Do not feed immature plants and peanut hay to livestock.			

PEARS (14)

PEST	DOSAGE		COMMENTS
	LB AI/A	FL OZ/A	
Aphids, Codling Moth, Cutworms, Green Fruitworm, Leafhoppers, Leafminers, Leafrollers, Lygus spp., Plant bugs, Plum Curculio, San Jose Scale (Crawlers), Stink Bugs, Tarnished Plant Bug	0.04 to 0.2	2.6 to 12.8	Ground application: Apply in water in a minimum of 200 gallons per acre (dilute) and 50 gallons per acre (concentrate). Air application: Apply in water in a minimum of 10 gallons per acre.
Twospotted Spider Mite, Yellow Mite	0.06 to 0.2	3.8 to 12.8	
European Red Mite	0.08 to 0.2	5.12 to 12.8	
<ul style="list-style-type: none">Do not apply more than 0.5 pound active ingredient (32 ounces formulated) per acre per season with no more than 0.45 (28.8 ounces formulated) pound active per acre applied after petal fall.To maintain a proper spray interval, do not make applications less than 30 days apart.Do not graze livestock in treated orchards or cut treated cover crops for feed.			

ROOT CROPS (except Sugar Beets)*

Burdock, edible; Carrot; Celeriac; Chervil, turnip rooted; Chicory; Ginseng; Horseradish; Parsley, turnip rooted; Parsnip; Radish; Radish, Oriental; Rutabaga; Salsify; Salsify, Black; Salsify, Spanish; Skirret; Turnip (21)

PEST	DOSAGE		COMMENTS
	LB AI/A	FL OZ/A	
Aphids, Beet Armyworm, Celery Leafminer, Corn Earworm, Cross-striped Cabbageworm, Cutworm species, Diamondback Moth, European Corn Borer, Fall Armyworm, Fire Ants, Fleabeetles, Green Cloverworm, Hornworms, Imported Cabbageworm, Loopers, Southern Armyworm, Spider Mites, Tobacco Budworm, Velvetbean Caterpillar, Whitefly, Yellowstriped Armyworm	0.08-0.10	5.12-6.4	Ground application: Apply in water in a minimum of 25 gallons of finished spray per acre. Air application: Apply in water in a minimum of 2 gallons per acre. Apply in sufficient water to obtain uniform coverage as needed.
<ul style="list-style-type: none"> Do not apply more than 0.5 pound active ingredient (32 ounces formulated) per acre per season. To maintain a proper spray interval, do not make applications less than 7 days apart. *Use in California not permitted unless accompanied by a supplemental label. 			

Garden Beet (1)

PEST	DOSAGE		COMMENTS
	LB A/A	FL OZ/A	
Aphids, Fire Ants, Fleabeetles, Lepidopterous Larvae, Spider Mites, Whitefly	0.08-0.10	5.12-6.4	Ground application: Apply in water in a minimum of 25 gallons of finished spray per acre. Air application: Apply in water in a minimum of 2 gallons per acre. Apply in sufficient water to obtain uniform coverage as needed.
<ul style="list-style-type: none"> Do not apply more than 0.40 pound active ingredient (25.6 ounces formulated) per acre per season. To maintain a proper spray interval, do not make applications less than 7 days apart. *Use in California not permitted unless accompanied by a supplemental label. 			

SOYBEAN (18)

PEST	DOSAGE		COMMENTS
	LB AI/A	FL OZ/A	
Soybean Aphid	0.033-0.10	2.1-6.4	Ground application: Apply in water in a minimum of 10 gallons per acre. Air application: Apply in water in a minimum of 2 gallons per acre.
Spider Mites	0.08-0.10	5.12-6.4	
<ul style="list-style-type: none">• To maintain a proper spray interval, do not make applications less than 30 days apart.• Do not apply more than 0.3 lb. active ingredient (12.8 ounces formulated) per acre per season.			

SPINACH (40)

PEST	DOSAGE		COMMENTS
	LB A/A	FL OZ/A	
Amyworms, Colorado Potato Beetle, Corn Earworm, Cucumber Beetles, Cutworms, European Corn Borer, Flea Beetles, Leafminers, Loopers, Pepper Weevil, Thrips, Tomato Hornworm, Tomato Pinworm, Whitefly	0.033-0.10	2.1-6.4	Ground application: Apply in water in a minimum of 10 gallons per acre. Air application: Apply in water in a minimum of 5 gallons per acre. For whitefly and fireant control, either at planting or as a foliar treatment, apply up to 6.4 oz. (0.1 lb. active) per acre being careful not exceed reapplication intervals or maximum dosage rates specified in this section.
Banks Grass Mite, Broad Mite, Carmine Mite, Fire Ants, Lygus spp., Pacific Spider Mite, Twospotted Spider Mite	0.08-0.10	5.12-6.4	
<ul style="list-style-type: none">To maintain a proper spray interval, do not make applications less than 7 days apart.Do not apply more than 0.4 lb. active ingredient (25.6 ounces formulated) per acre per season.			

SUCCULENT PEAS AND BEANS (3)

Pea (*Pisum* spp.): Dwarf pea, Edible-pod pea, English pea, Garden pea, Green pea, Snow pea, Sugar snap pea, Pigeon pea

Bean (*Phaseolus* spp.): Broadbean (succulent), Lima bean (green), Runner bean, Snap bean, Wax bean

Bean (*Vigna* spp.): Asparagus bean, Blackeyed pea, Chinese longbean, Cowpea, Moth bean, Southern pea, Yardlong bean, Jackbean, Soybean (immature seed), Sword bean

PEST	DOSAGE		COMMENTS
	LB A/A	FL OZ/A	
Aster Leafhopper, Flea Beetle, Grasshoppers, Leafhoppers	0.025-0.10	1.6-6.4	Ground application: Apply in water in a minimum of 10 gallons per acre. Air application: Apply in water in a minimum of 2 gallons per acre. Emulsified oil may be substituted for water. See section entitled MIXING INSTRUCTIONS for details on amount of oil to use in the spray tank.
Alfalfa Caterpillar, Aphids, Bean Leaf Beetle, Beet Armyworm, Cloverworm, Corn Earworm, Corn Rootworm Adult, Cucumber Beetle, Cutworms, European Corn Borer, Fall Armyworm, Japanese Beetle Adult, Loopers, Pea Leaf Weevil, Pea Weevil, Plant Bugs, Sap Beetle, Southern Armyworm, Stink Bugs, Tarnished Plant Bug, Thrips, Webworms, Western Bean Cutworm, Whitefly, Yellowstriped Armyworm	0.033-0.10	2.1-6.4	
Banks Grass Mite, Carmine Mite, Lygus spp., Twospotted Spider Mite	0.08-0.10	5.12-6.4	
<ul style="list-style-type: none">Do not apply more than 0.2 lb. active ingredient (12.8 ounces formulated) per acre per season.			

TOBACCO

PEST	DOSAGE		COMMENTS
	LB AI/A	FL OZ/A	
Armyworm spp., Cutworm spp., Mole Crickets, Stalkborers, Tobacco Flea Beetle (larvae), White Grubs, Wireworms	0.625-0.10	4.0-6.4	Pre-transplant soil applications: Apply 0.0625-0.1 active ingredient per acre in a minimum of 10 gallons per acre to control soil pests. Use of suitable equipment to incorporate into top 4" of the soil is required to control below ground pests. At-transplant water treatment application: Apply 0.0625-0.1 lb. active ingredient per acre in a water treatment application volume of the 10-200 gallons per acre. May be tank mixed with Command®, Spartan®, and other herbicides approved for tobacco use. Foliar applications: Apply 0.04-0.10 lb. active ingredient per acre foliar application up to and including layby in a minimum of 10 gallons per acre. May be tank mixed with Command, Spartan, and other herbicides approved for tobacco use.
Aphid spp., Armyworm spp., Chinch Bugs, Cutworm spp., Flea Beetle (adults), Grasshoppers, Green Bugs, Japanese Beetles, Stink Bugs, Tarnished Plant Bugs, Thrips, Whiteflies	0.04-0.10	2.56-6.4	
Lygus spp., Spider mites	0.1	6.4	
<ul style="list-style-type: none">• For foliar applications, do not make more than 2 applications per season.• For all applications, do not apply more than 0.2 lb. active ingredient (12.8 ounces formulated) per acre per season.• Do not apply later than layby.			

TOMATOES, TOMATILLO (1)

PEST	DOSAGE		COMMENTS
	LB AI/A	FL OZ/A	
Aphids, Armyworms (including Beet Armyworm, Fall Armyworm, Southern Yellowstriped Armyworm), Bean Leaf Beetle, Cabbageworms, Carmine Mite, Cloverworm, Corn Earworm, Corn Rootworm, Cucumber Beetle, Cutworms, Diamondback Moth, European Corn Borer, Flea Beetles, Flea Hoppers, Grasshoppers, Japanese Beetle (Adult), Leafhoppers, Loopers, Lygus spp., Melonworms, Pea Weevil, Pea Leaf Weevil, Pickleworms, Plant Bugs, Rindworms, Salt Marsh Caterpillar, Sap Beetle, Seedpod Weevil, Squash Bugs, Stink Bug spp., Tobacco Budworm, Tarnished Plant Bug, Thrips, Whitefly	0.033-0.08	2.1-5.2	Ground application: Apply in water in a minimum of 15 gallons per acre. Air application: Apply in water in a minimum of 3 gallons per acre.
Two Spotted Spider Mite	0.08-0.10	5.12-5.4	
<ul style="list-style-type: none">• To maintain a proper spray interval, do not make applications less than 10 days apart.• Do not make more than 4 applications per season.			

TUBEROUS AND CORM VEGETABLES (21)

Arracacha; Arrowroot; Chinese Artichoke; Jerusalem Artichoke; Edible Canna; Cassava (bitter and sweet); Chayote (root); Chufa; Dasheen (taro); Ginger; Leren; Potato; Sweet Potato; Tanier; Turmeric; Yam Bean; True Yam

PEST	DOSAGE		COMMENTS
	LB AI/A	FL OZ/A	
Corn Wireworm, Tobacco Wireworm	0.30 (at plant)	19.2 (at-plant)	In-furrow planting time treatment: Fanfare 2EC may be applied as an in-furrow planting time treatment for the control of wireworms, rootworms, and white grubs. Apply Fanfare 2EC at the rate of 0.3 pounds active ingredient per acre as an in-furrow spray or T-band spray at planting time. Lay-by treatment: Fanfare 2EC may be applied as a lay-by treatment for the control of wireworms, rootworms and white grubs. Apply Fanfare 2EC to the drill area and cover with soil utilizing cultivation equipment set to throw soil to the drill area. Apply Fanfare 2EC as a banded spray over the row at a rate of 0.05-0.15 pounds active ingredient per acre (3.2 to 9.6 ounces formulated) in 10 gallons per acre of spray. Foliar spray: Fanfare 2EC may be applied as a foliar spray for the control of the adult life stages of flea beetles, click beetles (wireworms), cucumber beetles (rootworms), white fringed beetles, and May/June beetles (white grubs). Apply Fanfare 2EC at the rate of 0.1 lb. active ingredient per acre (6.4 ounces formulated) in 10 gallons of spray by ground and 3 gallons of spray by air.
Southern Potato Wireworm, Japanese Beetle Grubs, June Beetle	0.05-0.15 (lay-by)	3.2-9.6 (lay-by)	
Sweetpotato Flea Beetle, Cucumber Beetle, Sweetpotato Weevil, Banded Cucumber Beetle, Black Flea Beetle, Whitefringed Beetle, White Grub, Sugarcane Beetle, Rootworms	0.033-0.10 (foliar)	2.1-6.4 (foliar)	
<ul style="list-style-type: none">For foliar applications, do not make more than 2 foliar applications per season and do not make application less than 21 days apart.Do not apply more than 0.5 lb. active ingredient (32 ounces formulated) per acre per season, including soil applications.			

STORAGE AND DISPOSAL

PROHIBITIONS: Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited. Do not reuse empty container.
PESTICIDE STORAGE: DO NOT ALLOW PRODUCT TO FREEZE. Do not store below 40° F. If crystals are observed, warm material to above 60° F by placing container in warm location. Shake or roll container periodically to redissolve solids. Keep out of reach of children and animals. Store in original containers only. Store in a cool, dry place and avoid excess heat. Do not contaminate other pesticides, fertilizers, water, food, or feed by storage or disposal.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative of the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL:

NON-REFILLABLE CONTAINER: Do not reuse or refill this container. Clean container promptly after emptying. Offer for recycling if available. If recycling is not available, puncture or dispose of in a sanitary landfill or incineration or if allowed by state and local authorities by burning. If burned, stay out of smoke.

Non-refillable container 5 gallons or less: Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Offer for recycling if available.

Non-refillable container more than 5 gallons: Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Offer for recycling if available.

REFILLABLE CONTAINER: Refill this container with Fanfare 2EC (containing the active ingredient bifenthrin) only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10% full with water. Agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

SPILL, FIRE, LEAK or OTHER CHEMICAL EMERGENCY: In case of spill or leak on floor or paved surfaces, soak up with sand earth, or synthetic absorbent. Remove to chemical waste area. In the event of a major spill, fire, or other emergency, call Infotrac at 1-800-535-5053, day or night.

LIMITATION OF WARRANTY AND LIABILITY

Read the entire directions for use, conditions of warranties and limitations of liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following **CONDITIONS, DISCLAIMER OF WARRANTIES** and **LIMITATIONS OF LIABILITY**.

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Makhteshim Agan of North America, Inc. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: To the extent consistent with applicable law, Makhteshim Agan of North America, Inc. makes no other warranties, express or implied, of merchantability or of fitness for a particular purpose or otherwise, that extend beyond the statements made on this label. No agent of Makhteshim Agan of North America, Inc. is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. To the extent consistent with applicable law, Makhteshim Agan of North America, Inc. disclaims any liability whatsoever for special, incidental or consequential damages resulting from the use or handling of this product.

LIMITATIONS OF LIABILITY: To the extent consistent with applicable law, the exclusive remedy of the user or buyer for any and all losses, injuries or damages resulting from the use or handling of this product, whether in contract, warranty, tort, negligence, strict liability or otherwise, shall not exceed the purchase price paid or at Makhteshim Agan of North America, Inc.'s election, the replacement of product.

Ammo, Capture, Mustang, U-Turn—trademarks of FMC Corporation
Ambush, Karate—Trademarks of Zeneca, Inc.
Asana—trademark of E.I. duPont de Nemours & Company
Baythroid—trademark of Bayer Aktiengesellschaft
Danitol—trademark of Sumitomo Chemical Company, Ltd.
Scout Xtra—trademark of Hoechst Schering Agrevo A.A.
Fanfare is a registered trademark of Makhteshim Chemical Works, Ltd.

Fanfare 2EC (66222-99)(EPA app 08-25-08)(Notif to EPA 01-10-09)