
66222-121





packet 116 UNITED STATES ENVIRONMENTAL PROTECTION AGENCE WASHINGTON, DC 20460

> OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

Makhteshim Agan of North America, Inc. **Brooke Hedrick** 3120 Highwoods Blvd., #100 Raleigh, NC 27604

AUG 1 4 2012

Product Name: Ag Streptomycin EPA Reg. No.: 66222-121 Amendment dated 6/26/2012 Subject: EPA Decision Number: 468004

Dear Mr. Hedrick:

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended is acceptable provided the following change is made:

1. On page 5, change the first sentence under Storage and Disposal to read "Do not contaminate water, food, or feed by storage and disposal."

One copy of the label stamped "Accepted with comments" is enclosed for your records. This label supersedes all labels previously accepted for this product. Please submit one copy of the final printed label before the product is released for shipment.

If you have any questions, please contact Erin Malone at (703) 347-0253 or via email at malone.erin@epa.gov.

Sincerely,

Mary L. Waller

Mary L. Waller Product Manager (21) Fungicide Branch Registration Division (7504P)

Enclosure

Ag Streptomycin Bactericide

ACTIVE INGREDIENT

Streptomycin Sulfate (equivalent to 17% streptomycin) (CAS #3810 74 0) OTHER INGREDIENTS

ACCEPTED with COMMENTS In EPA Letter Dated:

8/14/2012

Under the Foderal Insecticide, Fungicide, and Rodenticide Act as amended for she postietae registered under EIA hey. No. 6(222-121

EPA Reg No 66222 121

Manufactured for Makhteshim Agan of North America Inc 3120 Highwoods Blvd Suite 100 Raleigh NC 27604

KEEP OUT OF REACH OF CHILDREN

CAUTION

EPA Est No _____

Net Contents ____ Lbs

	FIRST AID
IF IN EYES	 Hold eye open and rinse slowly and gently with water for 15 20 minutes Remove contact lenses if present after the first 5 minutes then continue rinsing eye Call a poison control center or doctor for treatment advice
IF SWALLOWED	Call a poison control center or doctor immediately for treatment advice Have person sip a glass of water if able to swallow Do not induce vomiting unless told to do so by a poison control center or doctor • Do not give anything by mouth to an unconscious person
IF ON SKIN OR CLOTHING	 Take off contaminated clothing Rinse skin immediately with plenty of water for 15 20 minutes Call a poison control center or doctor for treatment advice
IF INHALED	 Move person to fresh air If person is not breathing call 911 or an ambulance then give artificial respiration preferably mouth to mouth if possible Call a poison control center or doctor for further treatment advice
	ntainer or label with you when calling a poison control center or doctor or going for treatment. You may also 1 877 250 9291 for emergency medical treatment information

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if absorbed through skin Causes moderate eye irritation Avoid contact with skin eyes or clothing May cause allergic skin reactions. Do not breathe dust or spray mist

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical resistant to this product are listed below. If you want more options follow the instructions for Category A on an EPA chemical resistance category selection chart

Applicators and other handlers must wear

Long sleeved shirt and long pants

Chemical resistant gloves made of any waterproof material

Shoes plus socks

A NIOSH approved respirator with any N $\,$ R $\,$ P $\,$ or HE filter

Follow manufacturers instructions for cleaning/maintaining PPE. If no such instructions for washables use detergent and hot water. Keep and wash PPE separately from other laundry

This material is not to be used for medical veterinary or human purposes

Page 1 of 5

% BY WT 22 4% <u>77 6%</u> TOTAL 100 0%

2/6

USER SAFETY RECOMMENDATION

Users should

Wash hands before eating drinking chewing gum using tobacco or using the toilet Remove PPE immediately after handling this product Wash outside of gloves before removing. As soon as possible wash thoroughly and change into clean clothing

316

ENVIRONMENTAL HAZARDS

This product may be hazardous to aquatic plants. Do not apply directly to water to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of wastes

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling

Do not apply this product through any type of irrigation system

Do not apply this product in a way that will contact workers or other persons either directly or through drift Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard 40 CFR Part 170 This Standard contains requirements for the protection of agricultural workers on farms forests nurseries and greenhouses and handlers of agricultural pesticides. It contains requirements for training decontamination notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted entry interval. The requirements in this box only apply to uses that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated such as plants soil or water is

Coveralls over long sleeved shirt and long pants

Chemical resistant gloves made of any waterproof material

NIOSH approved respirator with any N R P or HE filter

Chemical resistant shoes plus socks

MIXING INSTRUCTIONS						
Concentration	Quantity Ag Streptomycin					
desired	per Volume of Water					
ppm	50 gals	100 gals	500 gals			
50	2 oz	4 oz	20 oz			
60	2 4 oz	4 8 oz	1 1/2 lb			
100	4 oz	1/2 lb	2 1/2 lb			
200	1/2 lb	1 lb	5 lb			

ppm = parts per million

|--|

CROPS & DISEASES	CONCENTRATION	FIRST SPRAY	FOLLOW UP SPRAY SCHEDULE	
Apples Fire Blight	24-48 oz Ag Streptomycin per acre (equivalent to 50 100 ppm at 600 gals /A)	Spray trees at 20% 30% bloom	Spray trees every 3 4 days during blossom time Apply sprays after petal fall every 10 14 days to control twig blight (This could mean an additional 6 8 applications after blossom sprays) Do not apply within 50 days of harvest	
Apples Fire Blight (CA OR and WA)	28 8 oz Ag Streptomycin per acre (equivalent to 60 ppm at 600 gals /A)	Full bloom	Apply at petal fall and late secondary bloom Continue to spray at 5 to 7 day intervals to maintain disease control but not later than 50 days before harvest	
Celery Bacterial Blight (Florida)	200 ppm	Apply first spray when seedlings are in the 2 leaf stage when first true leaves appear	Apply at 4 to 5 day intervals Continue applications until celery is transplanted in the field	
Chrysanthemums Bacterial Wilt	50 ppm		streptomycin solution for 4 hours plant as	
Dieffenbachia Cuttings Bacterial Stem Rot	200 ppm	Soak cuttings in streptomycin solution for 20 minutes Plant cuttings in sterilized rooting medium		
	100 ppm	To check spread of stem rot in stock plants use 100 ppm streptomycin spray every 5 7 days		
Home Garden Apple Trees Home Garden Pear Trees Pyracantha (Fire Thorn Bush) Fire Blight of the Rosaceae (CA)	100 ppm	Apply Ag Streptomycin in foliar and blossom sprays Apply first spray at start of blossoming period Continue spray application every 3-4 days during blossom time Apply additional sprays every 5 7 days after blossom period when weather favors spread of fire blight Do not apply after fruit is visible		
Pears Fire Blight	24-48 oz Ag Streptomycin per acre (equivalent to 50 100 ppm at 600 gals /A)	Spray trees at 20% 30% bloom	Spray trees every 3 4 days during blossom time Apply sprays after petal fall every 10 14 days to control twig blight (This could mean an additional 6 8 applications after blossom sprays) Do not apply within 30 days of harvest	
Pears Fire Blight (CA OR and WA)	28 8 oz Ag Streptomycin per acre (equivalent to 60 ppm at 600 gals /A)	10% bloom	Repeat at 5 day intervals until all late bloom is over (This could mean 12 15 applications) Continue to spray at 5 to 7 day intervals to control shoot and fruit infections Do not apply within 30 days of harvest	
Peppers Bacterial Spot	200 ppm	Apply first spray when seedlings are in the 2 leaf stage when first true leaves appear	Apply at 4 to 5 day intervals Continue applications until transplanted in the field	
Philodendron Bacterial Leaf Rot	200 ppm	Apply as preventative or at first signs of water soaked areas on leaf	Apply every 4 5 days	
	For curative action	Remove all rotted leaves from plant and then spray at 200 ppm every 4 days		
Potatoes Soft Rot and Blackleg	100 ppm	Soak cut seed pieces in streptomycin solution for several minutes plant as usual Note A suitable fungicide (such as Captan Phygon dithiocarbamates) should be used as an adjunct to this treatment for the control of fungal diseases associated with potato seed pieces		

	(/	5/	
Roses (New Jersey) Crown Gall	200 p _r	Remove infected plan, ut out gall tissue Soak the plant root system and cut surfaces of the infected area in streptomycin solution for 15 minutes Replant rose bushes in soil free of the crown gall organisms		
	50 ppm	Use 50 ppm Ag Streptomycin in watering solution and in foliar sprays applied weekly starting one week after planting as an adjunct to this treatment		
Tobacco Wildfire and Blue Mold	100 ppm for preventative action	Apply first spray when plants are in the 2 leaf stage or about the size of a dime or when blue mold first appears in the area	Repeat application at 5 to 7 day intervals until plants are set in the field Additional protection may be obtained by spraying field plants with 100 ppm in a weekly spray schedule	
	200 ppm for curative action	where applications hav	fire has been a problem in recent years or the been delayed until disease appears a streptomycin is recommended Follow the ve	
Tomatoes Bacterial Spot	200 ppm	Apply first spray when seedlings are in the 2 leaf stage when first true leaves appear	Apply at 4 to 5 day intervals Continue applications until transplanted in the field	

Additional information regarding use of Ag Streptomycin may be obtained from your local Agricultural Extension Agent or State Experimental Station

STORAGE AND DISPOSAL

Do not contaminate water food or feed by storage or disposal

(

PESTICIDE STORAGE Keep tightly closed Storage should be at a cool temperature when possible and within minimum exposure to the atmosphere

PESTICIDE DISPOSAL Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility

CONTAINER HANDLING

Nonrefiliable Container (flexible bag all weights) Nonrefiliable container Do not reuse or refili this container. Offer for recycling if available. Then dispose of bag in a sanitary landfill by incineration or if allowed by State and local authorities by burning. If burned stay out of smoke

Nonrefillable Container (rigid fifty lbs or less) Nonrefillable container Do not reuse or refill this container Offer for recycling if available Clean container promptly after emptying Triple rinse as follows Empty the remaining contents into application equipment or a mix tank Fill the container 1/4 full with water and recap Shake for 10 seconds Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal Drain for 10 seconds after the flow begins to drip Repeat this procedure two more times Then dispose of bag in a sanitary landfill by incineration or if allowed by State and local authorities by burning If burned stay out of smoke

Nonrefillable Container (rigid greater than fifty lbs) Nonrefillable container Do not reuse or refill this container Offer for recycling if available Clean container promptly after emptying Triple rinse as follows Empty the remaining contents into application equipment or a mix tank Fill the container 1/4 full with water Replace and tighten closures Tip container on its side and roll it back and forth ensuring at least one complete revolution for 30 seconds Stand the container on its end and tip it back and forth several times Turn the container over onto its other end and tip it back and forth several times Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal Repeat this procedure two more times Then dispose of bag in a sanitary landfill by incineration or if allowed by State and local authorities by burning If burned stay out of smoke

Refillable Container Refillable container Refill this container with streptomycin sulfate only Do not reuse this container for any other purpose Cleaning the container before final disposal is the responsibility of the person disposing of the container Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

FOR 24 HOUR EMERGENCY ASSISTANCE (SPILL LEAK OR FIRE) CALL INFOTRAC AT 1 800 535 5053

IMPORTANT Read the information below before using this product. If the terms are not acceptable, you should return the unopened product container immediately for a complete refund

LIMITATION OF WARRANTY AND LIABILITY

Read the entire directions for use conditions of warranties and limitations of liability before using this product If terms are not acceptable return the unopened product container at once

By using this product user or buyer accepts the following CONDITIONS DISCLAIMER OF WARRANTIES and LIMITATIONS OF LIABILITY

CONDITIONS The directions for use of this product are believed to be adequate and must be followed carefully However it is impossible to eliminate all risks associated with the use of this product. Crop injury ineffectiveness or other unintended consequences may result because of such factors as weather conditions presence of other materials or the manner of use or application all of which are beyond the control of Makhteshim Agan of North America. Inc. To the extent consistent with applicable law all such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES To the extent consistent with applicable law Makhteshim Agan of North America Inc makes no other warranties express or implied of merchantability or of fitness for a particular purpose or otherwise that extend beyond the statements made on this label. No agent of Makhteshim Agan of North America. Inc. is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. To the extent consistent with applicable law Makhteshim Agan of North America. Inc. disclaims any liability whatsoever for special incidental or consequential damages resulting from the use or handling of this product.

LIMITATIONS OF LIABILITY To the extent consistent with applicable law the exclusive remedy of the user or buyer for any and all losses injuries or damages resulting from the use or handling of this product whether in contract warranty tort negligence strict liability or otherwise shall not exceed the purchase price paid or at Makhteshim Agan of North America Inc s election the replacement of product

Harbour (66222 121)

(Notif to add product identifier 05 21 12) (Notif to modify state descriptors)