

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, D.C. 20460

OFFICE OF PREVENTION. PESTICIDES AND TOXIC SUBSTANCES

1/10

FEB 2 3 2009

Christie Hitchcock Regulatory Specialist Makhteshim Agan of North America, Inc. 4515 Falls of Neuse Road, Suite 300 Raleigh, NC 27609

Subject: Label Notifications for Pesticide Registration Notice 2007-4; Updated Warranty Statements per Pesticide Registration Notice 98-10

Dear Ms. Hitchcock,

The Agency is in receipt of your Applications for Pesticide Notification under Pesticide Registration Notices (PRNs) 98-10 and 2007-4 dated August 29, 2008 (and the label resubmissions sent February 19, 2009) for the following products:

Ag Streptomycin	EPA Registration Number 66222-121
Klean Pik [®]	EPA Registration Number 66222-124
Parallel [®] Plus	EPA Registration Number 66222-132

The Registration Division (RD) has conducted its review of these requests for their applicability under PRNs 98-10 and 2007-4 and finds that the label changes requested fall within the scope of PRNs 98-10 and 2007-4. The labels submitted with the application have been stamped "Notification" and will be placed in our records.

Please be reminded that 40 CFR Part 156.140(a)(4) requires that a batch code, lot number, or other code identifying the batch of the pesticide distributed and sold be placed on <u>nonrefillable</u> containers. The code may appear either on the label (and can be added by nonnotification/PR Notice 98-10) or durably marked on the container itself. If you have any questions, please call me directly at 703-305-6249 or Steve Schaible of my staff at 703-308-9362.

C. Hollonar Sincerely, Linda Arrington

Notifications & Minor Formulations Team Leader Registration Division (7505P) Office of Pesticide Programs 2/10

Please read instructions of	reverse before complete	ting form		Form A	pproved		2070-008	. Approval expires 2-28-9
Please read instructions on reverse before completing form. Form A United States Environmental Protection Agency Washington, DC 20460				Registra Amenda Other	ation	OPP Identifier Number		
		Applicati	on for Pest	ticide - Sec	ction	I	· · · ·	······································
1. Company/Product Number 66222-124 4. Company/Product (Name)		2. EPA Product Ma Linda Arrington PM# 23		nager 3.		3. Pr	oposed Classification	
Klean-Pik® 500SC			Re	gistration Su	pport I	Branch		
5. Name and Address of A Makhteshim Agan o 4515 Falls of Neuse Raleigh, NC 27609	f North America, I	•	(b)(to: EP	Expedited Re i), my product A Reg. No.	t is sim NO	In accorda	Ince with	FIFRA Section 3(c)(3)
			Section	- 11		(· · · · · · · · · · · · · · · · · · ·
Amendment - Explain Resubmission in result Notification - Explain Explanation: Use addition Notification of label change regulations at 40 CFR 152. 156.146, and 156.156. No violation of 18 U.S.C. Sec.	ponse to Agency letter n below. onal page(s) if necessar per PR Notice 98-10 and 46, and with the guidance other changes have beer	y. (For section d PR Notice 20 e of PR Notice n made to the l	007-4. This notific 2007-4 and requirabeling or the Co	Agency le "Me Too" Other - Ex II.) cation is consiste irrements of EPA nfidential Staten	ent with Applica	ation. slow. the provision ations at 40 (Formula of thi	s of PR Not CFR §§156. is product, I	.10, 156.140, 156.144, I understand that it is a
terms of PR Notice 98-10 a violation of FIFRA and I ma	y be subject to enforcem	the requireme	nts of 40 CFR §§ I penalties under Section	sections 12 and	14 of Fl	14, 156.146, a FRA.	and 156.15	6, this product may be in
1. Material This Product W Child-Resistant Packaging ↓ Yes ↓ No * Certification must be submitted	Unit Packaged In:	No. per container	Water Solub ✓ Yes ✓ No If "Yes" Peckage wg	le Packaging No. per t contain		2. Type of	Container Metal Plastic Glass Paper Other (S	pecify)
3. Location of Net Contents	Information	4. Size(s) Re	atail Container		5. Lo	cation of La		ns
6. Manner in Which Label is Affixed to Product []Litho		or less; more than 5 gals on label graph r glued cifed						
			Section	- IV				
1. Contact Point (Complet	e items directly below f	for identificati	on of individual	to be contacted	l, if nec	essary, to pi	ocess this	application.)
Name Christie Hitchcock			Title Regulatory S	pecialist			Telephone 919-256-9	9 No. (Include Area Code) 9342
-	ements I have made on ny knowlinglly false or I aw.		d all attachments				-	6. Date Application Received (Stamped)
2. Signature Christie Atcheoch			3. Title Regulatory Specialist					
4. Typed Name Christie Hitchcock			5. Date	8-29-08				

Yellow - Applicant Copy



August 29, 2008

Document Processing Desk (NOTIF) Registration Division (7504P) OPP, USEPA Ariel Rios Building 1200 Pennsylvania Ave, NW Washington DC 20460

Re: Klean-Pik[®] 500SC, EPA Reg. No. 66222-124 Notification per PRN 98-10 and 2007-4

To Whom It May Concern:

We are notifying the Agency of several label updates as allowed in PR Notices 98-10 and 2007-4. The changes are summarized below:

4/10

- Added "Manufactured for" in front of the company name (allowed per PRN 98-10 §II(N)(2))
- Updated the Warranty Statement that is now consistent with EPA's guidance document dated 10-17-06) (allowed per PRN 98-10 §II(J))
- Updated container disposal instructions per PRN 2007-4

In support of this submission, the following documents are attached:

- Application for Pesticide Registration (EPA Form 8570-1)
- One copy of final printed labeling

Please contact me at 919-256-9342 or by email at <u>chitchcock@manainc.com</u> if you have any questions regarding this submission or would like a copy of the annotated label showing the changes for your files.

Sincerely,

mitie Altehook

Christie Hitchcock Regulatory Specialist

Enclosures

4515 Falls of Neuse Road, Suite 300, Raleigh NC 27609 T: 919.256.9305 + F: 919.256.9308

KLEAN-PIK[®] 500SC

For Agricultural Use Only COTTON DEFOLIANT

ACTIVE INGREDIENT:	% BY WT.
Thidiazuron: (N-phenyl-N-1,2,3-thiadiazol-5-ylurea)*	
INERT INGREDIENTS:	
	TOTAL 100.00%

*Contains 4.17 lbs of active ingredient per gallon.

KEEP OUT OF REACH OF CHILDREN CAUTION

Manufactured for: Makhteshim Agan of North America, Inc. 4515 Falls of Neuse Rd., Suite 300 Raleigh, NC 27609

GALS.

NET CONTENTS:

EPA Reg. No. 66222-124

	NET CONTENTO: GALO.
· · · · · · · · · · ·	FIRST AID
IF SWALLOWED:	Immediately call a poison control center or doctor for treatment advice.
	Have person sip a glass of water if able to swallow.
	• Do not induce vomiting unless told to do so by the poison control center or doctor.
	Do not give anything by mouth to an unconscious person.
IF INHALED:	Move person to fresh air.
	• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.
	Call a poison control center or doctor for further treatment advice.
IF IN EYES:	Hold eye open and rinse slowly and gently with water for 15-20 minutes.
	• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
	 Call a poison control center or doctor for treatment advice.
IF ON SKIN OR	Take off contaminated clothing.
CLOTHING:	Rinse skin immediately with plenty of water for 15-20 minutes.
	Call a poison control center or doctor for treatment advice.
Have the product c	container or label with you when calling a poison control center or doctor or going for
treatment. You may	also contact PROSAR at 1-877-250-9291 for emergency medical treatment information.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if swallowed, inhaled, or absorbed through the skin. Avoid breathing spray mist. Avoid contact with skin, eyes, and clothing. Do not contaminate food or feedstuffs.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks
- Waterproof gloves

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

NOTIFICATION

FEB 2 3 2009

EPA Est. No.

Page 1 of 6

5/10

ENGINEERING CONTROLS STATEMENT: When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

Users should:

USER SAFETY RECOMMENDATIONS

• Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.

Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of cleaning equipment or disposing of equipment washwaters or rinsate. Do not apply when weather conditions favor drift from target area.

GENERAL INFORMATION

Klean-Pik 500SC is used for removal of leaves prior to harvesting. Klean-Pik 500SC disperses easily in water and can be used in any ground and aircraft equipment. It may take several days before the effect of Klean-Pik 500SC becomes noticeable. Adverse conditions, such as low temperatures, may require higher dosages and/or longer times for more complete defoliation. Klean-Pik 500SC inhibits regrowth after defoliation and thus reduces staining of lint during harvesting and ginning.

Use Precautions:

- Do not apply this product through any type of irrigation system.
- Rainfall within 24 hours after application will reduce the effectiveness of Klean-Pik 500SC. Use only freshly
 prepared sprays. Do not store spray mixture overnight.
- Do not feed foliage from treated cotton plants or gin trash to livestock.

• Do not plant the following crops earlier than the specified periods after the application of Klean-Pik 500SC :

Small grain, sorghum, corn, or root crops (except carrots, onions, and sugar beets)	Two (2) weeks
Legumes (including alfalfa) or leafy vegetables (except lettuce and spinach)	Two (2) months
Sugar beets	Four (4) months
Carrots, onions, or spinach	Nine (9) months
Lettuce	Nine (9) months and only following deep plowing (12-15 inches)

Do not use immature crops for food or feed.

Do not allow spray drift to contact crops other than the target crop of mature cotton, or cotton which you desire to defoliate, as this product may injure or defoliate other crops.

Particular care should be taken when applying Klean-Pik 500SC adjacent to lettuce, citrus, or cantaloupe. Tank mixes with organophosphates may increase nontarget crop phytotoxicity.

PRACTICES TO LOWER THE POTETNIAL FOR DRIFT ONTO NONTARGET CROPS

During applications, particularly under windy conditions, Klean-Pik 500SC may drift to nontarget crops.

To help reduce the drift potential, use the following practices:

- Do not apply Klean-Pik 500SC by ground or air when wind speeds exceed ten (10) miles per hour at the time of application. Follow local recommendations if wind speeds of less than ten (10) miles per hour are specified in those recommendations.
- Use of low nozzle pressure (20-30 psi) is recommended.
- Do not apply Klean-Pik 500SC when a temperature inversion is present or when conditions favor an inversion prior to completing applications(s).
- Additionally, do not apply Klean-Pik 500SC by air within one-half (1/2) mile of lettuce. Do not apply Klean-Pik 500SC by ground equipment within 100 feet of lettuce.

PRACTICES TO LOWER DRIFT POTENTIAL ONTO NON-TARGET CROPS IN THE RIO GRANDE VALLEY OF TEXAS

To help reduce the drift potential, use the following practices:

• Do not apply Klean -Pik 500SC by ground or air when wind speeds exceed ten (10) miles per hour at the time of

application. Follow local recommendations if wind speeds of less than ten (10) miles per hour are specified in those recommendations.

- Use of low nozzle pressure (20-30 psi) is recommended.
- Do not apply Klean-Pik 500SC when a temperature inversion is present or when conditions favor an inversion prior to completing applications(s).
- Use the largest nozzle orifice possible which permits proper deposition and coverage of product.
- In addition, for citrus crops, do not apply Klean-Pik 500SC by air when citrus in flush is within five (5) miles downwind of the point of application. Do not apply Klean-Pik 500SC by ground when citrus in flush is within onehalf (1/2) mile downwind of the point of application.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

PPE required for early entry into treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Waterproof gloves
- Shoes plus socks

TIME OF APPLICATION

Apply Klean-Pik 500SC only to mature cotton plants when the last boll you expect to harvest is mature. A boll can be described as "mature" when it is too hard to be dented when squeezed between thumb and fingers, is difficult to slice with a sharp knife, and/or when seeds cut in cross sections have fully developed cotyledons, as evidenced by an absence of jelly within the seed. Apply Klean-Pik 500SC at least 5 days prior to anticipated harvest.

NOTE: Good defoliation with Klean-Pik 500SC is dependent upon three factors:

1. Actively growing cotton

2. High humidity

3. High moisture content in leaf of cotton plant

Use of Klean-Pik 500SC alone (without a tank mix partner) when nighttime temperatures are expected to fall below 60°F can result in less than desirable defoliation and/or regrowth inhibition.

Performance of Klean-Pik 500SC is variable under low temperatures, ideally, nighttime temperatures 2-3 days prior to and following application should be above 60°F or total defoliation and regrowth inhibition can be reduced. Use of Klean-Pik 500SC on heat and drought-stressed cotton (low leaf moisture, thick cuticle, etc.) may result in less than satisfactory defoliation and regrowth inhibition.

USE OF ADJUVANTS (CROP OILS)

The use of adjuvants such as petroleum-based crops oils or penetrating oils approved for use on growing crops has been shown to improve performance during low nighttime temperatures (60°F-65°F) or when defoliating drought-stressed cotton. Refer to specific adjuvant label for rate recommendations.

The use of adjuvants is recommended when applying Klean-Pik 500SC in the desert Southwest (i.e. Arizona and California).

PRECAUTION: The addition of adjuvants can cause desiccation and/or leaf freezing during periods of high temperatures. The use of compounds that desiccate leaf tissue is not recommended.

MIXING INSTRUCTIONS: Fill the spray tank with one-half of the total amount of water to be used. Begin agitation and

Page 3 of 6

add the recommended amount of Klean-Pik 500SC. Fill the tank with the additional quantity of water required. Maintain sufficient agitation to ensure a uniform spray mixture during application.

TANK MIX ADJUVANTS: When using tank mixes of Klean-Pik 500SC and organophosphates, the use of a surfactant or compatibility agent is recommended to improve tank clean-out and overall defoliation. Prior to mixing in the spray tank, a small scale compatibility test should be conducted.

Use only products which are exempt from tolerance under 40 CFR 180.1001.

Always follow the mixing instructions on the label of the appropriate adjuvant.

APPLICATION: Use the specified dosage of Klean-Pik 500SC in sufficient amount of water to give complete coverage of the foliage with uniform wetting of leaf surfaces. Apply in 10-25 gallons of spray per acre by ground equipment and 3-10 gallons per acre by air. Agitate the spray mix during application.

DOSAGE: One gallon of Klean-Pik 500SC treats 20 - 80 acres (0.05-0.2 lb. active ingredient/acre). Use the higher rates during periods of low temperatures (60°-65°F).

Two applications not exceeding a total of 9.6 ounces (0.3 lb. active ingredient/acre) may be necessary to defoliate rank cotton.

To Achieve an Application Rate of:	Use This Amount of Klean-Pik 500SC*	At the Indicated Rate, One Gallon of Klean-Pik 500SC Will Treat		
0.05 lb. ai/Acre	1.6 ounces/Acre	80 Acres		
0.10 lb. ai/Acre	3.2 ounces/Acre	40 Acres		
0.15 lb. ai/Acre	4.8 ounces/Acre	26.6 Acres		
0.20 lb. ai/Acre	6.4 ounces/Acre	20 Acres		

KLEAN-PIK 500SCAPPLICATION RATE TABLE

*Do not exceed a total of 9.6 ounces of Klean-Pik 500SC per acre per season.

BOTTOM DEFOLIATION FOR USE IN ARIZONA ONLY: Klean-Pik 500SC may be used as an aid to improve air circulation and help reduce boll rot caused by moisture.

Apply Klean-Pik 500SC at 1.6 ounces (0.05 lb. active ingredient/acre) formulated product per acre.

Application to bottom 1/3 of plant only recommended.

Timing of application should be based upon the latest date when ground application equipment can enter the field without crop damage.

PRECONDITIONING: Klean-Pik 500SC may be used as a pre-conditioner to enhance the activity of a defoliant application. Apply Klean-Pik 500SC at 1.6 – 3.2 ounces (0.05 – 0.1 lb. active ingredient/acre) formulated product per acre.

Timing of application is recommended to be 7 – 14 days prior to the use of another defoliant. Refer to the second product label prior to use for complete recommendations.

TANK MIX OF KLEAN-PIK 500SC PLUS FINISH® 6 PRO OR PREP™ FOR BOLL OPENING: The tank mix of Klean-Pik 500SC plus FINISH®6 PRO or Prep™ is recommended to improve overall defoliation, and as an aid in accelerating the opening of mature, unopened cotton bolls. Best activity will be obtained where the tank mix is applied to mature cotton plants. (See TIME OF APPLICATION SECTION of this label). Do not apply tank mix before sufficient unopened bolls have matured to produce the desired cotton yield. If nighttime temperatures are expected to fall below 60°F, unsatisfactory defoliation and regrowth inhibition may result. Certified cotton seed producers should determine boll and seed maturity prior to treatment.

Apply Klean-Pik500SC at a rate of 1.6 – 6.4 ounces (0.05-0.2 lb. active ingredient/acre) formulated product plus FINISH®6 PRO at 1.3 – 2 pt/A (1.0 – 1.5 lbs. active ingredient per acre) or Prep[™] at 1.3 – 2.6 pt/A (1.0- 2.0 lbs. active/ingredient per acre). The higher rate of Klean-Pik500SC should be used where excessive regrowth is anticipated, or during cooler temperatures.

TANK MIX OF KLEAN-PIK 500SCPLUS FINISH® 6PRO FOR DEFOLIATION ENHANCEMENT: Lower rates of FINISH® 6 PRO may be tank mixed with Klean-Pik 500SC to enhance defoliation without providing boll opening activity. FINISH® 6 PRO may still be applied as a sequential treatment for boll opening following an application of Klean-Pik 500SC plus FINISH® 6 PRO at a defoliation enhancement rate. Read and follow the FINISH® 6 PRO label, and do not exceed a maximum of 2.0 lb. ethephon a.i./acre for combined uses per acre per year.

Apply Klean-Pik 500SC at a rate of 1.6 - 6.4 ounces (0.05 - 0.2 lb. active ingredient/acre) formulated product plus FINISH @6 PRO at 5 ounces per acre (0.23 lb. active ingredient/acre). The higher rates of Klean-Pik 500SC should be used where excessive regrowth is anticipated, or during cooler temperatures.

TANK MIX OF KLEAN-PIK 500SC PLUS DEF® 6 OR KLEAN-PIK 500SC PLUS FOLEX® 6 EC: The tank mix of Klean Pik 500SC and Def® 6 or Folex® 6 EC is recommended to improve defoliation and inhibit regrowth under less than ideal conditions. Less than ideal conditions are those when the nighttime temperatures are expected to be 60°F on the Page 4 of 6

date of application and for three days following application.

Best activity will be obtained when the tank mix is applied to mature cotton plants with 60% or more open bolls. **RATE OF APPLICATION**

Klean-Pik 500SC + Def® 6 = 1.6 – 6.4 ounces Klean-Pik 500SC+ 0.5 – 2.0 pints Def® 6.

Klean-Pik 500SC+ Folex® 6 EC = 1.6 – 6.4 ounces Klean Pik 500SC + 0.5 – 2.0 pints Folex® 6 EC.

Higher rate of Klean-Pik 500SC should be used where excessive regrowth is anticipated.

Maximum rates of Klean-Pik 500SC and phosphate defoliants applied in a tank mix during very high air temperatures can cause desiccation.

Use lower rate of Def® 6 or Folex® 6 EC under ideal nighttime temperature conditions (65°F); higher rate during less than ideal or adverse conditions.

When applying Def® 6 or Folex® 6 EC defoliant plus Klean-Pik 500SC as a tank mix, the following mixing sequence is recommended:

1. Add required water to spray tank, agitate.

2. Add Klean-Pik 500SCaccording to label directions (see MIXING INSTRUCTIONS).

3. Add Def® 6 or Folex® 6 EC according to rate desired after the Klean Pik 500SC has completely dispersed.

A second application of the labeled tank mix may be made where necessary, but not exceeding a total of 0.3 lb. active ingredient of Klean-Pik 500 SC per acre.

Refer to Def® 6 and Folex® 6 EC labeling for additional use directions and cautions when using tank mixtures of Klean Pik 500SC and those products.

DO NOT USE TANK MIX OF KLEAN-PIK 500SC PLUS DEF® 6 OR FOLEX® 6 EC IN THE RIO GRANDE VALLEY COUNTIES OF STARR, HIDALGO, WILLACY, AND CAMERON.

IMPORTANT CLEANOUT INSTRUCTIONS

Tank mixes of Klean-Pik 500SC with organophosphates may form a residue in application equipment. While still fresh and moist, this residue can be removed by flushing the entire system with a commercial tank cleaner.

Do not allow the spray solution to dry in the application equipment. Immediately following application, clean all equipment (mix tanks, pumps, transfer lines, application tanks, sumps, booms, nozzles, and all related equipment) thoroughly with cleaner and water.

Should small quantities of Klean-Pik 500SC remain in inadequately cleaned equipment, they may be released during subsequent applications and may cause damage to crops. Makhteshim Agan of North America, Inc. accepts no liability for damage to crops due to inadequately cleaned equipment.

SEQUENTIAL APPLICATION

PRECAUTION: Do not apply sequential application of Klean-Pik 500SC following any defoliant or desiccant **except** Klean-Pik 500SCused alone or Klean-Pik 500SC tank mixes. Reduced activity of the second treatment will result.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store at temperatures below 100°F. If allowed to freeze, remix before using.

PESTICIDE DISPOSAL: Wastes resulting form the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING:

Nonrefillable Container (five gallons or less): Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Nonrefillable Container (greater than five gallons): Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Refillable Container: Refillable container. Refill this container with thidiazuron only. Do not reuse this container Page 5 of 6 for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

FOR 24-HOUR EMERGENCY ASSISTANCE (SPILL, LEAK, OR FIRE), CALL INFOTRAC AT 1-800-535-5053.

LIMITATION OF WARRANTY AND LIABILITY

Read the entire directions for use, conditions of warranties and limitations of liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following CONDITIONS, DISCLAIMER OF WARRANTIES and LIMITATIONS OF LIABILITY.

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Makhteshim Agan of North America, Inc. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: To the extent consistent with applicable law, Makhteshim Agan of North America, Inc. makes no other warranties, express or implied, of merchantability or of fitness for a particular purpose or otherwise, that extend beyond the statements made on this label. No agent of Makhteshim Agan of North America, Inc. is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. To the extent consistent with applicable law, Makhteshim Agan of North America, Inc. disclaims any liability whatsoever for special, incidental or consequential damages resulting from the use or handling of this product.

LIMITATIONS OF LIABILITY: To the extent consistent with applicable law, the exclusive remedy of the user or buyer for any and all losses, injuries or damages resulting from the use or handling of this product, whether in contract, warranty, tort, negligence, strict liability or otherwise, shall not exceed the purchase price paid or at Makhteshim Agan of North America, Inc.'s election, the replacement of product.

Klean-Pik[®] is a registered trademark of FarmSaver.com, LLC. DEF[®], Finish[®] 6 PRO and Prep[™] are trademarks of Bayer CropScience. FOLEX[®] 6EC is a trademark of AMVAC Chemical Corporation

Klean-Pik 500SC (66222-124)(EPA app 4-10-06)(notif to EPA 08-29-08)(notif 02-19-09)

NOTIFICATION FEB 2 3 2009

Page 6 of 6