

66222-136

6.26.2007

1/12



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

Anne Stout
Registration Specialist
Makhteshim Agan of North America, Inc.
4515 Falls of Neuse Rd., Suite 300
Raleigh, NC 27609

JUN 26 2007

SUBJECT: Application for Pesticide Notification – Updated Warranty Statement
Ethephon 6SC
EPA Reg. No. 66222-136
Application Dated April 30, 2007

Dear Ms. Stout:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 for the above product. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the actions requested fall within the scope of PRN 98-10. The label submitted with the application has been stamped "Notification" and will be placed in our records.

If you have any questions, please me directly at 703-305-6249 or Terri Stowe of my staff at 703-305-6117.

Sincerely,

A handwritten signature in black ink, appearing to read "Linda Arrington".

Linda Arrington
Notifications & Minor Formulations Team Leader
Registration Division (7505P)
Office of Pesticide Programs



United States
Environmental Protection Agency
 Washington, DC 20460

<input type="checkbox"/>	Registration
<input type="checkbox"/>	Amendment
<input checked="" type="checkbox"/>	Other

OPP Identifier Number

Application for Pesticide - Section I

1. Company/Product Number 66222-136	2. EPA Product Manager Dennis McNeilly	3. Proposed Classification <input type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Ethephon 6SC	PM# 22	
5. Name and Address of Applicant (Include ZIP Code) Makhteshim Agan of North America, Inc. 4515 Falls of Neuse Rd., Suite 300 Raleigh, NC 27609 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

Section - II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.
<input type="checkbox"/> Notification - Explain below.	<input checked="" type="checkbox"/> Other - Explain below.

NOTIFICATION
JUN 26 2007

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

Notification of updated warranty statement per EPA guidance issued 10-17-06 and PRN 98-10 Section II(J)
 This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula for this product. I understand that it is a violation of 18 USC Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under Sections 12 and 14 of FIFRA.

Section - III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No		<input type="checkbox"/> Metal	<input type="checkbox"/> Plastic
* Certification must be submitted		If "Yes" Unit Packaging wgt.	No. per container	If "Yes" Package wgt	No. per container
				<input type="checkbox"/> Glass	<input type="checkbox"/> Paper
				<input type="checkbox"/> Other (Specify) _____	
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container		5. Location of Label Directions <input type="checkbox"/>	
6. Manner in Which Label is Affixed to Product <input type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled			<input type="checkbox"/> Other _____		

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name Anne Stout	Title Registration Specialist	Telephone No. (Include Area Code) 901-861-4400
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.		6. Date Application Received (Stamped)
2. Signature <i>Anne Stout</i>	3. Title Registration Specialist	
4. Typed Name Anne Stout	5. Date 4-30-07	

PRECAUTIONARY STATEMENTS

DANGER

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Corrosive. Causes irreversible eye damage and skin burns. Harmful if swallowed or absorbed through skin. Do not get in eyes, on skin, or on clothing.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Coveralls over short-sleeved shirts and short pants
- Chemical-resistant gloves made of any waterproof material such as nitrile, butyl, neoprene and/or barrier laminate
- Chemical-resistant footwear plus socks
- Protective eyewear
- Chemical-resistant headgear for overhead exposure
- Chemical-resistant apron when mixing, loading or cleaning equipment

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Do not contaminate water used for irrigation or domestic purposes.

SPRAY DRIFT

Avoid spray drift. Do not apply when weather conditions may cause drift. Do not allow this product to drift on to non-target areas. Drift may result in illegal residues or injury to adjacent crops and vegetation, in the form of leaf yellowing and defoliation. To avoid spray drift, DO NOT apply aerially when wind speed is greater than 10 mph or during periods of temperature inversions. Use of larger droplet size will also reduce spray drift.

AVOIDING SPRAY DRIFT AT THE APPLICATION SITE IS THE RESPONSIBILITY OF THE APPLICATOR.

The interaction of many equipment- and-weather-related factors determine the potential for spray drift. The applicator is responsible for considering all these factors when making decisions.

The following drift management requirements must be followed to avoid off-target movement from aerial applications to agricultural field crops. These requirements do not apply to forestry applications, public health uses, or to applications using dry formulations.

1. The distance of the outer most nozzles on the boom must not exceed 3/4 the length of the wingspan or rotor.

2. Nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees.

Where states have more stringent regulations, they should be observed.

The applicator should be familiar with and take into account the information covered in the **AERIAL DRIFT REDUCTION ADVISORY** below.

AERIAL DRIFT REDUCTION ADVISORY

This section is advisory in nature and does not supersede the mandatory label requirements.

Information on Droplet Size

The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential but will not prevent drift if applications are made improperly or under unfavorable environmental conditions (see **Wind, Temperature and Humidity, and Temperature Inversions.**)

Controlling Droplet Size

Volume – Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.

Pressure – Do not exceed the nozzle manufacturer's recommended pressures. For many nozzle types, lower pressure produces larger droplets. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.

Number of Nozzles- Use the minimum number of nozzles that provide uniform coverage.

Nozzle Orientation- Orienting nozzles so that the spray is released parallel to the airstream produces larger droplets than other orientations and is the recommended practice. Significant deflection from horizontal will reduce droplet size and increase drift potential.

Nozzle Type- Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce the largest droplets and the lowest drift.

Boom Length

For some use patterns, reducing the effective boom length to less than ¾ of the wingspan or rotor length may further reduce drift without reducing swath width.

Application Height

Applications should be not be made at a height greater than 10 feet above the top of the target plants unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind.

Swath Adjustment

When applications are made with a crosswind, the swath will be displaced downwind. Therefore, on the up and downwind edges of the field, the applicator should compensate for this displacement by adjusting the path of the aircraft upwind. Swath adjustment distance should increase with increasing drift potential (higher wind, smaller drops, etc.).

Wind

Drift potential is lowest between wind speeds of 2 – 10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given speed. Application should be avoided below 2 mph due to variable wind direction and high inversion potential. NOTE: Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

Temperature and Humidity

When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

Temperature Inversions

Applications should not occur during a temperature inversion because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

Sensitive Areas

The pesticide should only be applied when the potential for drift to adjacent sensitive areas (e.g. residential areas, bodies of water, known habitat for threatened or endangered species, non-target crops) is minimal (e.g. when wind is blowing away from the sensitive areas).

USE PRECAUTIONS

DO NOT apply Setup 6SL through any type of irrigation system.

Avoid spray drift to nearby crops as this product will cause modifications in plant growth. Plant injury or reduced yields will result.

Mix only the amount of spray you expect to use each day. DO NOT allow mixed solutions to stand overnight.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State of Tribe, consult the agency responsible for pesticide regulations.

Read entire label before using this product.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers, and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 48 hours. The REI is 72 hours in areas where average rainfall is less than 25 inches per year.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated such as plants, soil or water is:

- Coveralls over long-sleeved shirt and long pants
- Chemical-resistant gloves made of any waterproof material, such as nitrile, butyl, neoprene or barrier laminate
- Chemical-resistant footwear plus socks
- Protective eyewear

• For overhead exposure, chemical-resistant headgear is also required.
Notify workers of the application by warning them orally and posting warning signs at entrances to treated areas.

GENERAL INFORMATION

Cotton

Setup 6SL will accelerate opening of mature unopened cotton bolls and enhance defoliation which can result in earlier harvest with an increased recoverable yield. Treating cotton with Setup 6SL allows increased efficiency from a once-over harvest.

Tobacco (Flue-Cured)

Setup 6SL promotes early, uniform "yellowing" of mature tobacco. Setup 6SL reduces curing time, allows more efficient use of curing barn space, and increases control over harvest schedules.

Spray Preparation

Add 1/2 to 3/4 of the required amount of water to the spray tank. Start agitation. Add the required amount of product, then the remaining amount of water. Prepare only as much spray solution as can be used on the day of mixing. Do not allow spray solution to stand overnight. Do not spill the concentrated product on spray equipment or any airplane parts.

ANY SPILLS SHOULD BE RINSED IMMEDIATELY WITH PLENTY OF WATER.

Use of a nurse tank is highly recommended for avoiding possible spills of concentrated formulation on spray equipment or any airplane parts.

Tank Mixtures with Defoliants and Insecticides

Setup 6SL is compatible with DEF¹, FOLEX², Dropp³, Dropp® Ultra^{TM3}, Ginstar^{®3}, Harvade⁴, Methyl Parathion, Guthion[®] and malathion. Follow all applicable use precautions and rate recommendations on labels of products applied as tank mixtures or in sequence with this product. In some cases, slight reduction in boll opening response has been observed when tank mixes with defoliants were used.

Good agitation in the spray tank is essential. DO NOT allow tank mixtures to stand without agitation for more than 5 to 10 minutes. Read and observe all appropriate label use directions and precautions for the defoliants and insecticides used.


- DO NOT tank mix Setup 6SL with desiccants if cotton is to be spindle harvested.
- DO NOT tank mix Setup 6SL with products containing sodium chlorate, since doing so may result in the formation of Hypochlorous Acid that will release toxic chlorine fumes upon heating.
- DO NOT tank mix Setup 6SL with ammonium thiosulfate, since doing so may result in the release of toxic fumes.

Equipment Cleaning

Because of the acidic nature of this product, prolonged exposure to spray deposit will damage acrylic plastics, certain paints, and metals.

Thoroughly rinse all exposed acrylic plastic-type materials (e.g. aircraft windshields) and painted surfaces with detergent and water within one hour after exposure to spray deposits.

At the end of each day, thoroughly rinse all the metal parts of the spray equipment exposed to the spray deposits with detergent and water.



COTTON

Use	Conditions	Rate of Setup 6SL		One Gallon of Setup 6SL Treats:	Minimum Spray Volumes (Gals/A) ^a		Application Timing
		Pints/A	Lbs. A.I.	Acres	Ground	Air ^d	
Boll Opener ^b	Hot, dry, above 80°F	1 1/3	1.0	6	10	2	Apply when the number of mature unopened bolls is sufficient to produce the desired crop. See Boll Maturity section below for test of boll maturity. Treatment uniformly opens bolls 7 to 14 days earlier.
	Dry and 75-80°F	2	1.5	4			
	Cool, above 65° or Rank Cotton	2 2/3	2.0	3			
Setup 6SL + FOLEX [®] Defoliant Tank Mix ^c	High soil moisture or High fertility level or Rank Cotton	1/3	0.25	24	10	5	Apply 4 to 7 days prior to boll opening treatment. Use as a sequential treatment with, not in place of, boll opening treatment.
Setup 6SL + Dropp Defoliant Tank Mix ^c	High soil moisture or High fertility level or Rank Cotton	1/3	0.25	24	10	3	Apply 4 to 7 days prior to boll opening treatment. Use as a sequential treatment with, not in place of, boll opening treatment.
Pre-Conditioner for Defoliation	Hot, dry, above 80°F	2/3	0.5	12	10	2	Apply 4 to 7 days prior to defoliant. Enhances top crop defoliation, reduces deterioration of bottom crop and allows for earlier harvest.
	Cool, above 65°F or Rank Cotton	1 1/3	1.0	6			

^a For best performance, whether by ground or air application, choose equipment and spray voluínés that will ensure uniform coverage of foliage and bolls.

^b Pretreatment with defoliant before boll opening treatment with Setup 6SL: If the cotton is overly rank or laying down in the middles and good spray coverage of the bolls is difficult, pretreatment with a defoliant will

often improve boll coverage in subsequent boll opening treatment with Setup 6SL at recommended rates (above). Read and observe all appropriate label use directions and precautions for the defoliant used.

Can use up to full label rate for each product.

In California and Arizona, use a volume of no less than 5 gallons per acre for aerial applications.

Boll Maturity

A cotton boll is mature when it is too hard to be dented when squeezed between thumb and fingers, too hard to be sliced with a sharp knife, and when the seed coat becomes light brown in color.

Use Limitations

- **Boll Opening:** DO NOT tank mix Setup 6SL with a desiccant if the cotton is to be spindle harvested.
- **Pre-Condition for Defoliation:** DO NOT tank mix Setup 6SL with desiccants unless plant desiccation is required. DO NOT use a defoliant before there are sufficient mature unopened bolls to produce the desired yield (see **Boll Maturity** section, above, about how to test for boll maturity).
- DO NOT apply this product if rain is expected within 6 hours. Rainfall within 6 hours of application may reduce product performance.
- DO NOT harvest cotton within 7 days after treatment.
- DO NOT plant another crop within 30 days after treatment. Small grains planted earlier than 1 month or intercropped with the cotton crop to which Setup 6SL will be applied may only be used as cover crops and may not be harvested for food or feed. Setup 6SL may cause yellowing and growth inhibition of treated small grains.
- DO NOT exceed a maximum of 2.0 lbs. ethephon active ingredient per acre per year through combined or repeated uses of any ethephon products.

Observe the treated crop and harvest when optimum boll opening has been attained. Harvesting too early might reduce the full advantage of the treatment and too late may result in reduced quality and loss of lint which will drop from the plant.

**TOBACCO
(Flue-Cured Only)**

Application Method	Rate of Setup 6SL Pints/Acre	Minimum Spray Volumes (Gals/A)	Specific Directions
Directed Spray	1 1/3	50	Use drop nozzles. Choose TG or OC spray tips designed to apply 50 – 60 GPA at 35 – 40 psi and tractor speed of 2-3 mph. Use 2 nozzles per row; one on each side of the row dropped low enough to direct the spray onto the leaves to be ripened for harvest. Thorough spray coverage is essential. Harvest all leaves with 20% or more yellowing.
Over-the-Top	1 1/3 – 2 2/3	40	Treat when only mature leaves remain on the stalk. To determine whether remaining leaves are mature, test spray several tobacco plants as described below in the Application Timing section. Use the lower rate under most conditions when experience indicates that minimum ripening inducement is required. Use the higher rate when the crop is heavy or rank or during cool (temperatures below 65°F), slow ripening conditions. Always test spray to determine if the tobacco is mature enough to respond to treatment with Setup 6SL. Apply over-the-top sprays as a fine mist using three nozzles (one nozzle tip over the center of the plant, and one on each side) so all leaves are covered thoroughly, similar to application methods for

systemic sucker control agents. Use a spray pressure of 40 to 60 psi.

Use Limitations

- DO NOT use Setup 6SL with additives other than those recommended on this label.
- DO NOT apply Setup 6SL to immature leaves, since this can result in unsatisfactory coloring, weight loss, and reduced leaf quality.
- DO NOT allow the crop to over ripen in the field after using Setup 6SL, since this may cause some reduction in yield and quality.
- DO NOT apply this product if rain is expected within 6 hours. Rainfall within 6 hours of application may reduce product performance.
- DO NOT treat before anticipated heavy rainstorms that could prevent harvest and result in crop loss.
- DO NOT plant another crop within 30 days after treatment.
- DO NOT exceed a maximum of 2 lbs. ethephon active ingredient per acre per year through repeated uses of any ethephon products.

Application Timing

Successful results with Setup 6SL require treatment when leaves are mature, not overly rank green. To determine the proper treatment timing and the number of leaves per stalk ready for harvest, test spray several plants in more than one location in each field and observe the response. Mature leaves will begin to yellow in 24 to 72 hours. Test leaves that do not yellow within 72 hours are not mature and not ready for treatment with Setup 6SL. Wait a few days to permit further natural maturing, then make another test spray to assess crop maturity.

Prepare test spray by mixing 1 teaspoonful of Setup 6SL in 1 quart of water. Spray each test plant with about 1 oz. of this mixture, covering all leaves with a fine mist. Setup 6SL will not color immature sprayed leaves.

Once the proper treatment timing has been determined, one can determine the number of acres to treat in order to fill the curing barn.

When to Harvest

Mature, treated leaves will begin to color within 24 to 72 hours after treatment with Setup 6SL. The yellowing process is weather dependent; cool weather will delay, while hot, sunny weather can accelerate the process. Harvest treated tobacco when leaves have achieved the desired color intensity.

Harvest can begin 48 hours after treatment. Closely monitor treated crop and weather conditions to determine harvest timing and avoid quality loss or leaf drop.

Curing Treated Tobacco

Curing procedures depend on crop condition, interval between treatment and harvest, weather, and type of curing facility. To obtain maximum quality, observe and control the curing process closely, especially during the late "coloring" and early "drying" stages of the leaves.

Tobacco treated with Setup 6SL will have begun the coloring process when harvested and will reduce the time required in the coloring phase of curing. Treated tobacco should be dried faster. If tobacco leaves are green or contain some green when harvested, it may be necessary to color them for a few hours. If the leaves are completely yellow, temperature and ventilation must be adjusted to dry the tobacco as fast as possible without scalding. Once the leaf is dried (3/4 dry), follow normal curing procedures. Since tobacco treated with Setup 6SL will cure faster than untreated tobacco, treated and untreated tobacco should not be cured together in the same barn.

STORAGE AND DISPOSAL
Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store in a cool, dry place and away from food, feed and other pesticides. **IF SPILLED:** If container is broken or contents have spilled, follow all precautions indicated above and clean up immediately. Before cleaning up, put on full-length trousers, long-sleeved shirt, protective gloves, and goggles or face shield. Soak up spill with absorbent media such as sand, earth, or other suitable material and dispose of waste at an approved waste disposal facility. In the event of a major spill, fire, or other emergency, call Infotrac at 1-800-535-5053, day or night.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Triple rinse (or equivalent) the empty containers. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or by incineration or, if allowed by state and local authorities, by burning. If container is burned, stay out of smoke.

IMPORTANT: Read the entire **DIRECTIONS FOR USE** and the **LIMITATION OF WARRANTY AND LIABILITY** statement before using this product. If terms are not acceptable, return the unopened product container at once.

LIMITATION OF WARRANTY AND LIABILITY

Read the entire Directions for Use, Conditions of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Makhteshim Agan of North America, Inc. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: To the extent consistent with applicable law, Makhteshim Agan of North America, Inc. makes no other warranties, express or implied, of merchantability or of fitness for a particular purpose or otherwise, that extend beyond the statements made on this label. No agent of Makhteshim Agan of North America, Inc. is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. To the extent consistent with applicable law, Makhteshim Agan of North America, Inc. disclaims any liability whatsoever for special, incidental or consequential damages resulting from the use or handling of this product.

LIMITATIONS OF LIABILITY: To the extent consistent with applicable law, the exclusive remedy of the user or buyer for any and all losses, injuries or damages resulting from the use or handling of this product, whether in contract, warranty, tort, negligence, strict liability or otherwise, shall not exceed the purchase price paid or at Makhteshim Agan of North America, Inc.'s election, the replacement of product.

Setup is a trademark of MAKHTESHIM AGAN OF NORTH AMERICA, INC.

- 1 DEF and Guthion are registered trademarks of Bayer Crop Science.
- 2 FOLEX is a registered trademark of the AMVAC Chemical Corporation.
- 3 Dropp and Ginstar are registered trademarks and Ultra is a trademark of Bayer Crop Science.
- 4 Harvade is a registered trademark of Crompton/Uniroyal Co.

