

66222-156

01-18-2012

1/29



UNITED STATES ENVIRONMENTAL PROTECTION  
AGENCY  
WASHINGTON, D.C. 20460

**NOTIFICATION**

**JAN 18 2012**

OFFICE OF CHEMICAL SAFETY  
AND POLLUTION PREVENTION

Kristen Knox  
Makhteshim Agan of North America, Inc.  
4515 Falls of Neuse Rd., Suite 300  
Raleigh, NC 27609

Subject: Notification to delete almond from product label per CDPR instructions  
EPA Registration No. 66222-156  
Primary Brand Name: MANA ALIAS 4F  
Submission Date: January 6, 2012

Dear Ms. Knox:

The Agency is in receipt of your Application for Pesticide Notification under PRN 98-10 dated January 6, 2012 for products EPA Reg No. 66222-156 and finds that the action requested falls within the scope of PRN 98-10. The label submitted with the application has been stamped "Notification" and will be placed in our records. If you have any questions, please contact Gene Benbow at (703) 347-0235 or via email at [benbow.gene@epa.gov](mailto:benbow.gene@epa.gov).

Sincerely,

A handwritten signature in black ink, appearing to read "Gene Benbow".

Gene Benbow  
Wildlife Biologist  
Insecticide-Rodenticide Branch  
Registration Division (7505P)

Please read instructions on reverse before completing form.

Form Approved. OMB No. 2070-0060, Approval expires 2-28-95



United States  
Environmental Protection Agency  
Washington, DC 20460

<input type="checkbox"/>	Registration
<input type="checkbox"/>	Amendment
<input checked="" type="checkbox"/>	Other

OPP Identifier Number

### Application for Pesticide - Section I

1. Company/Product Number 66222-156	2. EPA Product Manager Venus Eagle	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) MANA ALIAS 4F	PM# 01	
5. Name and Address of Applicant (Include ZIP Code) Makhteshim Agan of North America, Inc. 4515 Falls of Neuse Rd., Suite 300 Raleigh, NC 27609 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

### Section - II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____	<b>NOTIFICATION</b> JAN 18 2012
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.	
<input type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.	

**Explanation:** Use additional page(s) if necessary. (For section I and Section II.)

Notification of deletion of a crop; almonds, per CDPR instructions  
This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula for this product. I understand that it is a violation of 18 USC Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under Sections 12 and 14 of FIFRA.

### Section - III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No		<input type="checkbox"/> Metal	<input type="checkbox"/> Plastic
* Certification must be submitted		If "Yes" Unit Packaging wgt.	No. per container	If "Yes" Package wgt	No. per container
				<input type="checkbox"/> Glass	<input type="checkbox"/> Paper
				<input type="checkbox"/> Other (Specify) _____	
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container		5. Location of Label Directions <input type="checkbox"/>	
6. Manner in Which Label is Affixed to Product <input type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled		<input type="checkbox"/> Other _____			

### Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name Kristen Knox	Title Product Registration Manager	Telephone No. (include Area Code) 919-256-9337
<b>Certification</b> I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.		6. Date Application Received (Stamped)
2. Signature 	3. Title Product Registration Manager	
4. Typed Name Kristen Knox	5. Date 1-6-12	

3  
29



**M A N A**

January 6, 2012

Document Processing Desk (NOTIF)  
Office of Pesticide Programs (7504P)  
U.S. Environmental Protection Agency  
Room S-4900, One Potomac Yard  
2777 S. Crystal Drive  
Arlington, VA 22202

Attn: Ms. Venus Eagle  
Insecticide-Rodenticide Branch (Team 01)

Re: MANA Alias 4F, EPA Reg. No. 66222-156  
Quali-Pro Imidacloprid 2F, EPA Reg. No. 66222-203  
Pasada 1.6F Flowable Insecticide, EPA Reg. No. 66222-228  
Notification of the Deletion of a crop per PRN 98-10

To Whom It May Concern:

Makhteshim Agan of North America, Inc. is notifying the Agency of its intention to delete almonds from the list of tree crops of the referenced product labels, as allowed in PRN 98-10. As a result of California Notice 2009-2, Notice of Decision to Initiate Reevaluation of Chemicals in the Nitroguanidine Insecticide Class of Neonicotinoids, we have agreed to delete almonds from the labels of all registered imidacloprid products.

We certify the following:

This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula for this product. I understand that it is a violation of 18 USC Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under Sections 12 and 14 of FIFRA.

In support of this submission, the following documents are attached:

- Application for registration (EPA Form 8570-1)
- One copy of labeling with annotations marked in red
- One copy of "clean" labeling

Should you have any questions regarding this submission, please contact me at 919-256-9337 or by email at [kknox@manainc.com](mailto:kknox@manainc.com).

Sincerely,

Kristen B. Knox  
Product Registration Manager

**4515 Falls of Neuse Road, Suite 300, Raleigh NC 27609**  
**T: 919.256.9305 • F: 919.256.9308**

4  
29

**NOTIFICATION**

**ANNOTATED**

JAN 18 2012

**MANA ALIAS® 4F**  
**Flowable Insecticide**

<b>ACTIVE INGREDIENT</b>	<b>% BY WT.</b>
Imidacloprid; 1-[(6-Chloro-3-pyridinyl)methyl]-N-nitro-2-imidazolidinimine.....	40.6%
<b>OTHER INGREDIENTS:</b> .....	59.4%
	<b>TOTAL 100.0%</b>

Contains 4 lbs. of active ingredient per gallon

**KEEP OUT OF REACH OF CHILDREN**  
**CAUTION**

Manufactured for:  
Makhteshim Agan of North America, Inc.  
4515 Falls of Neuse Rd., Suite 300  
Raleigh, NC 27609

EPA Reg. No. 66222-156

EPA Est. No. [REDACTED]

**NET CONTENTS: \_\_\_\_\_ GALLON**

**FIRST AID**

<b>IF SWALLOWED:</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by a poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
<b>IF ON SKIN OR CLOTHING:</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>IF IN EYES:</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>IF INHALED:</b>	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact PROSAR at 1-877-250-9291 for emergency medical treatment information.

**NOTE TO PHYSICIAN:** No specific antidote is available. Treat the patient symptomatically.

**PRECAUTIONARY STATEMENTS**  
**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**  
**CAUTION**

Harmful if swallowed, absorbed through skin, or inhaled. Avoid contact with skin, eyes, or clothing. Avoid breathing vapor or spray mist. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.


**PERSONAL PROTECTIVE EQUIPMENT (PPE)**

**Applicators and other handlers must wear:**

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of any waterproof material such as barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, natural rubber, polyethylene, polyvinylchloride (PVC), or Viton
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining personal protective equipment, PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

5  
19



6  
29

### ENGINEERING CONTROLS STATEMENT

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

### USER SAFETY RECOMMENDATIONS

**Users should:**

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

### ENVIRONMENTAL HAZARDS

Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area. This product is toxic to wildlife and highly toxic to aquatic invertebrates.

This chemical demonstrates the properties and characteristics associated with chemicals detected in ground water. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground water contamination.

**OBSERVE THE FOLLOWING PRECAUTIONS WHEN MIXING AND APPLYING IN THE VICINITY OF AQUATIC AREAS SUCH AS LAKES, RESERVOIRS, RIVERS, PERMANENT STREAMS, MARSHES OR NATURAL PONDS, ESTUARIES, AND COMMERCIAL FISH FARM PONDS.**

### SPRAY DRIFT MANAGEMENT

The interaction of many equipment- and weather-related factors determine the potential for spray drift. The applicator is responsible for considering all of these factors when making application decisions. Avoiding spray drift is the responsibility of the applicator.

**For Aerial Applications**

For aerial applications, the spray boom should be mounted on the aircraft so as to minimize drift caused by wing tip vortices. The minimum practical boom length should be used, and must not exceed 75% of the wing span or rotor diameter. Spray should be released at the lowest possible height consistent with good pest control and flight safety. Applications more than 10 feet above the crop canopy should be avoided.

**Importance of Droplet Size**

An important factor influencing drift is droplet size. Small droplets (<150-200 microns) drift to a greater extent than large droplets. Within typical equipment specifications, applications should be made to deliver the largest droplet spectrum that provides sufficient control and coverage. Formation of very small droplets may be minimized by appropriate nozzle selection, by orienting nozzles away from the airstream as much as possible and by avoiding excessive spray boom pressure.

**Wind Speed Restrictions**

Drift potential increases at wind speeds of less than 3 mph (due to inversion potential) or more than 10 mph. However, many factors, including droplet size, canopy, and equipment specifications determine drift potential at any given wind speed. Do not apply when winds are greater than 15 mph and avoid gusty and windless conditions. Risk of exposure to sensitive aquatic areas can be reduced by avoiding applications when wind direction is toward the aquatic area.

**Restrictions During Temperature Inversions**

Do not make aerial or ground applications during temperature inversions. Drift potential is high during temperature inversions. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain close to the ground and move laterally in a concentrated cloud. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical mixing.

**Mixing and Loading Requirements**

To avoid potential contamination of groundwater, the use of a properly designed and maintained containment pad for mixing and loading of any pesticide into application equipment is recommended. If containment pad is not used, maintain a minimum distance of 25 feet between mixing and loading area and potential surface to groundwater conduits such as field ditches, uncased well heads, sinkholes, or field drains.

**Airblast (Air Assist) Specific Recommendations for Tree Crops and Vineyards**

Airblast sprayers carry droplets into the canopy of trees/vines via a radially, or laterally directed air stream. The following specific drift management practices should be followed:

- Adjust deflectors and aiming devices so that spray is only directed into the canopy.
- Block off upward pointed nozzles when there is no overhanging canopy.
- Use only enough air volume to penetrate the canopy and provide good coverage.
- Do not allow the spray to go beyond the edge of the cultivated area (i.e., turn off sprayer when turning at end rows).

7  
29

- Only spray inward, toward the orchard or vineyard, for application to the outside rows.

**No-Spray Zone Requirements for Soil and Foliar Applications**

Do not apply by ground within 25 feet or by air within 150 feet of lakes, reservoirs, rivers, permanent streams, marshes or natural ponds, estuaries, and commercial fish farm ponds.

**Runoff Management**

Do not cultivate within 10 feet of the aquatic areas to allow growth of a vegetative filter strip. When using MANA Alias 4F on erodible soils, Best Management Practice for minimizing runoff should be employed. Consult your local Natural Resources Conservation Service for recommendations in your use area.

**Endangered Species Notice**

Under the Endangered Species Act, it is a federal offense to use any pesticide in a manner that results in the death of a member of an endangered species. Consult your local county bulletin, County Extension Agent, or Pesticide State Lead Agency for information concerning endangered species in your area.

**Resistance Management**

Some insects are known to develop resistance to insecticides after repeated use. As with any insecticide, the use of this product should conform to resistance management strategies established for the use area.

MANA Alias® 4F contains a Group 4A insecticide. Insect biotypes with acquired or inherent tolerance to Group 4A insecticides may eventually dominate the insect population if Group 4A insecticides are used repeatedly as the predominant method of control for targeted species. This may eventually result in partial or total loss of control of those species by MANA Alias 4F and to other Group 4A insecticides.

The active ingredient in MANA Alias 4F is a member of the neonicotinoid chemical class. Insect pests resistant to other chemical classes have not shown cross-resistance to MANA Alias 4F. In order to maintain susceptibility to this class of chemistry in insect species with high resistance development potential, for each crop season: 1) make only a single, soil application of MANA Alias 4F; 2) foliar applications of products from the same class not be made following a long residual, soil application of MANA Alias 4F, or other neonicotinoid products.

If a soil application of MANA Alias 4F has not been made during a crop season and foliar applications are to be made, avoid using a block of more than three consecutive applications of MANA Alias 4F and/or other Group 4A products having the same or similar mode of action. Following a neonicotinoid block of treatments, MANA strongly encourages the rotation to a block of applications with effective products with a different mode of action before using additional applications of neonicotinoid products. Using a block rotation or windowed approach, along with other IPM practices, is considered an effective use strategy for preventing or delaying an insect's ability to develop resistance to this class of chemistry.

Do not use MANA Alias 4F or other Group 4A products from the neonicotinoid chemical class for foliar applications on crops previously treated with long-residual, soil-applied products from the neonicotinoid chemical class.

Other Group 4A, neonicotinoid products, used as foliar treatments include: Actara, Assail, Calypso, Centric, Clutch, Couraze, Galian, Impulse, Intruder, Leverage, Nuprid, Pasada, Provado, Trimax Pro, and Venom.

Other Group 4A, neonicotinoid products, used as soil/seed treatments include Admire Pro, Advise, Alias, Belay, Couraze, Cruiser, Gaucho, Macho, Macho Max, Nuprid, Platinum, Venom, and Widow.

Contact your Cooperative Extension specialist, certified crop advisor, and/or product manufacturer for additional insect resistance management recommendations. Also, for more information on Insect Resistance Management (IRM), visit the Insecticide Resistance Action Committee (IRAC) on the web at <http://www.irac-online.org/>.

**DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the Agency responsible for pesticide regulation.

**AGRICULTURAL USE REQUIREMENTS**

Use this product in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours. Exception: If the product is soil injected or soil incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Chemical resistant gloves made of any waterproof material such as barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, natural rubber, polyethylene, polyvinylchloride (PVC), or Viton
- Shoes plus socks

8  
29

### APPLICATION DIRECTIONS

For soil applications of MANA Alias 4F, direct product into the seed or root-zone of crop. Failure to place MANA Alias 4F into root-zone may result in loss of control or delay in onset of activity. Apply MANA Alias 4F with ground or chemigation application equipment.

Do not apply MANA Alias 4F in enclosed structures such as planthouses or greenhouses except as specifically instructed in the **TOBACCO, CUCURBIT VEGETABLES, FRUITING VEGETABLES** and **GREENHOUSE VEGETABLES**, (Mature plants in production greenhouses): Cucumber, Tomato only sections of this label.

Apply foliar applications of MANA Alias 4F as a directed or a broadcast foliar spray. Thorough coverage of foliage is necessary without runoff for optimum insecticidal efficacy. Use adequate spray volumes, properly calibrated application equipment, and spray adjuvant if necessary to obtain thorough coverage. Failure to provide adequate coverage and retention of MANA Alias 4F on leaves and fruit may result in loss of insect control or delay in onset of activity. Apply MANA Alias 4F with properly calibrated ground or aerial application equipment. Minimum spray volumes, unless otherwise specified on crop specific application sections, are 10 gallons per acre by ground and 5 gallons per acre by air. MANA Alias 4F may also be applied by overhead chemigation (see additional information in **CHEMIGATION** section of this label below) if allowed in crop specific application section.

When applied as a soil application, optimum activity of MANA Alias 4F results from applications to the root-zone of plants to be protected. The earlier MANA Alias 4F is available to a developing plant, the earlier the protection begins. MANA Alias 4F is continuously taken into the roots over a long period of time and the systemic nature of MANA Alias 4F allows movement from roots through the xylem tissue to all vegetative parts of the plant. This results in extended residual activity of MANA Alias 4F, the control of insects, and the prevention and/or reduction of virus transmission or symptom expression, and plant health benefits. The rate of MANA Alias 4F applied affects the length of the plant protection period. Use higher rates when infestations occur later in crop development or where pest pressure is continuous. MANA Alias 4F will generally not control insects infesting flowers, blooms, or fruit. Additional crop protection may be required for insects feeding in, or on these plant parts, and for insects not listed in the crop-specific, pests-controlled sections of this label. Additionally, specific MANA Alias 4F application instructions are also provided in the crop-specific sections of this label.

Suppression, or less than complete control of certain insect pests that may carry diseases including reduced feeding, may also result from a MANA Alias 4F application. Complete control of these pests may require supplemental control measures.

Generally MANA Alias 4F is not used on crops grown for production of true seed intended for private or commercial planting but may be allowed under state specific, supplemental labeling. As with any insecticide, minimize exposure of MANA Alias 4F to honey bees and other pollinators. Do not use MANA Alias 4F on crops requiring bee pollination during bloom and a minimum of 10 days prior to bloom. Additional information on MANA Alias 4F uses for these crops and other questions may be obtained from the Cooperative Extension Service, PCAs, consultants, or local Makhteshim Agan of North America, Inc. representatives.

Apply only to plants grown in field-type soils, potting media, or mixtures thereof. Do not apply to plants grown in non-soil medias such as perlite, vermiculite, rock wool, or other soil-less media, or plants growing hydroponically. Pre-mix MANA Alias 4F with water or other appropriate diluent prior to application. Keep MANA Alias 4F and water suspension agitated to avoid settling.

Do not apply more than 0.5 lb. active ingredient per acre per crop season regardless of formulation or method of application, unless specified within a crop-specific application section for a given crop.

### MIXING INSTRUCTIONS

To prepare the application mixture, add a portion of the required amount of water to the spray tank and with agitation, add MANA Alias 4F. Complete filling tank with balance of water needed. Maintain sufficient agitation during both mixing and application. MANA Alias 4F may also be used with other pesticides and/or fertilizer solutions. Please see **Compatibility** section of this label. When tank mixtures of MANA Alias 4F and other pesticides are involved, prepare the tank mixture as instructed above and follow suggested **Mixing Order** below.

#### Mixing Order

When pesticide mixtures are needed, add wettable powders or wettable granules first, MANA Alias 4F and other suspension concentrate (flowable) products second, and emulsifiable concentrates last. Ensure good agitation as each component is added. Do not add an additional component until the previous is thoroughly mixed. If a fertilizer solution is added, a fertilizer/pesticide compatibility agent may be needed. Maintain constant agitation during both mixing and application to ensure uniformity of spray mixture.

#### Compatibility

Test compatibility of the intended mixture before adding MANA Alias 4F to the spray or mix tank. Add proportionate amounts of each ingredient in the appropriate order to a pint or quart jar, cap, shake for 5 minutes, and let set for 5 minutes. Do not use if poor mixing or formation of precipitates that do not readily redisperse occurs which indicates an incompatible mixture.

### CHEMIGATION

**Types of Irrigation Systems:** Make foliar chemigation applications of MANA Alias 4F to crops through overhead sprinkler systems if specified in crop-specific application sections. Make soil chemigation applications of MANA Alias 4F only to crops through chemigation as specified in crop-specific application sections and only through low-pressure systems specifically listed for a given crop. Do not apply MANA Alias 4F through any other type of irrigation system.



Make foliar chemigation applications of MANA Alias 4F as concentrated as possible. Retention of MANA Alias 4F on target site of insect infestation is necessary for optimum activity. Do not chemigate MANA Alias 4F in water volumes exceeding 0.10 inches per acre. See crop-specific application sections of the label for more information.

**Uniform Water Distribution and System Calibration:** The irrigation system must provide uniform distribution of treated water. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water. The system must be calibrated to uniformly apply the rates specified. If you have questions about calibration, contact Cooperative Extension Service specialists, equipment manufacturers, or other experts.

**Chemigation Monitoring:** A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

**Drift:** Do not apply when wind speed favors drift beyond the area intended for treatment.

**Required System Safety Devices:** The system must contain a functional check valve, vacuum relief valve, and low-pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected. Systems must use a metering pump such as a positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

**Using Water from Public Water Systems:** Public water systems means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year. Chemigation systems connected to public water systems must contain a functional reduced-pressure zone, back flow preventer (RPZ), or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and to top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe. The pesticide injection pipeline must contain a functional automatic quick-closing check valve to prevent the flow of fluid back toward the injection. The pesticide injection pipeline must contain a functional normally closed solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected. Systems must use a metering pump such as a positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

**ROTATIONAL CROPS\***

Replant treated areas with any crop specified on an imidacloprid label or any crop for which a tolerance exists for the active ingredient as soon as practical following the last application. For crops not listed on an imidacloprid label, or for crops for which no tolerances for the active ingredient have been established, a 12-month plant-back interval must be observed.
<b>IMMEDIATE PLANT-BACK:</b> All crops on this label plus the following crops not on this label: barley, canola, corn (field, pop, and sweet), rapeseed, sorghum, sugarbeet, and wheat
<b>30-DAY PLANT-BACK:</b> Cereals (including buckwheat, millet, oats, rice, rye, and triticale), soybeans, and safflower
<b>10-MONTH PLANT-BACK:</b> Onion and bulb vegetables
<b>12-MONTH PLANT-BACK:</b> All Other Crops

\*Plant cover crops for soil building or erosion control at any time, but do not graze or harvest for food or feed.

19  
29

## FIELD CROPS

### COTTON – soil treatment

Pests Controlled	Rate: Fluid ounces per 1000 row-feet	Rate: Fluid ounces per acre
Cotton aphid, Plant bugs, Thrips, Whiteflies	0.65	8.5 –10.55 (depending on row-spacing)
<b>Restrictions:</b> <ul style="list-style-type: none"> <li>• Maximum MANA Alias 4F allowed per crop season when making soil applications: 10.55 fluid ounces per acre (0.33 lb active ingredient per acre).</li> <li>• Regardless of formulation or method of application, apply no more than 0.5 lb. active ingredient per acre per season, including seed treatment, soil, and foliar uses.</li> <li>• Do not apply more than a total of 6 applications of the active ingredient per season.</li> <li>• Do not graze treated fields after any application of MANA Alias 4F. See <b>RESISTANCE MANAGEMENT</b> section of this label.</li> </ul> <b>Applications: Apply specified dosage in one of the following methods:</b> <ul style="list-style-type: none"> <li>• In-furrow spray during planting directed on or below seed.</li> <li>• In a narrow band directly below the eventual seed row in a bedding operation 7 or fewer days before planting.</li> <li>• Chemigation into root-zone through low-pressure drip or trickle irrigation.</li> </ul>		

### COTTON – foliar treatment

Pests Controlled	Rate: Fluid ounces per acre
Cotton aphid, Cotton fleahopper, Bandedwinged whitefly, Plant bugs (excludes <i>Lygus hesperus</i> ), Green stink bug, Southern green stink bug, Bollworm/budworm (ovicidal effect)	1-2
<b>Pest Suppressed</b>	<b>Rate: Fluid ounces per acre</b>
Lygus bug ( <i>Lygus hesperus</i> ), Whiteflies (other than bandedwinged whitefly)	1.52-2
<b>Restrictions:</b> <ul style="list-style-type: none"> <li>• Pre-harvest interval (PHI): 14 days</li> <li>• Minimum interval between applications: 7 days</li> <li>• Maximum MANA Alias 4F allowed per crop season when making foliar applications: 10 fluid ounces per acre (0.31 lb. AI per acre)</li> <li>• Regardless of formulation or method of application, apply no more than 0.5 lb. active ingredient per acre per season, including seed treatment, soil, and foliar uses.</li> <li>• Do not graze treated fields after any application of MANA Alias 4F.</li> <li>• Apply MANA Alias 4F through properly calibrated ground, aerial, or chemigation application equipment.</li> <li>• Do not apply more than a total of 6 applications of the active ingredient per season.</li> </ul> <b>Applications:</b> Apply specified rate per acre as a broadcast or directed foliar spray to an infested area as pest populations begin to build. Thorough uniform coverage is necessary to achieve optimum control. Use a spray adjuvant to improve coverage. MANA Alias 4F may not knockdown established and heavy insect populations. Two applications may be required to achieve control. Scout fields and retreat if needed. Tank mix MANA Alias 4F with other insecticides for knockdown of pests or for improved control of other pests.	

#### Tank Mix Instructions

Pests Controlled (in addition to pests listed above)	MANA Alias 4F Rate fluid ounces per acre	Bidrin® 8* Rate: Fluid ounces per acre
For early season control of: Thrips	1-1.52	1.6-3.2
For mid to late season control of: Plant bugs, Stink bugs (including Brown stink bug), Grasshoppers, Saltmarsh caterpillar, Cotton leafperforator	1-1.52	4.0-8.0

**Restrictions (in addition to Restrictions listed above):**

\*Refer to the Bidrin 8 product label for specific use directions. Observe all restrictions and precautions that appear on the label.

### PEANUT<sup>1</sup> – soil treatment

Pests Controlled	Rate: Fluid ounces per acre
Aphids Leafhoppers Whiteflies	8 - 12
<b>Pest Suppressed</b>	<b>Rate: Fluid ounces per acre</b>
Thrips	8 – 12

11  
79

**Restrictions:**

- Pre-Harvest Interval (PHI): 14 days
- Maximum MANA Alias 4F allowed per season: 12 fluid ounces/Acre (0.38 ai/Acre)

Use not permitted in California unless otherwise directed by supplemental labeling.

**Applications:** Apply specified dosage in one of the following methods:

- In-furrow spray during planting directed on or below seed;
- Chemigation into root-zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment.

**Important Note:** Increases in Tomato spotted wilt virus (TSWV) incidence have been observed with applications of MANA Alias 4F on multiple varieties of peanut. This may also be the case with other tospoviruses, or other viruses transmitted by various thrips species or perhaps other pests. Prior to applying MANA Alias 4F to peanuts, consult with the State, Cooperative Extension Service, or a MANA representative, for recommendations. Growers are advised to weigh insect control benefits against potential increase in viral disease levels. In areas where TSWV or other tospovirus are endemic, growers are encouraged to use virus resistant varieties and consult the University of Georgia, Tomato spotted wilt virus index, before applying MANA Alias 4F.

**POTATO – soil treatment**

Pests Controlled	Rate: Fluid ounces per 1000 row-feet	Rate: Fluid ounces per acre
Aphids, Colorado potato beetle, Flea beetles, Leafhoppers, Potato psyllid	0.45-0.65	6.5-10.0
Pests/Diseases Suppressed	Rate: Fluid ounces per 1000 row-feet	Rate: Fluid ounces per acre
Symptoms of: Potato leaf roll virus (PLRV), Potato yellows, Net necrosis, Wireworms (with in-furrow spray at-planting)	0.45-0.65	6.5-10.0

**Restrictions:**

- Maximum MANA Alias 4F allowed per crop season when making soil applications: 10.0 fluid ounces per acre (0.31 lb active ingredient per acre)

**Applications:** Apply specified dosage in one of the following methods:

- In-furrow spray during planting directed on seed pieces or seed potatoes.
- Subsurface side-dress on both sides of the row covered with 3 or more inches of soil.
- Narrow band spray at ground cracking directly over the row during hilling covered with 3 or more inches of soil.
- Narrow band directly below the eventual seed row in a bedding operation 7 or fewer days before planting. For effective pest control or suppression, MANA Alias 4F applications must be placed below soil-surface and in contact with seed piece or within root-zone. For potatoes grown on highly permeable soils with shallow water table, make at-plant applications of MANA Alias 4F in a 2 to 4 inch band (width of planter shoe opening) and completely cover.

**POTATO – seed piece treatment**

Pests Controlled	Rate: Fluid ounces per 100 lbs. of seed	Rate: Fluid ounces per acre*
Aphids, Colorado potato beetle, Flea beetles, Leafhoppers, Potato psyllid, Wireworms (seed-piece protection)	0.2-0.4	4.0-8.0
Pests/Diseases Suppressed	Rate: Fluid ounces per 1000 row-feet	Rate: Fluid ounces per acre
Symptoms of: Potato leaf roll virus (PLRV), Potato yellows, Net necrosis	0.4	8.0

**Restrictions:**

- Maximum MANA Alias 4F allowed per crop season when making seed piece treatment applications: 10.0 fluid ounces per acre (0.31 lb. active ingredient per acre)
- Do not use treated seed-pieces for food, feed, or fodder.
- Do not apply any subsequent application of MANA Alias 4F (in-furrow), Gaucho, Leverage, or Provado following an MANA Alias 4F seed-piece treatment.

**Application:** Apply specified dosage as a diluted spray onto seed-pieces using a shielded spray system. Dilute with 3 parts water, or less, to 1 part MANA Alias 4F. Agitate or stir spray solution as needed. Apply fungicidal or inert absorbent dusts after MANA Alias 4F application. Apply only in areas with adequate ventilation or in areas that are equipped to remove spray mist or dust. Plant seed pieces as soon as possible after treating, avoiding prolonged exposure of MANA Alias 4F treated seed-pieces to sunlight and in accordance with the recommendation of your local Extension service.

\* Based on a seeding rate of 2000 lbs. per acre.

**POTATO – foliar treatment**

Pests Controlled	Rate: Fluid ounces per acre
Aphids, Colorado potato beetle, Flea beetles, Leafhoppers, Psyllids	1.52

**Applications:** Apply specified rate per acre as a broadcast or directed foliar spray to an infested area as pest populations begin to build. Thorough uniform coverage is necessary to achieve optimum control. Use a spray adjuvant to improve coverage. MANA Alias 4F may not knockdown established and heavy insect populations. Two applications may be required to achieve control. Scout fields and retreat if needed. Tank mix MANA Alias 4F with other insecticides for knockdown of pests or for improved control of other pests.

**Restrictions:**

- Pre-Harvest Interval (PHI): 7 days
- Minimum Interval between applications: 7 days
- Maximum MANA Alias 4F allowed per crop season when making foliar applications: 6.4 fluid ounces per acre (0.2 lb. AI per acre)

12  
29

**SOYBEANS – foliar treatment**

PESTS CONTROLLED	RATE (FL OZ/ACRE)
Aphids, Bean leaf beetle, Cucumber beetles/Rootworm adults, Japanese beetle (adults), Leafhoppers, Whiteflies	1.5
<b>Notes and Restrictions:</b>	
<ul style="list-style-type: none"> <li>• Pre-harvest Interval (PHI): 7 days</li> <li>• Minimum interval between applications: 7 days</li> <li>• Maximum MANA Alias 4F allowed per crop season: 4.5 fl oz /acre (0.14 lb AI/A)</li> </ul>	

**TOBACCO – soil treatment**

Pests Controlled	Rate: Fluid ounces per 1000 plants (as seedling tray drench)	Rate: Fluid ounces per 1000 plants (in-furrow or transplant-water)
Aphids, Flea beetles	0.5	0.7
Mole crickets, Whiteflies, Wireworms	0.7-1.4	0.9-1.4
Pests/Disease Suppressed	Rate: Fluid ounces per 1000 plants (as seedling tray drench)	Rate: Fluid ounces per 1000 plants (in-furrow or transplant-water)
Cutworms Symptoms of: Tomato spotted wilt virus (TSWV)	0.7-1.4	0.9-1.4
<b>Restrictions:</b>		
<ul style="list-style-type: none"> <li>• Maximum MANA Alias 4F allowed per crop season when making soil applications or foliar sprays to seedlings: 16.0 fluid ounces per acre (0.5 lb. active ingredient per acre)</li> <li>• Pre-Harvest Interval (PHI): 14 days</li> </ul>		
<b>Applications:</b> Apply specified dosage in one of the following methods:		
<ul style="list-style-type: none"> <li>• Uniform, broadcast foliar spray to seedlings in trays (tray drench) not more than 7 days prior to transplanting followed immediately by overhead irrigation to wash MANA Alias 4F from foliage into potting media. Failure to wash MANA Alias 4F from foliage may result in a reduction in pest control. Handle transplants carefully during setting to avoid dislodging treated potting media from roots.</li> <li>• In-furrow spray or transplant-water drench during setting.</li> <li>• Chemigation into root-zone through low-pressure drip, trickle, micro-sprinkler, or equivalent equipment.</li> </ul>		
<b>Important Note:</b> Proper tray drench applications of MANA Alias 4F have been shown to be the most efficacious method of application. However, apply the specified rate of MANA Alias 4F as a combination of the tray drench in the planthouse and/or transplant-water drench in field. Adverse growing conditions may cause a delay in uptake of MANA Alias 4F into the plant and a delay in control.		

**TOBACCO – foliar treatment**

Pests Controlled	Rate: Fluid ounces per acre
Aphids	0.8-1.6
Flea beetles, Japanese beetle	1.6
<b>Restrictions:</b>	
<ul style="list-style-type: none"> <li>• Pre-Harvest Interval (PHI): 14 days</li> <li>• Minimum interval between applications: 7 days</li> <li>• Maximum MANA Alias 4F allowed per crop season when making foliar applications: 8.9 fluid ounces per acre (0.28 lb. AI per acre)</li> </ul>	
<b>Applications:</b> Apply specified rate per acre as a broadcast or directed foliar spray to an infested area as pest populations begin to build. Thorough uniform coverage is necessary to achieve optimum control. Use a spray adjuvant to improve coverage. MANA Alias 4F may not knockdown established and heavy insect populations. Two applications may be required to achieve control. Scout fields and retreat if needed. Tank mix MANA Alias 4F with other insecticides for knockdown of pests or for improved control of other pests.	

**VEGETABLE AND SMALL FRUIT CROPS**

**CUCURBIT VEGETABLES<sup>1</sup> – soil treatment**

**Crops of Crop Group 9 Including:** Chayote (fruit), Chinese waxgourd (Chinese preserving melon), Citron melon, Cuban pumpkin, Cucumber, Gherkin, Gourd (edible, includes hyotan, cucuzza, hechima, Chinese okra), *Momordica* spp. (includes balsam apple, balsam pear, bitter melon, Chinese cucumber), Muskmelon (hybrids and/or cultivars of *Cucumis melo* including true cantaloupe, cantaloupe, casaba, Crenshaw melon, golden pershaw melon, honeydew melon, honey balls, mango melon, Persian melon, pineapple melon, Santa Claus melon, snake melon, and Winter melon), Pumpkin, Squash (includes summer squash types such as: butternut squash, calabaza, crookneck squash, Hubbard squash, scallop squash, straightneck squash, vegetable marrow and zucchini, and winter squash types such as acorn squash and spaghetti squash), Watermelon (includes hybrids and/or varieties of *Citrullus lanatus*)

**Field Applications. See details below for additional planthouse application instructions.**

Pests Controlled	Rate: Fluid ounces per acre
Aphids, Cucumber beetles, Leafhoppers, Thrips (foliage feeding thrips only), Whiteflies	8.0-12.0
Pests/ Diseases Suppressed	Rate: Fluid ounces per acre
Bacterial wilt (as vectored by various cucumber beetles), Leaf silvering resulting from whitefly feeding	8.0-12.0

13  
29

**Restrictions:**

- Pre-Harvest Interval (PHI): 21days
- Maximum MANA Alias 4F allowed per application when making soil applications: 12.0 fluid ounces per acre (0.38 lb/active ingredient per acre)

<sup>1</sup> Not for use on crops grown for seed unless allowed by state-specific supplemental labeling. [Moved from below per EPA request.]

**Applications:** Apply the specified dosage in one of the following methods:

- Chemigation into root-zone through low- pressure drip, trickle, micro-sprinkler, or equivalent equipment.
- In-furrow spray directed on or below seed.
- Narrow (2" or less) surface band spray over seed-line during planting incorporated to a depth of 1 to 1 ½ inches with sufficient irrigation within 24 hours of application.
- Narrow band spray directly below eventual seed row in bedding operation 14 or fewer days before planting.
- Post-seeding drench, transplant-water drench, or hill drench.
- Subsurface side-dress on both sides of each row. MANA Alias 4F must be incorporated into root-zone.

**CUCURBIT VEGETABLES<sup>1</sup> – soil treatment (continued)**

**Planthouse Applications\***

Pest Controlled	Rate: Fluid ounces per 1000 plants
Aphids, Whiteflies	0.05

**Restrictions:**

- Maximum amount MANA Alias 4F applied in the planthouse: 0.05 fluid ounces (0.00156 lb. active ingredient per 1000 plants)
- Maximum number MANA Alias 4F applications in planthouse: 1

<sup>1</sup> Not for use on crops grown for seed unless allowed by state-specific supplemental labeling.

\* Use not permitted in CA unless otherwise directed by supplemental labeling.

**Applications:** Apply specified dosage to seedlings in trays in the planthouse, targeting soil media (tray drench), not more than 7 days prior to transplanting, in one of the following methods:

- Uniform, broadcast high-volume foliar spray, followed immediately by sufficient overhead irrigation to wash MANA Alias 4F from foliage into potting media without loss of gravitational liquid from the bottom of the tray. Failure to wash MANA Alias 4F from foliage may result in reduced pest control.
- Injection into overhead irrigation system, using adequate volume to thoroughly saturate soil media without loss of gravitational solution from the bottom of the tray.

The application made in the planthouse will only provide short-term protection and is not intended as a substitution for a field application. An additional field application must be made within 2 weeks following transplanting to provide continuous protection. Applications of higher rates or increased number of applications in planthouse may result in significant plant injury. Transplants should be handled carefully during setting to avoid dislodging treated potting media from roots.

**Important Note:** Not all varieties of cucurbit vegetables have been tested for tolerance to MANA Alias 4F applied to seedling flats. Treat a small number of plants and confirm tolerance for 7 days prior to treating entire planthouse.

**GREENHOUSE VEGETABLES<sup>1</sup> – soil treatment**

(Mature plants in production greenhouses): Cucumber, Tomato only

Pests Controlled	Rate: Fluid ounces per 1000 plants
Aphids, Whiteflies	0.7

**Restrictions:**

- Pre-Harvest Interval (PHI): 0 day
- Maximum number MANA Alias 4F applications per crop season when making soil applications: 1

<sup>1</sup> Not for use on crops grown for seed unless allowed by state-specific supplemental labeling.

**Applications:** Apply specified dosage in a minimum of 16 gallons of water for tomatoes and 21 gallons of water for cucumbers using soil drenches, micro-irrigation, drip irrigation, or hand-held or motorized calibrated irrigation equipment. Make applications only to plants grown in field-type soils, potting media, or mixtures thereof. Do not apply to plants grown in non-soil medias such as perlite, vermiculite, rock wool or other soil-less media, or plants growing hydroponically. Do not apply to immature plants since phytotoxicity may occur.

Make applications when infestation pressure surpasses threshold and beneficials are not able to maintain pest populations below damage thresholds. Repellency of bumble bee pollinators and negative effects on some beneficials (*Orius* spp.) can occur when MANA Alias 4F is applied.

Many varieties of vegetables have been tested for tolerance to MANA Alias 4F and show good safety. However, certain varieties may show more sensitivity to MANA Alias 4F. Therefore, treat a few plants before treating the whole greenhouse.

**FRUITING VEGETABLES<sup>1</sup> – soil treatment**

Crops of Crop Group 8 plus Okra including: Eggplant, Ground cherry, Okra, Pepper (including bell, chili, cooking, pimento, and sweet), Tomato, Pepinos, Tomatillo

**Field Applications. See details below for additional planthouse applications.**

Pests Controlled	Rate: Fluid ounces per Acre
Aphids, Colorado potato beetle, Flea beetles, Leafhoppers, Thrips (foliage feeding thrips only), Whiteflies	Okra and Pepper; 8.0-16.0 Other Crops; 8.0-12.0
Diseases Suppressed	Rate: Fluid ounces per Acre
Symptoms of: Tomato mottle virus, Tomato spotted wilt virus, Tomato yellow leaf curl virus	Okra and Pepper; 8.0-16.0 Other Crops; 8.0-12.0

**Restrictions:**

- Pre-Harvest Interval (PHI): 21 days
- Maximum MANA Alias 4F allowed on pepper and okra crops per application when making soil applications: 16.0 fluid ounces/Acre (0.5 lb A.I. per acre)
- Maximum MANA Alias 4F allowed on other fruiting vegetable crops per application when making soil applications: 12.0 fluid ounces/Acre (0.38 lb AI/ per acre)

<sup>1</sup> Not for use on crops grown for seed unless allowed by state-specific supplemental labeling.

14  
29

**Applications:** Apply specified dosage in one of the following methods:

- Chemigation into root-zone through low-pressure drip, trickle, micro-sprinkler, or equivalent equipment.
- In-furrow spray directed on or below seed.
- Narrow (2" or less) surface band spray over seed-line during planting incorporated to a depth of 1 to 1 ½ inches with sufficient irrigation within 24 hours of application.
- Narrow band spray directly below eventual seed row in bedding operation 14 or fewer days before planting.
- Post-seeding drench, transplant-water drench, or hill drench.
- Subsurface side-dress on both sides of each row. Incorporate MANA Alias 4F into root-zone.

**FRUITING VEGETABLES<sup>1</sup> – soil treatment (continued)**

**Planthouse Applications\***

Pests Controlled	Rate: Fluid ounces per 1000 plants
Aphids, Whiteflies	0.05

**Restrictions:**

- Maximum amount MANA Alias 4F applied in the planthouse: 0.05 fluid ounces (0.00156 lb A.I.) per 1000 plants.
  - Maximum number MANA Alias 4F applications in planthouse: 1
- <sup>1</sup> Not for use on crops grown for seed unless allowed by state-specific supplemental labeling. [Moved up to restrictions section]  
 \*Use not permitted in CA unless otherwise directed by supplemental labeling.

**Applications:** Apply specified dosage to seedlings in trays in the planthouse, targeting soil media (tray drench), not more than 7 days prior to transplanting, in one of the following manners:

- Uniform, broadcast high-volume foliar spray, followed immediately by sufficient overhead irrigation to wash MANA Alias 4F from foliage into potting media without loss of gravitational liquid from the bottom of the tray. Failure to wash MANA Alias 4F from foliage may result in reduced pest control.
- Injection into overhead irrigation system, using adequate volume to thoroughly saturate soil media without loss of gravitational solution from the bottom of the tray.

The application made in the planthouse will only provide short-term protection and is not intended as a substitution for a field application. An additional field application must be made within 2 weeks following transplanting to provide continuous protection. Applications of higher rates or increased number of applications in planthouse may result in significant plant injury. Handle transplants carefully during setting to avoid dislodging treated potting media from roots.

**Important Note:** Not all varieties of fruiting vegetables have been tested for tolerance to MANA Alias 4F applied to seedling flats. Treat a small number of plants and confirm tolerance for 7 days prior to treating entire planthouse.

**FRUITING VEGETABLES<sup>1</sup> – foliar treatment**

Crops of Group 8 plus Okra, Including: Eggplant, Ground cherry, Okra, Pepper (including bell, chili, cooking, pimento, and sweet), Tomato, Pepinos, Tomatillo

Pests Controlled	Rate: Fluid ounces per acre
Aphids, Colorado potato beetle, Leafhoppers, Whiteflies	1.5- 2.4
Pepper weevil	2.4

**Restrictions:**

- Pre-harvest interval (PHI): 0 day
- Minimum interval between applications: 5 days
- Maximum MANA Alias 4F allowed per crop season when making foliar applications: 7.6 fluid ounces per acre (0.24 lb AI per acre)

<sup>1</sup> Not for use on crops grown for seed unless allowed by state-specific supplemental labeling.

**Applications:**

Apply specified rate per acre as a broadcast or directed foliar spray to infested area as pest populations begin to build. Thorough uniform coverage is necessary to achieve optimum control. Use a spray adjuvant to improve coverage. MANA Alias 4F may not knockdown established and heavy insect populations. Two applications may be required to achieve control. Scout fields and retreat if needed. Tank mix MANA Alias 4F with other insecticides for knockdown of pests or for improved control of other pests.

Crops contained within certain Crop Groups recognized by EPA are subject to change. Refer to the EPA website ([www.epa.gov](http://www.epa.gov)) for latest Crop Groups.

For pepper weevil, apply specified dosage of MANA Alias 4F by ground equipment only, timing applications prior to a damaging population becoming established. Good coverage of foliage and fruit is necessary for optimum control. Applications of MANA Alias 4F must be incorporated into a full-season program where alternations of effective products from multiple classes of chemistry and different modes of action are utilized in a blocked or windowed approach.

For additional information, please contact your Makhteshim Agan of North America, Inc. representative, Extension Specialist or crop advisor.

When targeting adult whiteflies, use higher rates.

**GLOBE ARTICHOKE – foliar treatment**

Pests Controlled	Rate: Fluid ounces per acre
Aphids, Leafhoppers	1.6-4.0

**Restrictions:**

- Pre-harvest interval (PHI): 7 days
- Minimum interval between applications: 14 days
- Maximum MANA Alias 4F allowed per crop season when making foliar applications: 16 fluid ounces per acre (0.5 lb AI per acre)

**Applications:** Apply specified rate per acre as a broadcast or directed foliar spray to infested area as pest populations begin to build. Thorough uniform coverage is necessary to achieve optimum control. Use a spray adjuvant to improve coverage. MANA Alias 4F may not knockdown established and heavy insect populations. Two applications may be required to achieve control. Scout fields and retreat if needed. Tank mix MANA Alias 4F with other insecticides for knockdown of pests or for improved control of other pests.

Crops contained within certain Crop Groups recognized by EPA are subject to change. Refer to the EPA website ([www.epa.gov](http://www.epa.gov)) for latest Crop Groups.

15  
29

**HERBS– soil treatment**

**Crops of Crop Subgroup 19A including:** Angelica, Balm (lemon balm), Basil (fresh and dried), Borage, Bumet, Camomile, Catnip, Chervil (dried), Chinese chive, Chive, Clary, Coriander (cilantro or Chinese parsley leaves), Costmary, Culantro (leaf), Curry (leaf), Dillweed, Horehound, Hyssop, Lavender, Lemongrass, Lovage (leaf), Marigold, Marjoram, Nasturtium, Parsley (dried), Pennyroyal, Rosemary, Rue, Sage, Savory (summer and winter), Sweet bay (bay leaf), Tansy, Tarragon, Thyme, Wintergreen, Woodruff, Wormwood.

<b>Pests Controlled</b>	<b>Rate: Fluid ounces per acre</b>
Aphids, Flea beetles, Leafhoppers, Whiteflies	8.0-12.0
<b>Pests Suppressed</b>	<b>Rate: Fluid ounces per acre</b>
Thrips (foliage feeding thrips only)	8.0-12.0
<b>Restrictions:</b>	
<ul style="list-style-type: none"> <li>• Pre-Harvest Interval (PHI): 14 days</li> <li>• Maximum MANA Alias 4F per season when making soil applications: 12.0 fluid ounces/Acre (0.38 lb AI/Acre)</li> </ul>	
<b>Applications:</b> Apply specified dosage in one of the following methods:	
<ul style="list-style-type: none"> <li>• In-furrow spray during planting directed on or below seed.</li> <li>• In-furrow spray or transplant-water drench during setting or transplanting.</li> <li>• Shankd-into or below eventual seed-line.</li> <li>• Chemigation into root-zone through low-pressure drip, trickle, micro-sprinkler, or equivalent equipment.</li> </ul>	
<b>Notes:</b> Not all crops and/or varieties listed above have been tested for phytotoxic effects. Without specific knowledge about a particular crop and variety, treat only a small area or small number of plants of each listed above and evaluate prior to commercial use.	

**HERBS – foliar treatment**

**Crops of Crop Subgroup 19A including:** Angelica, Balm (lemon balm), Basil (fresh and dried), Borage, Bumet, Camomile, Catnip, Chervil (dried), Chinese chive, Chive, Clary, Coriander (cilantro or Chinese parsley leaves), Costmary, Culantro (leaf), Curry (leaf), Dillweed, Horehound, Hyssop, Lavender, Lemongrass, Lovage (leaf), Marigold, Marjoram, Nasturtium, Parsley (dried), Pennyroyal, Rosemary, Rue, Sage, Savory (summer and winter), Sweet bay (bay leaf), Tansy, Tarragon, Thyme, Wintergreen, Woodruff, Wormwood.

<b>Pests Controlled</b>	<b>Rate: Fluid ounces per Acre</b>
Aphids, Flea beetles, Leafhoppers, Whiteflies	1.4
<b>Restrictions:</b>	
<ul style="list-style-type: none"> <li>• Pre-Harvest Interval (PHI): 7 days</li> <li>• Minimum interval between applications: 5 days</li> <li>• Maximum MANA Alias 4F allowed per season when making foliar applications: 4.2 fluid ounces per acre (0.13 lb AI per acre)</li> </ul>	
<b>Applications:</b> Apply specified rate per acre as a broadcast or directed foliar spray to infested area as pest populations begin to build. Thorough uniform coverage is necessary to achieve optimum control. Use a spray adjuvant to improve coverage. MANA Alias 4F may not knockdown established and heavy insect populations. Two applications may be required to achieve control. Scout fields and retreat if needed. Tank mix MANA Alias 4F with other insecticides for knockdown of pests or for improved control of other pests.	
Crops contained within certain Crop Groups recognized by EPA are subject to change. Refer to the EPA website ( <a href="http://www.epa.gov">www.epa.gov</a> ) for latest Crop Groups.	
Apply MANA Alias 4F through properly calibrated ground and aerial application equipment. Thorough coverage with direct contact of the spray adjuvant to the target pests is required for optimum control. The addition of an organosilicone-based spray adjuvant at a rate not to exceed the adjuvant manufacturer's specified use rate may improve coverage and control.	
<b>Note:</b> Not all crops and/or varieties listed above have been tested for phytotoxic effects. Without specific knowledge about a particular crop and variety, treat only a small area or small number of plants of each listed above and evaluate prior to commercial use.	

**BRASSICA (COLE) LEAFY VEGETABLES<sup>1</sup> – soil treatment**

**Crops of Crop Group 5 including:** Broccoli, Broccoli raab (rapini), Brussels sprouts, Cabbage, Cauliflower, Cavalo broccoli, Chinese (gai lon) broccoli, Chinese (bok choy) cabbage, Chinese (napa) cabbage, Chinese mustard (gai choy) cabbage, Collards, Kale, Kohlrabi, Mizuna, Mustard greens, Mustard spinach, Rape greens, Turnip tops (leaves)

<b>Pests Controlled</b>	<b>Rate: Fluid ounces per acre (on 36 inch rows)</b>
Aphids, Leafhoppers, Thrips (foliage feeding thrips only), Whiteflies	5.0-12.0
<b>Restrictions:</b>	
<ul style="list-style-type: none"> <li>• Pre-Harvest Interval (PHI): 21 days</li> <li>• Maximum MANA Alias 4F allowed per application when making soil applications: 12.0 fluid ounces/Acre (0.38 lb AI per acre)</li> </ul>	
<sup>1</sup> Not for use on crops grown for seed unless allowed by state-specific supplemental labeling.	
<b>Applications:</b> Apply specified dosage in one of the following methods:	
<ul style="list-style-type: none"> <li>• Chemigation into root-zone through low-pressure drip, trickle, micro-sprinkler, or equivalent equipment.</li> <li>• In-furrow spray directed on or below seed.</li> <li>• Narrow (2 inches or less) surface band spray over seed-line during planting incorporated to a depth of 1 to 1 ½ inches with sufficient irrigation within 24 hours of application.</li> <li>• Narrow band spray directly below eventual seed row in bedding operation 14 or fewer days before planting.</li> <li>• Post-seeding drench, transplant-water drench, or hill drench.</li> <li>• Subsurface side-dress on both sides of each row. MANA Alias 4F must be incorporated into root-zone.</li> </ul>	

16  
29

**BRASSICA (COLE) LEAFY VEGETABLES<sup>1</sup> – foliar treatment**

**Crops of Crop Group 5 including:** Broccoli, Broccoli raab (rapini), Brussels sprouts, Cabbage, Cauliflower, Cavalo broccoli, Chinese (gai lon) broccoli, Chinese (bok choy) cabbage, Chinese (napa) cabbage, Chinese mustard (gai choy) cabbage, Collards, Kale, Kohlrabi, Mizuna, Mustard greens, Mustard spinach, Rape greens, Turnip tops (leaves)

Pests Controlled	Rate: Fluid ounces per acre
Aphids, Fleabeetles, Leafhoppers, Whiteflies	1.5
<b>Restrictions:</b> <ul style="list-style-type: none"> <li>• Pre-Harvest Interval (PHI): 7 days</li> <li>• Minimum interval between applications: 5 days</li> <li>• Maximum MANA Alias 4F allowed per crop season when making foliar applications: 7.68 fluid ounces/Acre (0.24 lb AI per acre)</li> </ul> <sup>1</sup> Not for use on crops grown for seed unless allowed by state-specific supplemental labeling. <p><b>Applications:</b> Apply specified rate per acre as a broadcast or directed foliar spray to infested area as pest populations begin to build. Thorough uniform coverage is necessary to achieve optimum control. Use a spray adjuvant to improve coverage. MANA Alias 4F may not knockdown established and heavy insect populations. Two applications may be required to achieve control. Scout fields and retreat if needed. Tank mix MANA Alias 4F with other insecticides for knockdown of pests or for improved control of other pests. Crops contained within certain Crop Groups recognized by EPA are subject to change. Refer to the EPA website (<a href="http://www.epa.gov">www.epa.gov</a>) for latest Crop Groups.</p>	

**LEAFY VEGETABLES<sup>1</sup> – soil treatment**

**Crops of Crop Subgroup 4A plus Watercress including:** Amaranth (leafy amaranth, Chinese spinach, tampala), Arugula (Roquette), Chervil, Chrysanthemum (edible leaved and garland), Cilantro, Corn salad, Cress (garden), Cress (upland, yellow rocket, winter cress), Dandelion, Dock (sorrel), Endive (escarole), Lettuce (head and leaf), Orach, Parsley, Purslane (garden and winter), Radicchio (red chicory), Spinach (including New Zealand and vine (Malabar spinach, Indian spinach)), Watercress (commercial production only, applications must not be made to native cress growing in streams or other bodies of water), Watercress (upland)

Pests Controlled	Rate: Fluid ounces per acre (on 36 inch rows)
Aphids, Leafhoppers, Thrips (foliage feeding thrips only), Whiteflies	5.0-12.0
<b>Restrictions:</b> <ul style="list-style-type: none"> <li>• Pre-Harvest Interval (PHI): 21 days</li> <li>• Maximum MANA Alias 4F allowed per application when making soil applications: 12.0 fluid ounces per acre (0.38 lb AI per Acre)</li> </ul> <sup>1</sup> Not for use on crops grown for seed unless allowed by state-specific supplemental labeling. <p><b>Applications:</b> Apply specified dosage in one of the following methods:</p> <ul style="list-style-type: none"> <li>• Chemigation into root-zone through low-pressure drip, trickle, micro-sprinkler, or equivalent equipment.</li> <li>• In-furrow spray directed on or below seed.</li> <li>• Narrow (2 inches or less) surface band spray over seed-line during planting incorporated to a depth of 1 to 1 ½ inches with sufficient irrigation within 24 hours of application.</li> <li>• Narrow band spray directly below eventual seed row in bedding operation 14 or fewer days before planting.</li> <li>• Post-seeding drench, transplant-water drench, or hill drench.</li> <li>• Subsurface side-dress on both sides of each row. MANA Alias 4F must be incorporated into root-zone.</li> </ul>	

**LEAFY VEGETABLES<sup>1</sup> – foliar treatment**

**Crops of Crop Subgroup 4A plus Watercress including:** Amaranth (leafy amaranth, Chinese spinach, tampala), Arugula (Roquette), Chervil, Chrysanthemum (edible leaved and garland), Cilantro, Corn salad, Cress (garden), Cress (upland, yellow rocket, winter cress), Dandelion, Dock (sorrel), Endive (escarole), Lettuce (head and leaf), Orach, Parsley, Purslane (garden and winter), Radicchio (red chicory), Spinach (including New Zealand and vine (Malabar spinach, Indian spinach)), Watercress (commercial production only, applications must not be made to native cress growing in streams or other bodies of water), Watercress (upland)

Pests Controlled	Rate: Fluid ounces per acre
Aphids, Flea beetles, Leafhoppers, Whiteflies	1.5
<b>Restrictions:</b> <ul style="list-style-type: none"> <li>• Pre-Harvest Interval (PHI): 7 days</li> <li>• Minimum interval between applications: 5 days</li> </ul> Maximum MANA Alias 4F allowed per crop season when making foliar applications: 7.6 fluid ounces per acre (0.24 lb AI per acre) <sup>1</sup> Not for use on crops grown for seed unless allowed by state-specific supplemental labeling. <p><b>Applications:</b> Apply specified rate per acre as a broadcast or directed foliar spray to infested area as pest populations begin to build. Thorough uniform coverage is necessary to achieve optimum control. Use a spray adjuvant to improve coverage. MANA Alias 4F may not knockdown established and heavy insect populations. Two applications may be required to achieve control. Scout fields and retreat if needed. Tank mix MANA Alias 4F with other insecticides for knockdown of pests or for improved control of other pests. Crops contained within certain Crop Groups recognized by EPA are subject to change. Refer to the EPA website (<a href="http://www.epa.gov">www.epa.gov</a>) for latest Crop Groups.</p> <p>For applications made to watercress, production fields must be drained of water at least 24 hours prior to application and water must not be reapplied to the field for a minimum of 24 hours following the applications. Applications must be made to fully leafed-up canopies only.</p>	



17  
29

**LEAFY PETIOLE VEGETABLES<sup>1</sup> – soil treatment**

**Crops of Crop Subgroup 4B including:** Cardoon, Celery, Celtuce, Chinese celery (fresh leaves and stalk only), Florence fennel (including sweet anise, sweet fennel, Finocchio), Rhubarb, Swiss chard

Pests Controlled	Rate: Fluid ounces per acre
Aphids, Leafhoppers, Thrips (foliage feeding thrips only), Whiteflies	5.0-12.0
<b>Restrictions:</b>	
<ul style="list-style-type: none"> <li>• Pre-Harvest Interval (PHI): 45 days</li> <li>• Maximum MANA Alias 4F allowed per application when making soil applications: 12.0 fluid ounces/Acre (0.38 lb AI per acre)</li> </ul>	
<sup>1</sup> Not for use on crops grown for seed unless allowed by state-specific supplemental labeling.	
<b>Applications:</b> Apply specified dosage in one of the following methods:	
<ul style="list-style-type: none"> <li>• Chemigation into root-zone through low-pressure drip, trickle, micro-sprinkler, or equivalent equipment.</li> <li>• In-furrow spray directed on or below seed.</li> <li>• Narrow (2 inches or less) surface band spray over seed-line during planting incorporated to a depth of 1 to 1 ½ inches with sufficient irrigation within 24 hours of application.</li> <li>• Narrow band spray directly below eventual seed row in bedding operation 14 or fewer days before planting.</li> <li>• Post-seeding drench, transplant-water drench, or hill drench.</li> <li>• Subsurface side-dress on both sides of each row. MANA Alias 4F must be incorporated into root-zone.</li> </ul>	

**LEGUME VEGETABLES<sup>1</sup> except soybean, dry-soil treatment**

**Crops of Crop Group 6 including: Edible Podded and Succulent Shelled Pea and Bean and Dried Shelled Pea and Bean**

**Bean** (*Lupinus* spp., includes grain lupin, sweet lupin, white lupin, and white sweet lupin)

**Bean** (*Phaseolus* spp., includes field bean, kidney bean, lima bean, navy bean, pinto bean, runner bean, snap bean, tepary bean, wax bean)

**Bean** (*Vigna* spp., includes adzuki bean, asparagus bean, blackeyed pea, catjang, Chinese longbean, cowpea, Crowder pea, moth bean, mung bean, rice bean, Southern pea, urd bean, yardlong bean)

**Pea** (*Pisum* spp., includes dwarf pea, edible-pod pea, English pea, field pea, garden pea, green pea, snow pea, sugar snap pea)

**Other Beans and Peas** [Broad bean (fava), Chickpea (garbanzo bean), Guar, Jackbean, Lablab bean (hyacinth bean), Lentil, Pigeon pea, Soybean (immature seed), Sword bean]

Pests Controlled	Rate: Fluid ounces per acre
Aphids, Leafhoppers, Thrips (foliage feeding thrips only), Whiteflies	8.0-12.0
<b>Diseases Suppressed</b>	
Symptoms of: Bean common mosaic virus (BCMV), Bean golden mosaic virus (BGMV), Beet curly top hybrigeminivirus (BCTV)	8.0-12.0
<b>Restrictions:</b>	
<ul style="list-style-type: none"> <li>• Pre-Harvest Interval (PHI): 21 days</li> <li>• Maximum MANA Alias 4F allowed per crop season when making soil applications: 12.0 fluid ounces/Acre (0.38 lb AI per acre)</li> </ul>	
<sup>1</sup> Not for use on crops grown for seed unless allowed by state-specific supplemental labeling.	
<b>Applications:</b> Apply specified dosage in one of the following methods:	
<ul style="list-style-type: none"> <li>• Chemigation into root-zone through low-pressure drip, trickle, micro-sprinkler, or equivalent equipment.</li> <li>• In-furrow spray at planting directed on or below seed.</li> <li>• In a narrow (2 inches or less) surface band over seed-line during planting incorporated to a depth of 1 to 1 ½ inches with sufficient irrigation with 24 hours following application.</li> <li>• In a narrow band directly below the eventual seed row in a bedding operation 7 or fewer days before planting.</li> <li>• As a post-seeding drench, transplant drench, or hill drench.</li> </ul>	

**LEGUME VEGETABLES<sup>1</sup> except soybean, dry-foliar treatment**

**Crops of Crop Group 6 including: Edible Podded and Succulent Shelled Pea and Bean and Dried Shelled Pea and Bean**

**Bean** (*Lupinus* spp., includes grain lupin, sweet lupin, white lupin, and white sweet lupin)

**Bean** (*Phaseolus* spp., includes field bean, kidney bean, lima bean, navy bean, pinto bean, runner bean, snap bean, tepary bean, wax bean)

**Bean** (*Vigna* spp., includes adzuki bean, asparagus bean, blackeyed pea, catjang, Chinese longbean, cowpea, Crowder pea, moth bean, mung bean, rice bean, Southern pea, urd bean, yardlong bean)

**Pea** (*Pisum* spp., includes dwarf pea, edible-pod pea, English pea, field pea, garden pea, green pea, snow pea, sugar snap pea)

**Other Beans and Peas** [Broad bean (fava), Chickpea (garbanzo bean), Guar, Jackbean, Lablab bean (hyacinth bean), Lentil, Pigeon pea, Soybean (immature seed), Sword bean]

Pests Controlled	Rate: Fluid ounces per acre
Aphids, Leafhoppers, Whiteflies	1.4
<b>Restrictions:</b>	
<ul style="list-style-type: none"> <li>• Pre-Harvest Interval (PHI): 7 days</li> <li>• Minimum interval between applications: 7 days</li> <li>• Maximum MANA Alias 4F allowed per crop season when making foliar applications: 4.2 fluid ounces per acre (0.13 lb AI per acre)</li> </ul>	
<sup>1</sup> Not for use on crops grown for seed unless allowed by state-specific supplemental labeling.	
<b>Applications:</b> Apply specified rate per acre as a broadcast or directed foliar spray to infested area as pest populations begin to build. Thorough uniform coverage is necessary to achieve optimum control. Use a spray adjuvant to improve coverage. MANA Alias 4F may not knockdown established and heavy insect populations. Two applications may be required to achieve control. Scout fields and retreat if needed. Tank mix MANA Alias 4F with other insecticides for knockdown of pests or for improved control of other pests. Crops contained within certain Crop Groups recognized by EPA are subject to change. Refer to the EPA website ( <a href="http://www.epa.gov">www.epa.gov</a> ) for latest Crop Groups.	

18  
29

**ROOT VEGETABLES<sup>1</sup> – soil treatment**

**Crops of Crop Subgroup 1B except Sugarbeet including:** Beet (garden)<sup>2</sup>, Burdock (edible)<sup>2</sup>, Carrot<sup>2</sup>, Celeriac<sup>2</sup>, Chervil (turnip-rooted)<sup>2</sup>, Chicory<sup>2</sup>, Ginseng, Horseradish, Kava<sup>2,3</sup>, Parsley (turnip-rooted), Parsnip<sup>2</sup>, Radish<sup>2</sup>, Oriental radish (diakon)<sup>2</sup>, Rutabaga<sup>2</sup>, Salsify (oyster plant), Salsify (black)<sup>2</sup>, Salsify (Spanish), Skirret, and Turnip<sup>2</sup>

Pests Controlled	Rate: Fluid ounces per 1000 row-feet	Rate: Fluid ounces per acre
Aphids, Flea beetles, Leafhoppers, Thrips (foliage feeding thrips only), Whiteflies	0.35-0.85	5.0-12.0

**Restrictions:**

- Pre-Harvest Interval (PHI): 21 days
- Maximum MANA Alias 4F allowed per crop season when making soil applications: 12.0 fluid ounces/Acre (0.38 lb AI per acre)
- Maximum MANA Alias 4F soil applications per crop season: 1

<sup>1</sup>Not for use on crops grown for seed unless allowed by a state-specific supplemental labeling.

<sup>2</sup>Tops or greens from these crops may be utilized for food or feed.

<sup>3</sup>Use not permitted in California unless otherwise directed by supplemental labeling.

**Application:** Apply specified dosage in one of the following methods:

- Chemigation into root-zone through low-pressure drip, trickle, micro-sprinkler, or equivalent equipment.
- In-furrow spray (rate specified per 1000 row-feet) or, shanked-in 1 to 2 inches below seed depth during planting.
- In a narrow (2 inches or less) band directly (1 to 2 inches) below the eventual seed row in a bedding operation 14 or fewer days before planting.

**Important Note:** The rate applied affects the length of control. Use higher rates where infestations occur later in crop development, or where pest pressure is continuous. MANA Alias 4F rates less than 0.7 fluid ounces/1000 row-feet will not provide adequate residual pest control. MANA Alias 4F treated crops grown on very high organic matter soils (muck) may also require additional pest management control.

**ROOT VEGETABLES<sup>1</sup> – foliar treatment**

**Crops of Crop Subgroup 1B except Sugarbeet including:** Beet (garden)<sup>2</sup>, Burdock (edible)<sup>2</sup>, Carrot<sup>2</sup>, Celeriac<sup>2</sup>, Chervil (turnip-rooted)<sup>2</sup>, Chicory<sup>2</sup>, Ginseng, Horseradish, Kava<sup>2,3</sup>, Parsley (turnip-rooted), Parsnip<sup>2</sup>, Radish<sup>2</sup>, Oriental radish (diakon)<sup>2</sup>, Rutabaga<sup>2</sup>, Salsify (oyster plant), Salsify (black)<sup>2</sup>, Salsify (Spanish), Skirret, Turnip<sup>2</sup>.

Pests Controlled	Rate: Fluid ounces per acre
Aphids, Flea beetles, Leafhoppers, Whiteflies	1.4

**Restrictions:**

- Pre-Harvest Interval (PHI): 7 days
- Minimum interval between applications: 5 days
- Maximum MANA Alias 4F allowed per crop season when making foliar applications: 1.4 fluid ounces per acre (0.044 lb AI per acre) on Radish, 4.2 fluid ounces per acre (0.13 lb AI per acre) on other crops.
- Maximum MANA Alias 4F application(s) per crop season: 1 on radish, 3 on all other crops

<sup>1</sup>Not for use on crops grown for seed unless allowed by state-specific supplemental labeling.

<sup>2</sup>Tops or greens from these crops may be utilized for food or feed.

<sup>3</sup>Use not permitted in California unless otherwise directed by supplemental labeling.

**Applications:** Apply specified rate per acre as a broadcast or directed foliar spray to infested area as pest populations begin to build. Thorough uniform coverage is necessary to achieve optimum control. Use a spray adjuvant to improve coverage. MANA Alias 4F may not knockdown established and heavy insect populations. Two applications may be required to achieve control. Scout fields and retreat if needed. Tank mix MANA Alias 4F with other insecticides for knockdown of pests or for improved control of other pests.

Crops contained within certain Crop Groups recognized by EPA are subject to change. Refer to the EPA website ([www.epa.gov](http://www.epa.gov)) for latest Crop Groups.

**TUBEROUS and CORM VEGETABLES<sup>1</sup> – soil treatment**

**Crops of Crop Subgroup 1C including:** Arracacha, Arrowroot, Artichoke (Chinese and Jerusalem), Canna (edible, Queensland arrowroot), Cassava (bitter and sweet)<sup>2</sup>, Chayote (root), Chufa, Dasheen (taro)<sup>2</sup>, Ginger, Leren, Sweet potato, Tanier (cocoyam)<sup>2</sup>, Turmeric, Yam bean (jicama, manioc pea), Yam (true)<sup>2</sup> (For applications on potato see **FIELD CROPS** section)

Pests Controlled	Rate: Fluid ounces per 1000 row-feet	Rate: Fluid ounces per acre
Aphids, Flea beetles, Leafhoppers Thrips (foliage feeding thrips only), Whiteflies	0.35-0.85	5.0-12.0

**Restrictions:**

- Pre-Harvest Interval (PHI) from planting application: 3 days (leaves); 125 days (corms)
- Maximum MANA Alias 4F allowed per crop season when making soil applications: 12.0 fluid ounces/Acre (0.38 lb AI per acre)
- Maximum MANA Alias 4F soil applications per crop season: 1

<sup>1</sup>Not for use on crops grown for seed unless allowed by state-specific supplemental labeling.

<sup>2</sup>Tops or greens from these crops may be utilized for food or feed.

**Applications:** Apply specified dosage in one of the following methods:

- In-furrow spray (rate specified per 1000 row-feet) over planting materials (hulis) or shanked-in 1 to 2 inches below hulis depth at planting.
- Side-dress not more than 0.3 fluid ounces/1000 row-feet no later than 45 days after planting. Observe the same PHI as above.

**Important Note:** The rate applied affects the length of control. Use higher rates where infestations occur late in crop development, or where pest pressure is continuous. MANA Alias 4F rates less than 0.35 fluid ounces/1000 row-feet may not provide adequate residual pest control. MANA Alias 4F treated crops grown on very high organic matter soils (muck) may also require additional pest management control.

19  
29

**TUBEROUS and CORM VEGETABLES<sup>1</sup> – foliar treatment**

**Crops of Crop Subgroup 1C including:** Arracacha, Arrowroot, Artichoke (Chinese and Jerusalem), Canna (edible, Queensland arrowroot), Cassava (bitter and sweet)<sup>2</sup>, Chayote (root), Chufa, Dasheen (taro)<sup>2</sup>, Ginger, Leren, Sweetpotato<sup>2</sup>, Tanier (cocoyam)<sup>2</sup>, Turmeric, Yam bean (jicama, manioc pea), Yam (true)<sup>2</sup> (For applications on potato see **FIELD CROPS** section)

Pests Controlled	Rate: Fluid ounces per acre
Aphids, Flea beetles, Leafhoppers, Whiteflies	1.4
<b>Restrictions:</b>	
<ul style="list-style-type: none"> <li>• Pre-Harvest Interval (PHI): 7 days</li> <li>• Minimum interval between applications: 5 days</li> <li>• Maximum MANA Alias 4F allowed per crop season when making foliar applications: 1.4 fluid ounces per acre (0.044 lb AI per acre) on Radish, 4.2 fluid ounces per acre (0.13 lb AI per acre) on other crops.</li> <li>• Maximum MANA Alias 4F application(s) per crop season: 3 on all crops</li> </ul>	
<sup>1</sup> Not for use on crops grown for seed unless allowed by state-specific supplemental labeling.	
<sup>2</sup> Tops or greens from these crops may be utilized for food or feed.	
<p><b>Applications:</b> Apply specified rate per acre as a broadcast or directed foliar spray to infested area as pest populations begin to build. Thorough uniform coverage is necessary to achieve optimum control. Use a spray adjuvant to improve coverage. MANA Alias 4F may not knockdown established and heavy insect populations. Two applications may be required to achieve control. Scout fields and retreat if needed. Tank mix MANA Alias 4F with other insecticides for knockdown of pests or for improved control of other pests.</p> <p>Crops contained within certain Crop Groups recognized by EPA are subject to change. Refer to the EPA website (<a href="http://www.epa.gov">www.epa.gov</a>) for latest Crop Groups.</p>	

**STRAWBERRY<sup>1</sup> – soil treatment**

Annual and Perennial Crops	
Pests Controlled	Rate: Fluid ounces per acre
Aphids, Whiteflies	12.0-16.0
<b>Restrictions:</b>	
<ul style="list-style-type: none"> <li>• Pre-Harvest Interval (PHI): 14 days</li> <li>• Maximum MANA Alias 4F allowed per crop season when making soil applications: 16.0 fluid ounces/Acre (0.50 lb AI per acre)</li> </ul>	
<sup>1</sup> Do not use both application methods on the same crop in the same season.	
<b>Applications:</b> Apply specified dosage in one of the following methods:	
<ul style="list-style-type: none"> <li>• Chemigation into root-zone through low-pressure drip, trickle, micro-sprinkler, or equivalent equipment after plants are established or on perennial crops in early spring prior to bud opening.</li> <li>• As a plant material or plant hole treatment just prior to, or during transplanting.</li> <li>• As a band spray over-the-row in a minimum of 20 gallons of water per acre, followed immediately by overhead irrigation to incorporate product into root-zone. Do not use plastic or other mulches that limit movement of MANA Alias 4F into root zone.</li> </ul>	
The rate applied affects the length of control. Use higher rates where infestations occur later in crop development or where pest pressure is continuous.	
Post-harvest Use on Perennial Crops	
Pests Controlled	Rate fluid ounces per acre
White grub complex (grubs of Asiatic garden beetle, European and Masked chafer, Japanese beetle, Oriental beetle)	8.0-12.0
<b>Restrictions:</b>	
<ul style="list-style-type: none"> <li>• Pre-Harvest Interval (PHI): 14 days</li> <li>• Maximum MANA Alias 4F allowed per season when making soil applications: 12.0 fluid ounces per acre (0.38 lb A.I. per acre)</li> </ul>	
<sup>1</sup> Do not use both application methods on the same crop in the same season.	
<b>Applications:</b> Apply a single application post harvest to coincide with renovation of strawberry fields and during active egg-laying period of beetles. Apply specified dosage of MANA Alias 4F in one of the following methods:	
<ul style="list-style-type: none"> <li>• As a ground spray via boom or backpack sprayer in a minimum of 20 gallons of water per acre.</li> <li>• As a row-band spray using an adjusted amount of product based on the treated row band area in proportion to the amount required per full acre. The bandwidth should be equivalent to the width of the anticipated fruiting bed.</li> <li>• As a chemigation application with 600 to 1000 gallons of water followed by 0.1 to 0.25 inches irrigation.</li> </ul>	
<b>Important Note:</b> Follow all soil-surface applications with 0.25 inches of rainfall or overhead irrigation water per acre within 2 hours of application. Failure to adequately incorporate MANA Alias 4F into egg-deposition zone may result in decreased activity.	

**STRAWBERRY – foliar treatment**

Pests Controlled	Rate: Fluid ounces per acre
Aphids, Spittlebugs, Whiteflies	1.5
<b>Restrictions:</b>	
<ul style="list-style-type: none"> <li>• Pre-harvest Interval (PHI): 7 days</li> <li>• Minimum interval between applications: 5 days</li> <li>• Maximum MANA Alias 4F allowed per crop season when making foliar applications: 4.5 fluid ounces per acre (0.14 lb. AI per acre).</li> <li>• Do not apply during bloom or within 10 days prior to bloom or when bees are actively foraging.</li> </ul>	
<p><b>Applications:</b> Apply specified rate per acre as a broadcast or directed foliar spray to infested area as pest populations begin to build. Thorough uniform coverage is necessary to achieve optimum control. Use a spray adjuvant to improve coverage. MANA Alias 4F may not knockdown established and heavy insect populations. Two applications may be required to achieve control. Scout fields and retreat if needed. Tank mix MANA Alias 4F with other insecticides for knockdown of pests or for improved control of other pests.</p> <p>Crops contained within certain Crop Groups recognized by EPA are subject to change. Refer to the EPA website (<a href="http://www.epa.gov">www.epa.gov</a>) for latest Crop Groups.</p>	

20  
29

**SUGARBEET<sup>1</sup> – soil treatment-For use only in CA**

<b>Pests Controlled</b>	<b>Rate: Fluid ounces per acre</b>
Aphids, Leafhoppers, Whiteflies, Flea beetles	3.0-6.0
<b>Diseases Suppressed</b>	<b>Rate: Fluid ounces per acre</b>
Symptoms of: Western yellows/Beet curly top hybrigeminivirus (BCTV)	3.0-6.0
<b>Restrictions:</b>	
<ul style="list-style-type: none"> <li>Maximum MANA Alias 4F allowed per crop season when making soil applications: 6.0 fluid ounces/Acre (0.18 lb AI per acre)</li> <li>Do not apply immediately prior to bud opening or during bloom or when bees are actively foraging.</li> </ul>	
<sup>1</sup> Not for use on crops grown for seed unless allowed by state-specific supplemental labeling.	
<b>Applications:</b> Apply specified dosage in the following method:	
<ul style="list-style-type: none"> <li>Apply specified dosage in sufficient carrier volume to insure uniform application. Apply directly below each seed furrow either during the bedding operation immediately prior to planting or at the time of planting.</li> </ul>	
Apply the low rate to aid establishment of stands in whitefly areas, or for early season control of the other pests listed.	

Rate: Fluid Ounces/Acre	Rate: Fluid ounces/1000 row-feet Based on <u>average</u> row spacing (in inches):							
	10	15	20	25	30	35	40	45
5	0.0475	0.07125	0.095	0.11875	0.1425	0.16625	0.19	0.21375
6	0.057	0.0855	0.114	0.1425	0.171	0.1995	0.228	0.2565
7	0.0665	0.09975	0.133	0.16625	0.1995	0.23275	0.266	0.29925
8	0.076	0.114	0.152	0.19	0.228	0.266	0.304	0.342
9	0.0855	0.12825	0.171	0.21375	0.2565	0.29925	0.342	0.38475
10	0.095	0.145	0.19	0.24	0.285	0.335	0.38	0.43
12	0.115	0.17	0.23	0.285	0.345	0.4	0.46	0.515
14	0.135	0.2	0.27	0.335	0.4	0.47	0.535	0.605
16	0.155	0.23	0.305	0.385	0.46	0.535	0.61	0.69
18	0.17	0.26	0.345	0.43	0.515	0.605	0.69	0.775
20	0.19	0.285	0.38	0.48	0.575	0.67	0.765	0.86
22	0.21	0.315	0.42	0.525	0.63	0.735	0.84	0.945
24	0.23	0.345	0.46	0.575	0.69	0.805	0.92	1.035
26	0.25	0.375	0.495	0.62	0.745	0.87	0.995	1.12
28	0.27	0.4	0.535	0.67	0.805	0.935	1.07	1.205
30	0.285	0.43	0.575	0.715	0.86	1.005	1.145	1.29
32	0.305	0.46	0.61	0.76	0.92	1.07	1.225	1.375

**TREE, BUSH, and VINE CROPS**

**BANANA and PLANTAIN – soil treatment**

<b>Pests Controlled</b>	<b>Rate: Fluid ounces per acre</b>
Aphids, Leafhoppers	8.0-16.0
<b>Pests Suppressed</b>	<b>Rate: Fluid ounces per acre</b>
Scales	8.0-16.0
<b>Restrictions:</b>	
<ul style="list-style-type: none"> <li>Pre-Harvest Interval (PHI): 0 day</li> <li>Maximum MANA Alias 4F allowed per crop season when making soil applications: 16.0 fluid ounces per Acre (0.5 lb AI per A)</li> </ul>	
<b>Applications:</b> Apply specified dosage in the following method:	
<ul style="list-style-type: none"> <li>Chemigation into root-zone through low-pressure drip, trickle, micro-sprinkler, or equivalent equipment.</li> </ul>	

**BANANA and PLANTAIN – foliar treatment**

<b>Pests Controlled</b>	<b>Rate: Fluid ounces per acre</b>
Aphids, Leafhoppers, Thrips	3.2
<b>Restrictions:</b>	
<ul style="list-style-type: none"> <li>Pre-Harvest Interval (PHI): 0 day</li> <li>Minimum interval between applications: 14 days</li> <li>Maximum MANA Alias 4F allowed per crop season when making foliar applications: 16.0 fluid ounces per acre (0.5 lb AI per acre)</li> </ul>	
<b>Applications:</b> Apply specified rate per acre as a broadcast or directed foliar spray to infested area as pest populations begin to build. Thorough uniform coverage is necessary to achieve optimum control. Use a spray adjuvant improve coverage. MANA Alias 4F may not knockdown established and heavy insect populations. Two applications may be required to achieve control. Scout fields and retreat if needed. Tank mix MANA Alias 4F with other insecticides for knockdown of pests or from improved control of other pests. Aerial application of MANA Alias 4F may result in slower activity and reduced control relative to results from ground application. For tree and vine crops, application rates are based on full-size, mature trees or vines. Crops contained within certain crop groups recognized by EPA are subject to change. Refer to EPA website ( <a href="http://www.epa.gov">www.epa.gov</a> ) for latest crop groups.	

21  
29

Apply specified dosage as a broadcast or directed spray to infested area insuring thorough coverage. MANA Alias 4F may be applied through properly calibrated ground or aerial application equipment. Aerial applications of MANA Alias 4F may result in slower activity and reduced control relative to results from ground application.  
To improve coverage and pest control, add a finished spray solution of an organosilicone adjuvant at a rate not to exceed 2.0 fluid ounces per 100 gallons.

**BUSHBERRY – soil treatment**

**Crops of Crop Subgroup 13B Including:** Blueberry, Currant, Elderberry, Gooseberry, Huckleberry, Juneberry, Ligonberry, Salal

Pests Controlled	Rate: Fluid ounces per acre
Japanese beetle: (adults, feeding on foliage) White grub complex: (grubs of Asiatic garden beetle, European and Masked chafer, Japanese beetle and Oriental beetle)	8.0-16.0
<b>Restrictions:</b> <ul style="list-style-type: none"> <li>Pre-Harvest Interval (PHI): 7 days</li> <li>Maximum MANA Alias 4F allowed per season when making soil applications: 16.0 fluid ounces/Acre (0.5 lb AI per acre)</li> <li>Do not apply pre-bloom or during bloom or when bees are actively foraging.</li> </ul> <b>Applications:</b> Apply specified dosage in one of the following methods: <ul style="list-style-type: none"> <li>Chemigation into root-zone through low-pressure drip, trickle, micro-sprinkler, or equivalent equipment.</li> <li>18-inch band on each side of the row followed by irrigation immediately after application.</li> </ul> For optimal grub control, apply MANA Alias 4F to control 1 <sup>st</sup> or 2 <sup>nd</sup> instar larvae. Make application post-bloom up to 7 days prior to harvest, or post-harvest until October 1 <sup>st</sup> . For optimum control of Japanese beetle larvae, make applications from June 1 to July 15. Application to grass covered rows, row middles, drive lanes, headlands, and other grassy areas in and around the berry field will control resident grub populations. Applications directed to the root-zone will help protect berry plant roots from grub feeding. Apply MANA Alias 4F to moist soil. If necessary, apply one hour of irrigation water immediately before application of MANA Alias 4F. To ensure maximum efficacy of soil surface spray, apply 1/2 to 1 inch of irrigation water or rainfall within 24 hours of application of MANA Alias 4F to facilitate movement into the soil and into the root-zone.	

**BUSHBERRY – foliar treatment**

**Crops of Crop Subgroup 13B Including:** Blueberry, Currant, Elderberry, Gooseberry, Huckleberry, Juneberry, Ligonberry, Salal

Pests Controlled	Rate: Fluid ounces per acre
Aphids, Leafhoppers/Sharpshooters	1.2-1.6
Blueberry maggot, Japanese beetle (adults), Thrips (foliage feeding thrips only)	2.4-3.2
<b>Restrictions:</b> <ul style="list-style-type: none"> <li>Pre-Harvest Interval (PHI): 3 days</li> <li>Minimum interval between applications: 7 days</li> <li>Maximum MANA Alias 4F allowed per crop season when making foliar applications: 16 fluid ounces per acre (0.5 lb AI per acre)</li> <li>Maximum number of MANA Alias 4F applications per crop season when making foliar applications: 5</li> <li>Minimum application volume (water): 20.0 GPA-ground, 5.0 GPA-aerial.</li> <li>Do not apply pre-bloom or during bloom or when bees are actively foraging.</li> </ul> <b>Applications:</b> Apply specified rate per acre as a broadcast or directed foliar spray to infested area as pest populations begin to build. Thorough uniform coverage is necessary to achieve optimum control. Use a spray adjuvant to improve coverage. MANA Alias 4F may not knockdown established and heavy insect populations. Two applications may be required to achieve control. Scout fields and retreat if needed. Tank mix MANA Alias 4F with other insecticides for knockdown of pests or from improved control of other pests. Aerial application of MANA Alias 4F may result in slower activity and reduced control relative to results from ground application. For tree and vine crops, application rates are based on full-size, mature trees or vines. Crops contained within certain crop groups recognized by EPA are subject to change. Refer to EPA website ( <a href="http://www.epa.gov">www.epa.gov</a> ) for latest crop groups.	

**CANEBERRY – soil treatment**

For use only in CA

**Crops of Crop Subgroup 13A including:**

**Blackberry** (*Rubus eubatus*, including bingleberry, black satin berry, boysenberry, Cherokee blackberry, Chesterberry, Cheyenne blackberry, coryberry, darrowberry, dewberry, Dirksen thornless berry, Himalayaberry, hullberry, Lavacaberry, Loganberry, lowberry, Lucretiaberry, mammoth blackberry, marionberry, nectarberry, olallieberry, Oregon evergreen berry, phenomenalberry, rangeberry, ravenberry, rossberry, Shawnee blackberry, youngberry, and varieties and/or hybrids of these)  
**Raspberry** (black and red, *Rubus occidentalis*, *Rubus strigosus*, *Rubus idaeus*)

Pests Controlled	Rate: Fluid ounces per acre
Aphids, Leafhoppers, Whiteflies	8.0-16.0
Rednecked cane borer	12.0-16.0
<b>Pests Suppressed</b>	
Thrips (foliage feeding thrips only)	8.0-16.0
<b>Restrictions:</b> <ul style="list-style-type: none"> <li>Pre-Harvest Interval (PHI): 7 days</li> <li>Maximum MANA Alias 4F allowed per season when making soil applications: 16.0 fluid ounces/Acre (0.5 lb AI per acre)</li> <li>Do not apply pre-bloom or during bloom or when bees are actively foraging.</li> </ul> <b>Soil Application:</b> Apply specified dosage in one of the following methods: <ul style="list-style-type: none"> <li>Chemigation into root-zone through low-pressure drip, trickle, micro-sprinkler, or equivalent equipment.</li> <li>Basal, soil drench in a minimum of 500 gallons solution per acre.</li> </ul>	

22  
29

**CITRUS (Containerized) – soil treatment**

**Crops of Crop Group 10 Including:** Calamondin, Citrus citron, Citrus hybrids (includes chironja, tangelo, and tangor), Grapefruit, Kumquat, Lemon, Lime, Mandarin (tangerine), Pummelo, Orange (sweet and sour), Satsuma mandarin, Tangelo, White sapote (*Casimiroa* spp), and other cultivars and/or hybrids of these.

<b>Pests Controlled</b>	<b>Rate: ml/ft<sup>3</sup> container media</b>
Aphid, Asian citrus psyllid, Blackfly, Citrus leafminer, Leafhoppers/Sharpshooters, Mealybugs, Scales, Whiteflies	0.37
Citrus root weevil (larval complex)	0.62-1.2
<b>Pests Suppressed</b>	<b>Rate: ml/ft<sup>3</sup> container media</b>
Citrus thrips (foliage feeding thrips only)	1.25
<b>Application:</b> Determine volume of container and calculate dosage necessary to treat container. Apply calculated dosage of MANA Alias 4F per container as a soil drench or through low-pressure drip or trickle irrigation water. Use sufficient carrier volume to ensure thorough uniform distribution throughout the media without loss of gravitational water from the container. For optimal results, make treatment at planting prior to insect infestation. Retreat if necessary. For control of larvae of the citrus root weevil complex, make application prior to neonate larvae entering potting media. Utilize higher dosage for heavy infestations.	

**CITRUS (Field) – soil treatment**

**Crops of Crop Group 10 Including:** Calamondin, Citrus citron, Citrus hybrids (includes chironja, tangelo, and tangor), Grapefruit, Kumquat, Lemon, Lime, Mandarin (tangerine), Pummelo, Orange (sweet and sour), Satsuma mandarin, Tangelo, White sapote (*Casimiroa* spp.), and other cultivars and/or hybrids of these.

<b>Pests Controlled</b>	<b>Rate: Fluid ounces per acre</b>
Aphids, Asian citrus psyllid, Blackfly, Citrus leafminer, Leafhoppers/Sharpshooters, Mealybugs, Scales, Termites (FL only), Whiteflies	8.0-16.0
<b>Pests/Diseases Suppressed</b>	<b>Rate: Fluid ounces per acre</b>
Citrus nematode, Symptoms of: Citrus tristeza virus (CTV) through vector control, Citrus yellows, Thrips (foliage feeding thrips only)	16.0
<b>Restrictions:</b>	
<ul style="list-style-type: none"> <li>• Pre-Harvest Interval (PHI): 0 day</li> <li>• Maximum MANA Alias 4F allowed per season when making soil applications: 16.0 fluid ounces/Acre (0.5 lb AI per Acre)</li> </ul>	
<b>Applications:</b> Apply specified dosage in one of the following methods:	
<ul style="list-style-type: none"> <li>• Chemigation into root-zone through low-pressure drip, trickle, micro-sprinkler, or equivalent equipment. For optimum results, apply to newly planted trees or those previously trained to drip, trickle or micro-sprinkler irrigation. Lightly pre-wet soil to break soil surface tension prior to applications of MANA Alias 4F. Chemigation application can be made separate to normal irrigation but followed by 10 to 20 minutes of additional watering to move MANA Alias 4F into root-zone. Allow 24 hours before initiating subsequent irrigations.</li> <li>• Soil surface band spray on both sides of the tree. Overlap bands at the tree base to create a continuous band within the drip-line area of the tree, to be followed immediately with light sprinkler irrigation sufficient to move the product into the upper portion of the root-zone. This method is suitable for very coarse soils with 0.75% organic matter or less.</li> <li>• Drench to base of tree not exceeding one-quart total solution per tree immediately around trunk of tree and extending outward covering the entire fibrous root system of the tree. Use only on trees up to 8 feet tall.</li> <li>• For control of existing termite infestations, apply specified dosage in 1 to 4 quarts of total solution volume, depending on size of tree, as a drench application to the basal portion of the tree trunk and surrounding soil in the immediate vicinity of the tree trunk.</li> <li>• For suppression of citrus nematode, apply specified dosage through low-pressure chemigation or soil surface band spray only, ensuring complete coverage of the root system and utilizing application directions stated above for the respective application method. Repeated and regular use of MANA Alias 4F over several consecutive growing seasons provides the greatest degree of nematode suppression and yields the greatest plant response.</li> </ul>	

**CITRUS (Field) – foliar treatment**

**Crops of Crop Group 10 Including:** Calamondin, Citrus citron, Citrus hybrids (includes chironja, tangelo, and tangor), Grapefruit, Kumquat, Lemon, Lime, Mandarin (tangerine), Pummelo, Orange (sweet and sour), Satsuma mandarin, Tangelo, White sapote (*Casimiroa* spp.), and other cultivars and/or hybrids of these.

<b>Pests Controlled</b>	<b>Rate: Fluid ounces per acre</b>
Aphids, Asian citrus psyllid, Blackfly, Leafhoppers/Sharpshooters, Leafminers, Mealybugs, Scales, Whiteflies	4.0-8.0 (depending on tree size, target pest and infestation pressure)
<b>Pest Suppressed</b>	<b>Rate: Fluid ounces per acre</b>
Thrips (foliage feeding thrips only)	4.0-8.0
<b>Restrictions:</b>	
<ul style="list-style-type: none"> <li>• Pre-Harvest Interval (PHI): 0 day</li> <li>• Minimum interval between applications: 10 days</li> <li>• Maximum MANA Alias 4F allowed per season when making foliar applications: 16 fluid ounces per acre (0.5 lb AI per acre)</li> <li>• Do not apply during bloom or within 10 days prior to bloom or when bees are actively foraging.</li> </ul>	
<b>Applications:</b> Scales – time applications to the crawler stage. Treat each generation. Apply specified rate per acre as a broadcast or directed foliar spray to infested area as pest populations begin to build. Thorough uniform coverage is necessary to achieve optimum control. Use a spray adjuvant to improve coverage. MANA Alias 4F may not knockdown established and heavy insect populations. Two applications may be required to achieve control. Scout fields and retreat if needed. Tank mix MANA Alias 4F with other insecticides for knockdown of pests or from improved control of other pests. Aerial application of MANA Alias 4F may result in slower activity and reduced control relative to results from ground application. For tree and vine crops, application rates are based on full-size, mature trees or vines. Crops contained within certain crop groups recognized by EPA are subject to change. Refer to EPA website ( <a href="http://www.epa.gov">www.epa.gov</a> ) for latest crop groups.	

23  
19

**COFFEE – soil treatment**

<b>Pests Controlled</b>	<b>Rate: Fluid ounces per acre</b>
Aphids, Leafhoppers, Leafminers	8.0-16.0
<b>Pest Suppressed</b>	<b>Rate: Fluid ounces per acre</b>
Scales	8.0-16.0
<b>Restrictions:</b>	
<ul style="list-style-type: none"> <li>• Pre-Harvest Interval (PHI): 7 days</li> <li>• Maximum MANA Alias 4F allowed per season when making soil applications: 16.0 fluid ounces per Acre (0.5 lb AI per acre)</li> <li>• Do not apply pre-bloom or during bloom or when bees are actively foraging.</li> </ul>	
<b>Applications:</b> Apply specified dosage in one of the following methods:	
<ul style="list-style-type: none"> <li>• Chemigation into root-zone through low-pressure drip, trickle, micro-sprinkler, or equivalent equipment.</li> <li>• Subsurface side-dress shanked into the root-zone on both side of the plants followed by irrigation.</li> <li>• Basal, soil drench in sufficient water to insure incorporation into the root-zone followed by irrigation.</li> </ul>	

**COFFEE – foliar treatment**

<b>Pests Controlled</b>	<b>Rate: Fluid ounces per acre</b>
Aphids, Leafhoppers, Whiteflies	3.2
<b>Pest Suppressed</b>	<b>Rate: Fluid ounces per acre</b>
Scales	3.2
<b>Restrictions:</b>	
<ul style="list-style-type: none"> <li>• Pre-Harvest Interval (PHI): 7 days</li> <li>• Minimum interval between applications: 7 days</li> <li>• Maximum MANA Alias 4F allowed per crop season when making foliar applications: 16 fluid ounces per Acre (0.5 lb AI/Acre)</li> <li>• Do not apply pre-bloom or during bloom or when bees are actively foraging.</li> </ul>	
<b>Applications:</b> Apply specified dosage as a broadcast or directed spray to infested area insuring thorough coverage. Apply MANA Alias 4F through properly calibrated ground or aerial application equipment. Aerial application of MANA Alias 4F may result in slower activity and reduced control relative to results from ground application.	
Apply specified rate per acre as a broadcast or directed foliar spray to infested area as pest populations begin to build. Thorough uniform coverage is necessary to achieve optimum control. Use a spray adjuvant to improve coverage. MANA Alias 4F may not knockdown established and heavy insect populations. Two applications may be required to achieve control. Scout fields and retreat if needed. Tank mix MANA Alias 4F with other insecticides for knockdown of pests or from improved control of other pests. Aerial application of MANA Alias 4F may result in slower activity and reduced control relative to results from ground application. For tree and vine crops, application rates are based on full-size, mature trees or vines. Crops contained within certain crop groups recognized by EPA are subject to change. Refer to EPA website ( <a href="http://www.epa.gov">www.epa.gov</a> ) for latest crop groups.	

**CRANBERRY – soil treatment**

<b>Pests Controlled</b>	<b>Rate: Fluid ounces per acre</b>
Rootgrubs ( <i>Scarabaeidae</i> ), Rootworms ( <i>Chrysomelidae</i> )	8.0-16.0
<b>Restrictions:</b>	
<ul style="list-style-type: none"> <li>• Pre-Harvest Interval (PHI): 30 days</li> <li>• Maximum MANA Alias 4F allowed per season when making soil applications: 16.0 fluid ounces/Acre (0.5 lb AI/Acre)</li> <li>• Do not apply pre-bloom or during bloom or when bees are actively foraging.</li> </ul>	
<b>Applications:</b> Apply MANA Alias 4F to moist soil. Apply specified dosage in one of the following methods:	
<ul style="list-style-type: none"> <li>• As a soil spray (ground application) directed to the root and crown area using a minimum of 20 gal of water per acre.</li> <li>• As a chemigation application with 600 to 1000 gal water.</li> </ul>	
Immediately upon application, incorporate MANA Alias 4F into root-zone by 0.1-0.3 inches water/Acre, either with the chemigation application or through irrigation/rainfall if not applied through chemigation. Inadequate incorporation within 24 hours of application may result in reduced control.	
<b>Rootgrubs and Rootworms</b>	
Make application post-bloom immediately after bees are removed. Target applications to early instar larvae. MANA Alias 4F has not been tested for crop response in tank mixes with other registered fungicides or insecticides. If tank mixing is desired, premix a sample of the MANA Alias 4F and the desired fungicide or insecticide partner at labeled rates and apply to a small area. Evaluate crop response within 48 hours and for at least two weeks prior to utilizing the tank mix on larger acreage. If crop injury results from the premix test, do not apply the tank mix to larger acreage.	

**GRAPE – soil treatment**

Including: American bunch grape, Muscadine grape and Vinifera grape

<b>Pests Controlled</b>	
European fruit lecanium, Leafhoppers/Sharpshooters, Mealybugs, <i>Phylloxera</i> * spp	8.0-16.0
<b>Pest/Disease Suppressed</b>	<b>Rate: Fluid ounces per acre</b>
Grapeleaf skeletonizer, Nematodes, Pierce's disease	12.0-16.0
<b>Restrictions:</b>	
<ul style="list-style-type: none"> <li>• Pre-Harvest Interval (PHI): 30 days</li> <li>• Maximum MANA Alias 4F allowed per season when making soil applications: 16.0 fluid ounces/Acre (0.5 lb AI/Acre)</li> </ul>	
<b>Applications:</b> Apply specified dosage in one of the following methods:	
<ul style="list-style-type: none"> <li>• Chemigation into root-zone through low-pressure drip, trickle, micro-sprinkler, or equivalent equipment.</li> <li>• Subsurface side-dress shanked into the root-zone on both sides of the plants followed by irrigation.</li> <li>• Hill drench in sufficient water to insure incorporation into the root-zone followed by irrigation.</li> <li>• For suppression of nematodes, apply 7 fluid ounces in a single application or two 3.5 fluid ounce applications on a 30 to 45 day interval. Apply treatment(s) only by 1) chemigation into root-zone through above ground low-pressure drip, trickle, micro-sprinkler, or equivalent</li> </ul>	

24  
29

equipment; or 2) French plow technique, followed immediately by sufficient irrigation to move the product into the entire root-zone of the plant. Repeated and regular use of MANA Alias 4F over several consecutive growing seasons provides the greatest degree of nematode suppression and yields the greatest plant response.

For optimum results, make application(s) between bud-break and the pea-berry stage. Use a total of 14.7 fluid ounces/Acre under any of the following conditions:

1. Where vigorous vine growth is expected;
2. In warmer growing areas;
3. Where mealybug and European fruit lecanium populations are expected to be heavy;
4. Where vine populations exceed 600 per acre, or;
5. For suppression of nematodes.

\*Repeated and regular use of MANA Alias 4F over several, consecutive growing seasons controls existing *Phylloxera* infestations over time or prevents *Phylloxera* from becoming established.

**GRAPE – foliar treatment**

Including: American bunch grape, Muscadine grape and Vinifera grape

Pests Controlled	Rate: Fluid ounces per acre
Leafhoppers/Sharpshooters, Mealybugs	1.2-1.6
Grapeleaf skeletonizer	1.5-1.6

**Restrictions:**

- Pre-Harvest Interval (PHI): 0 days
- Minimum interval between applications: 14 days
- Maximum MANA Alias 4F allowed per season when making foliar applications: 3.2 fluid ounces/Acre (0.1 lb AI/Acre)

Apply MANA Alias 4F by ground application only.

**Applications:** Apply specified rate per acre as a broadcast or directed foliar spray to infested area as pest populations begin to build. Thorough uniform coverage is necessary to achieve optimum control. Use a spray adjuvant to improve coverage. MANA Alias 4F may not knockdown established and heavy insect populations. Two applications may be required to achieve control. Scout fields and retreat if needed. Tank mix MANA Alias 4F with other insecticides for knockdown of pests or from improved control of other pests. Aerial application of MANA Alias 4F may result in slower activity and reduced control relative to results from ground application. For tree and vine crops, application rates are based on full-size, mature trees or vines. Crops contained within certain crop groups recognized by EPA are subject to change. Refer to EPA website ([www.epa.gov](http://www.epa.gov)) for latest crop groups.

**HOP – soil treatment**

Pest Controlled	Rate: Fluid ounces per acre
Aphids	9.6

**Restrictions:**

- Pre-Harvest Interval (PHI): 60 days
- Maximum MANA Alias 4F allowed per season when making soil applications: 9.6 fluid ounces/Acre (0.3 lb AI/Acre)

**Applications:** Apply specified dosage in one of the following methods:

- Chemigation into root-zone through low-pressure drip, trickle, micro-sprinkler, or equivalent equipment.
- Subsurface side-dress shanked into the root-zone on both sides of the plants followed by irrigation.
- Hill drench in sufficient water to insure incorporation into the root-zone followed by irrigation.

Use the higher dosage where extended residual control is desired or for treating larger vines or vines with dense foliage volume.

**HOP – foliar treatment**

Pest Controlled	Rate: Fluid ounces per acre
Aphids	3.2

**Restrictions:**

- Pre-Harvest Interval (PHI): 28 days
- Minimum interval between applications: 21 days
- Maximum MANA Alias 4F allowed per season when making foliar applications: 9.6 fluid ounces/Acre (0.3 lb AI/Acre)

**Applications:** Apply specified rate per acre as a broadcast or directed foliar spray to infested area as pest populations begin to build. Thorough uniform coverage is necessary to achieve optimum control. Use a spray adjuvant to improve coverage. MANA Alias 4F may not knockdown established and heavy insect populations. Two applications may be required to achieve control. Scout fields and retreat if needed. Tank mix MANA Alias 4F with other insecticides for knockdown of pests or from improved control of other pests. Aerial application of MANA Alias 4F may result in slower activity and reduced control relative to results from ground application. For tree and vine crops, application rates are based on full-size, mature trees or vines. Crops contained within certain crop groups recognized by EPA are subject to change. Refer to EPA website ([www.epa.gov](http://www.epa.gov)) for latest crop groups.

**POME FRUIT – soil treatment**

**Crops of Crop Group 11 Including:** Apple, Crabapple, Loquat, Mayhaw, Pear (including Oriental pear), Quince

Pests Controlled	Rate: Fluid ounces per acre
Aphids (including Woolly apple aphid), Leafhoppers	8.0-12.0

**Restrictions:** Pre-Harvest Interval (PHI): 21 days

- Maximum MANA Alias 4F allowed per season when making soil applications: 12.0 fluid ounces/Acre (0.38 lb AI/Acre)
- Do not apply pre-bloom or during bloom or when bees are actively foraging

**Applications:** Apply specified dosage in the following method:

- Chemigation into the root-zone through low-pressure drip, trickle, micro-sprinkler, or equivalent equipment.



25  
29

**POME FRUIT – foliar treatment**

**Crops of Crop Group 11 Including:** Apple, Crabapple, Loquat, Mayhaw, Pear (including Oriental pear), Quince

Pests Controlled	Rate: Fluid ounces per acre
Leafhoppers	1.6-3.2
Aphids (except Woolly apple aphid), Apple maggot, Leafminers, San Jose scale	3.2
<b>FOR PEAR ONLY:</b> Mealybugs, Pear psylla	8
<b>Restrictions:</b> <ul style="list-style-type: none"> <li>• Pre-Harvest Interval (PHI): 7 days</li> <li>• Minimum interval between applications: 10 days</li> <li>• Maximum MANA Alias 4F allowed per season when making foliar applications: 16 fluid ounces per acre (0.5 lb AI/Acre)</li> <li>• Do not apply pre-bloom or during bloom or when bees are actively foraging</li> </ul> <b>Applications:</b> Combine applications targeting apple maggots with an approved sticker at the manufacturer's specified rates. Apply specified rate per acre as a broadcast or directed foliar spray to infested area as pest populations begin to build. Thorough uniform coverage is necessary to achieve optimum control. Use a-spray adjuvant to improve coverage. MANA Alias 4F may not knockdown established and heavy insect populations. Two applications may be required to achieve control. Scout fields and retreat if needed. Tank mix MANA Alias 4F with other insecticides for knockdown of pests or from improved control of other pests. Aerial application of MANA Alias 4F may result in slower activity and reduced control relative to results from ground application. For tree and vine crops, application rates are based on full-size, mature trees or vines. Crops contained within certain crop groups recognized by EPA are subject to change. Refer to EPA website ( <a href="http://www.epa.gov">www.epa.gov</a> ) for latest crop groups.	

**POMEGRANATE – soil treatment**

Pests Controlled	Rate: Fluid ounces per acre
Aphids, Leafhoppers/Sharpshooters, Whiteflies	8.0-16.0
<b>Restrictions:</b> <ul style="list-style-type: none"> <li>• Pre-Harvest Interval (PHI): 0 day</li> <li>• Maximum MANA Alias 4F allowed per season when making soil applications: 16.0 fluid ounces/Acre (0.5 lb AI per acre)</li> <li>• Do not apply pre-bloom or during bloom or when bees are actively foraging.</li> </ul> <b>Applications:</b> Apply specified dosage in the following method: <ul style="list-style-type: none"> <li>• Chemigation into the root-zone through low-pressure drip, trickle, micro-sprinkler, or equivalent equipment.</li> </ul>	

**POMEGRANATE – foliar treatment**

Pests Controlled	Rate: Fluid ounces per acre
Aphids, Leafhoppers/Sharpshooters, Whiteflies	3.2
<b>Pests suppressed</b>	
Scales	3.2
<b>Restrictions:</b> <ul style="list-style-type: none"> <li>• Pre-Harvest Interval (PHI): 7 days</li> <li>• Minimum interval between applications: 7 days</li> <li>• Maximum MANA Alias 4F allowed per season when making foliar applications: 9.6 fluid ounces per acre (0.3 lb AI per acre)</li> <li>• Do not apply pre-bloom or during bloom or when bees are actively foraging</li> </ul> <b>Applications:</b> Apply specified rate per acre as a broadcast or directed foliar spray to infested area as pest populations begin to build. Thorough uniform coverage is necessary to achieve optimum control. Use a spray adjuvant to improve coverage. MANA Alias 4F may not knockdown established and heavy insect populations. Two applications may be required to achieve control. Scout fields and retreat if needed. Tank mix MANA Alias 4F with other insecticides for knockdown of pests or from improved control of other pests. Aerial application of MANA Alias 4F may result in slower activity and reduced control relative to results from ground application. For tree and vine crops, application rates are based on full-size, mature trees or vines. Crops contained within certain crop groups recognized by EPA are subject to change. Refer to EPA website ( <a href="http://www.epa.gov">www.epa.gov</a> ) for latest crop groups.	

**STONE FRUIT – soil treatment**

**Crops of Crop Group 12 Including:** Apricot, Cherry (including sweet and tart), Nectarine, Peach, Plum (including Chickasaw, Damson, and Japanese), Plumcot, Prune (fresh and dried)

In-field, Soil Application	
Pests Controlled	Rate: Fluid ounces per acre
Aphids (including Woolly apple aphid), Leafhoppers	8.0-12.0
<b>Restrictions:</b> <ul style="list-style-type: none"> <li>• Pre-Harvest Interval (PHI): 21 days</li> <li>• Maximum MANA Alias 4F allowed per season when making soil applications: 12.0 fluid ounces/Acre (0.38 lb AI/Acre)</li> <li>• Do not apply pre-bloom or during bloom or when bees are actively foraging.</li> </ul> <b>Applications:</b> Apply specified dosage in the following method: <ul style="list-style-type: none"> <li>• Chemigation into root-zone through low-pressure drip, trickle, micro-sprinkler, or equivalent equipment.</li> </ul>	
Pre-plant, Root Dip Application	
Pest Controlled	Rate fluid ounces per 10 gallons root-dip solution
Black peach aphid (infesting roots)	1.0
Mix MANA Alias 4F at 1.0 fluid ounces per 10 gallons of water. Thoroughly wet bare-root transplant to slightly above the graft union by soaking roots in the MANA Alias 4F solution for up to 5 minutes. Allow solution to dry on roots and transplant trees as soon as possible following treatment.	

26  
29

**STONE FRUIT – foliar treatment**

**Crops of Crop Group 12 Including:** Apricot, Cherry (including sweet and tart), Nectarine, Peach, Plum (including Chickasaw, Damson, and Japanese), Plumcot, Prune (fresh and dried)

Pests Controlled	Rate: Fluid ounces per Acre
Aphids, Green June beetle, Japanese beetle, Leafhoppers/Sharpshooters, Plant bugs, Rose chafer, San Jose scale	1.6-3.2
Cherry fruit fly	2.4-3.2
Pests Suppressed	Rate: Fluid ounces per Acre
Plum curculio, Stink bugs	3.2
<b>Restrictions for Apricot, Nectarine, Peach:</b>	
<ul style="list-style-type: none"> <li>• Pre-Harvest Interval (PHI): 0 day</li> <li>• Minimum interval between applications: 7 days</li> <li>• Maximum MANA Alias 4F allowed per season when making foliar applications: 9.6 fluid ounces/Acre (0.3 lb AI/Acre)</li> <li>• Minimum application volume (water): 50 GPA – ground application, 25 GPA- aerial application.</li> <li>• Do not apply pre-bloom or during bloom or when bees are actively foraging.</li> </ul>	
<b>Restrictions for Cherries, Plums, Plumcot, Prune:</b>	
<ul style="list-style-type: none"> <li>• Pre-Harvest Interval (PHI): 7 days</li> <li>• Minimum interval between applications: 10 days</li> <li>• Maximum MANA Alias 4F allowed per season when making foliar applications: 16.0 fluid ounces/Acre (0.5 lb AI/Acre)</li> <li>• Minimum application volume (water): 50 GPA – ground application, 25 GPA- aerial application.</li> <li>• Do not apply pre-bloom or during bloom or when bees are actively foraging.</li> </ul>	
<p><b>Applications:</b> Apply specified rate per acre as a broadcast or directed foliar spray to infested area as pest populations begin to build. Thorough uniform coverage is necessary to achieve optimum control. Use a spray adjuvant to improve coverage. MANA Alias 4F may not knockdown established and heavy insect populations. Two applications may be required to achieve control. Scout fields and retreat if needed. Tank mix MANA Alias 4F with other insecticides for knockdown of pests or from improved control of other pests. Aerial application of MANA Alias 4F may result in slower activity and reduced control relative to results from ground application. For tree and vine crops, application rates are based on full-size, mature trees or vines. Crops contained within certain crop groups recognized by EPA are subject to change. Refer to EPA website (<a href="http://www.epa.gov">www.epa.gov</a>) for latest crop groups.</p>	

**TREE NUTS – soil treatment**

**Crops of Crop Group 14 Including:** Almond, Beechnut, Brazil nut, Butternut, Cashew, Chestnut, Chinquapin, Filbert, Hickory nut, Macadamia nut, Pecan, Pistachio, Walnut [black and English]

Pests Controlled	Rate: Fluid ounces per acre
Aphids, Leafhoppers/Sharpshooters, Mealybugs, Spittlebugs, Termites, Whiteflies	8.0-16.0
Pests/Diseases Suppressed	Rate: Fluid ounces per acre
Pecan scab (from reduction in honeydew deposition)	8.0-16.0
Thrips (foliage-feeding thrips only)	16.0
<b>Restrictions:</b>	
<ul style="list-style-type: none"> <li>• Pre-Harvest Interval (PHI): 7 days</li> <li>• Maximum MANA Alias 4F allowed per season when making soil applications: 16.0 fluid ounces/Acre (0.5 lb AI/Acre)</li> <li>• Do not apply pre-bloom or during bloom or when bees are actively foraging.</li> </ul>	
<p><b>Applications:</b> Apply specified dosage prior to or at onset of pest infestation in one of the following methods:</p> <ul style="list-style-type: none"> <li>• Chemigation into root-zone through low-pressure drip, trickle, micro-sprinkler, or equivalent irrigation equipment. Pre-wet soil prior to applications of MANA Alias 4F and allow soil to dry following application and prior to subsequent irrigation.</li> <li>• Emitter or spot application in a minimum of 4 fluid ounces of mixture per emitter site.</li> <li>• Shank or subsurface side-dress, injected to a depth just above or just within the root zone and between the trunk and drip line of the tree canopy. Product should be applied in a minimum of 10 gallons per acre using multiple shanks on both sides of trees. Ensure product placement is below sod or orchard floor debris. Irrigation covering entire treated area should follow within 48 hours to promote uptake by root system.</li> <li>• For control of termites, apply specified dosage to slightly moist soil as a high-volume drench to the basal portion of the tree trunk and surrounding soil in the immediate vicinity of the tree trunk. Utilize sufficient carrier volume to penetrate the soil to a depth of 18-24 inches to obtain optimum control. Allow soil to dry following treatment and prior to applying any irrigation.</li> </ul>	
<p><b>Remarks:</b> Use the higher rates when applied by shank or subsurface side dress, used on larger trees, soils with high clay content, for high plant populations, and/or where extended control is desired. Under some conditions, control may not occur for 14 or more days or until two (2) irrigations have been made. Applications made later in the season may result in reduced efficacy.</p>	

**TREE NUTS – foliar treatment**

**Crops of Crop Group 14 Including:** Almond, Beechnut, Brazil nut, Butternut, Cashew, Chestnut, Chinquapin, Filbert, Hickory nut, Macadamia nut, Pecan, Pistachio, Walnut [black and English]

Pests Controlled	Rate: Fluid ounces per acre
Aphids (except black pecan aphid), Leafhoppers/Sharpshooters, Phylloxera spp. (leaf infestations), Spittlebugs, Whiteflies	1.4-2.8
Black pecan aphid, Mealybugs, San Jose scale	3.2
<b>Restrictions:</b>	
<ul style="list-style-type: none"> <li>• Pre-Harvest Interval (PHI): 7 days</li> <li>• Minimum interval between applications: 6 days</li> <li>• Maximum MANA Alias 4F allowed per crop season when making foliar applications: 11.5 fluid ounces/Acre (0.36 lb AI/Acre)</li> <li>• Minimum application volume (water): 50 GPA- ground application, 25 GPA- aerial application</li> <li>• Do not apply pre-bloom or during bloom or when bees are actively foraging.</li> </ul>	

27  
29

**Applications:** Applications for control of San Jose scale should be timed according to crawler stage, treating each successive generation. Two applications on a 10 to 14 day interval may be required to achieve control. Apply specified rate per acre as a broadcast or directed foliar spray to infested area as pest populations begin to build. Thorough uniform coverage is necessary to achieve optimum control. Use a spray adjuvant to improve coverage. MANA Alias 4F may not knockdown established and heavy insect populations. Two applications may be required to achieve control. Scout fields and retreat if needed. Tank mix MANA Alias 4F with other insecticides for knockdown of pests or from improved control of other pests. Aerial application of MANA Alias 4F may result in slower activity and reduced control relative to results from ground application. For tree and vine crops, application rates are based on full-size, mature trees or vines. Crops contained within certain crop groups recognized by EPA are subject to change. Refer to EPA website ([www.epa.gov](http://www.epa.gov)) for latest crop groups.

**TROPICAL FRUIT – soil treatment**

Including: Acerola, Atemoya, Avocado, Birida, Black sapote, Canistel, Cherimoya, Custard apple, Feijoa, Jaboticaba, Guava, Llama, Longan, Lychee, Mamey sapote, Mango, Papaya, Passionfruit, Persimmon, Pulasan, Rambutan, Sapodilla, Soursop, Spanish lime, Star apple, Starfruit, Sugar apple, Wax jambu

<b>Pests Controlled</b>	<b>Rate: Fluid ounces per acre</b>
Aphids, Avocado lacebug, Leafhoppers, Whiteflies	12.0-16.0
<b>Pests Suppressed</b>	
Scales, Thrips (foliage feeding thrips only)	16.0
<b>Restrictions:</b>	
<ul style="list-style-type: none"> <li>• Pre-Harvest Interval (PHI): 6 days</li> <li>• Maximum MANA Alias 4F allowed per season when making soil applications: 16.0 fluid ounces/Acre (0.5 lb AI/A).</li> <li>• Do not apply pre-bloom or during bloom or when bees are actively foraging.</li> </ul>	
<b>Applications:</b> Apply specified dosage in the following method:	
<ul style="list-style-type: none"> <li>• Chemigation through low-pressure drip, trickle, micro-sprinkler, or equivalent equipment.</li> </ul>	

**TROPICAL FRUIT – foliar treatment**

Including: Acerola, Atemoya, Avocado, Birida, Black sapote, Canistel, Cherimoya, Custard apple, Feijoa, Jaboticaba, Guava, Llama, Longan, Lychee, Mamey sapote, Mango, Papaya, Passionfruit, Persimmon, Pulasan, Rambutan, Sapodilla, Soursop, Spanish lime, Star apple, Starfruit, Sugar apple, Wax jambu

<b>Pests Controlled</b>	<b>Rate: Fluid ounces per acre</b>
Aphids, Leafhoppers/Sharpshooters, Mealybugs, Thrips (foliage feeding thrips only), Whiteflies	3.2
<b>Pest Suppressed</b>	<b>Rate: Fluid ounces per acre</b>
Scales	3.2
<b>Restrictions:</b>	
<ul style="list-style-type: none"> <li>• Pre-Harvest Interval (PHI): 7 days</li> <li>• Minimum interval between applications: 10 days</li> <li>• Maximum MANA Alias 4F allowed per crop season when making foliar applications: 16.0 fluid ounces per acre (0.5 lb AI/A).</li> <li>• Do not apply pre-bloom or during bloom or when bees are actively foraging.</li> </ul>	
<b>Applications:</b> Apply specified rate per acre as a broadcast or directed foliar spray to infested area as pest populations begin to build. Thorough uniform coverage is necessary to achieve optimum control. Use a spray adjuvant to improve coverage. MANA Alias 4F may not knockdown established and heavy insect populations. Two applications may be required to achieve control. Scout fields and retreat if needed. Tank mix MANA Alias 4F with other insecticides for knockdown of pests or from improved control of other pests. Aerial application of MANA Alias 4F may result in slower activity and reduced control relative to results from ground application. For tree and vine crops, application rates are based on full-size, mature trees or vines. Crops contained within certain crop groups recognized by EPA are subject to change. Refer to EPA website ( <a href="http://www.epa.gov">www.epa.gov</a> ) for latest crop groups.	

**OTHER CROPS**

**CHRISTMAS TREE – soil treatment**

<b>Pests Controlled</b>	<b>Rate: Fluid ounces per acre</b>
White grub complex (damage from grubs of Asiatic garden beetle, European and Masked chafer, Japanese beetle and Oriental beetle)	8.0-16.0
<b>Restrictions:</b>	
<ul style="list-style-type: none"> <li>• Maximum MANA Alias 4F allowed per season when making soil applications: 16.0 fluid ounces/Acre (0.5 lb AI/Acre)</li> </ul>	
<b>Applications:</b> Soil incorporation and movement of MANA Alias 4F to the root-zone is required for activity. MANA Alias 4F can be incorporated most readily when applied to moist soil. Apply specified dosage in one of the following methods:	
<ul style="list-style-type: none"> <li>• Chemigation into root-zone through low-pressure drip, trickle, micro-sprinkler, or equivalent equipment.</li> <li>• 18-inch band on each side of the row (small trees) to full broadcast application (large trees) followed by rainfall or 0.25-1 inch of irrigation within 12 hours after application.</li> </ul>	
For optimal grub control, apply MANA Alias 4F during adult flight activity, or up to mid-July, when 1 <sup>st</sup> instar larvae are present.	

28  
29

**CHRISTMAS TREE – foliar treatment**

<b>Pests Controlled</b>	<b>Rate: Fluid ounces per acre</b>
Aphids, Adelgids, Sawflies	1.6-3.2
<b>Restrictions:</b>	
<ul style="list-style-type: none"> <li>• Minimum interval between applications: 7 days</li> <li>• Maximum MANA Alias 4F allowed per crop season when making foliar applications: 16.0 fluid ounces/Acre (0.5 lb AI/Acre)</li> </ul>	
<p><b>Applications:</b> Gall-forming adelgids – time applications to coincide with full bud-swell of earliest bud-breaking trees. Once galls form spraying will be ineffective.</p> <p>Apply specified rate per acre as a broadcast or directed foliar spray to infested area as pest populations begin to build. Thorough uniform coverage is necessary to achieve optimum control. Use a spray adjuvant to improve coverage. MANA Alias 4F may not knockdown established and heavy insect populations. Two applications may be required to achieve control. Scout fields and retreat if needed. Tank mix MANA Alias 4F with other insecticides for knockdown of pests or for improved control of other pests.</p>	

**POPLAR/COTTONWOOD<sup>1</sup> – soil treatment**

(includes members of the genus *Populus* grown for pulp or timber)

<b>Field Applications See details below for Cuttings/Whips Applications.</b>	
<b>Pests Controlled</b>	
Aphids, Cottonwood leaf beetle	8.0-16.0
<b>Pest Suppressed</b>	<b>Rate: Fluid ounces per acre</b>
<i>Phylloxera popularia</i>	8.0-16.0
<b>Restrictions:</b>	
<ul style="list-style-type: none"> <li>• Maximum MANA Alias 4F allowed at-plant per crop season: 16.0 fluid ounces/Acre (0.5 lb AI/Acre)</li> <li>• Do not apply pre-bloom or during bloom or when bees are actively foraging.</li> </ul> <p><sup>1</sup>Use not permitted in California unless otherwise directed by supplemental labeling.</p> <p><b>Applications:</b> Apply specified dosage in the following method:</p> <ul style="list-style-type: none"> <li>• Chemigation through low-pressure drip irrigation.</li> <li>• For narrow-row, cutting orchards/nurseries used for plant propagation, shank into root-zone followed by adequate irrigation to promote uptake. (Adequate irrigation depends on soil moisture level at application. Under dry conditions, use 0.25 inches/Acre).</li> </ul> <p>For Cottonwood leaf beetle, protection against damage will occur when application is made early-season, when the beetles first begin feeding. Larger trees may require earlier treatment as a result of slower uptake.</p> <p>For <i>Phylloxera</i>, apply early in the year from break of dormancy through May.</p>	
<b>Cutting/Whip Applications. See details above for Field Applications.</b>	
<b>Pest Controlled</b>	<b>Cutting/Whip Soaking Solution fluid ounces MANA Alias 4F Needed per 100 gallons</b>
Cottonwood leaf beetle	6.6-13.3 (unhydrated cuttings/whips) 13.3-20.0 (partially hydrated cuttings/whips)
<b>Pests Suppressed</b>	<b>Cutting/Whip Soaking Solution fluid ounces MANA Alias 4F Needed per 100 gallons</b>
Aphids, <i>Phylloxera popularia</i>	6.6-13.3 (unhydrated cuttings/whips) 13.3-20.0 (partially hydrated cuttings/whips)
<b>Restrictions:</b>	
<ul style="list-style-type: none"> <li>• Maximum MANA Alias 4F allowed at-plant per crop season: 16.0 fluid ounces/Acre (0.5 lb AI/Acre)</li> </ul> <p><sup>1</sup>Use not permitted in California unless otherwise directed by supplemental labeling.</p> <p><b>Applications:</b> Moisture content of cuttings/whips prior to application, the solution concentration, and the length of soaking interval interact to affect the amount of product absorbed into plant material. For a constant soaking interval of 24 hours, drier cuttings/whips absorb a higher quantity of solution and require a lower concentration. Conversely, more hydrated cuttings/whips absorb less solution and require a higher concentration. Soaking of cuttings/whips should occur in a covered container in absence of UV light. Not all <i>Populus</i> spp. clones/varieties/hybrids have been tested for crop safety. Without specific knowledge about a particular <i>Populus</i> spp. clone/variety/hybrid, Makhteshim Agan of North America, Inc. suggests that small numbers of cuttings/whips of each be treated and evaluated prior to commercial use.</p> <p>Apply MANA Alias 4F in one of the following cuttings/whips soaking methods:</p> <ul style="list-style-type: none"> <li>• For freshly cut (unhydrated) cuttings/whips, soak plant material in specified solution concentration for 24 hours prior to cold storage. After removal from cold storage, plant as needed.</li> <li>• For previously hydrated cuttings/whips removed from cold storage, allow plant material to reach room temperature and soak in specified solution concentration for 24 hours prior to planting.</li> </ul> <p>Take proper care in disposal of any residual soaking solution. Apply solution to existing trees or other registered crops as long as all product label precautions and restrictions are observed.</p>	

**POPLAR/COTTONWOOD<sup>1</sup> – foliar treatment**

(includes members of the genus *Populus* grown for pulp or timber)

<b>Pests Controlled</b>	<b>Rate: Fluid ounces per acre</b>
Aphids, Leaf beetles	1.6-3.2
<b>Restrictions:</b>	
<ul style="list-style-type: none"> <li>• Minimum interval between applications: 10 days</li> <li>• Maximum MANA Alias 4F allowed per crop season when making foliar applications: 16 fluid ounces per acre (0.5 lb AI per acre)</li> <li>• Do not apply pre-bloom or during bloom or when bees are actively foraging.</li> </ul> <p><sup>1</sup>Use not permitted in California unless otherwise directed by supplemental labeling.</p> <p><b>Applications:</b> Apply specified rate per acre as a broadcast or directed foliar spray to infested area as pest populations begin to build. Thorough uniform coverage is necessary to achieve optimum control. Use a spray adjuvant to improve coverage. MANA Alias 4F may not knockdown established and heavy insect populations. Two applications may be required to achieve control. Scout fields and retreat if needed. Tank mix MANA Alias 4F with other insecticides as knockdown of pests or for improved control of other pests.</p>	

29  
29

#### STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal or cleaning of equipment. Open dumping is prohibited.

**PESTICIDE STORAGE:** Keep pesticide in original container. Do not put concentrate or dilute into food or drink containers. Store in cool, dry place. Do not store diluted spray.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

#### CONTAINER HANDLING:

**Nonrefillable Container (five gallons or less):** Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. If recycling is not available, puncture and dispose of in a sanitary landfill or incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

**Nonrefillable Container (greater than five gallons):** Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. If recycling is not available, puncture and dispose of in a sanitary landfill or incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

**Refillable Container (greater than 55 gallons):** Refillable container. Refill this container with imidacloprid only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. For final disposal, offer for recycling or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

#### LIMITATION OF WARRANTY AND LIABILITY

Read the entire Directions for Use, Conditions of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following **CONDITIONS, DISCLAIMER OF WARRANTIES AND LIMITATIONS OF LIABILITY.**

**CONDITIONS:** The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Makhteshim Agan of North America, Inc. All such risks shall be assumed by the user or buyer.

**DISCLAIMER OF WARRANTIES:** To the extent consistent with applicable law, Makhteshim Agan of North America, Inc. makes no other warranties, express or implied, of merchantability or of fitness for a particular purpose or otherwise, that extend beyond the statements made on this label. No agent of Makhteshim Agan of North America, Inc. is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. To the extent consistent with applicable law, Makhteshim Agan of North America, Inc. disclaims any liability whatsoever for special, incidental or consequential damages resulting from the use or handling of this product.

**LIMITATIONS OF LIABILITY:** To the extent consistent with applicable law, the exclusive remedy of the user or buyer for any and all losses, injuries or damages resulting from the use or handling of this product, whether in contract, warranty, tort, negligence, strict liability or otherwise, shall not exceed the purchase price paid or at Makhteshim Agan of North America, Inc.'s election, the replacement of product.

MANA Alias is a registered trademark of Celsius Property B.V.

MANA Alias 4F(66222-156)(EPA Approved 08-13-09)(Notif to EPA 08-27-10)(Amend to EPA 02-04-11)