



U.S. ENVIRONMENTAL PROTECTION AGENCY
 Office of Pesticide Programs
 Registration Division (7505C)
 1200 Pennsylvania Ave., N.W.
 Washington, D.C. 20460

EPA Reg. Number:

66222-173

Date of Issuance:

MAY 14 2008

NOTICE OF PESTICIDE:

Registration
 Reregistration

(under FIFRA, as amended)

Term of Issuance:

Conditional

Name of Pesticide Product:

Bromacil Plus Diuron Herbicide

Name and Address of Registrant (include ZIP Code):

Makheshim Agan of North America, Inc
 4515 Falls of Neuse Road, Suite 300
 Raleigh, NC 27609

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA section 3(c)(7)(A) provided that you:

1. Submit the results of the one-year storage stability (830.6317) and corrosion characteristics (830.6320) studies when they are available.
2. Submit and/or cite all data required for registration/reregistration of your product when the Agency requires all registrants of similar products to submit such data.
3. Make the labeling changes listed below before you release the product for shipment:

Signature of Approving Official:

James A. Tompkins, Product Manager (25)
 Herbicide Branch, Registration Division (7505P)

Date:

MAY 14 2008

- a. Add the phrase "EPA Registration No. 66222-173"
- b. On page 2, under Use Precautions and Restrictions, revise item 2a to read similar to "Do not spray except as **instructed** by this label." The Agency no longer allows use of the word "recommended" in this context.
- c. On pages 5 and 6 of the label under citrus item g must be revised to read similar to "Do not apply at less than 60-day intervals when making multiple applications to trees less than 4 years old or 80-day intervals to trees 4 years old and older. A maximum of 2 applications of product per year is permitted.
- d. On page 7, under Non-Agricultural Uses, Application Information, add statements stating the following:
 - 1. Apply a maximum of two applications per year.
 - 2. The minimum retreatment interval is 90 days.
 - 3. A maximum of 12 pounds active ingredient bromacil per year is allowed.
 - 4. A maximum of 12 pounds active diuron is allowed per year in areas of high rainfall or dense vegetation. A maximum of 8 pounds active diuron is allowed in all other areas.
- 4. Submit one (1) copy of your final printed label before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6 (e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Enclosure

ACCEPTED
with COMMENTS
In EPA Letter Dated:

MAY 14 2008

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under EPA Reg. No.
66222-173

BROMACIL PLUS DIURON

Herbicide
Dispersible granules

Contains bromacil and diuron, the active ingredients used in Dupont Krovar 1 DF
Bromacil plus Diuron is not manufactured or distributed by E.I. du Pont de Nemours and Company

ACTIVE INGREDIENT:

Bromacil: [5-bromo-3-sec-butyl-6-methyluracil]	40.0%
Diuron: [3-(3,4-dichlorophenyl)-1,1-dimethylurea].....	40.0%
INERT INGREDIENTS:	<u>20.0%</u>

TOTAL 100.0%

EPA Reg. No. 66222-xx

EPA Est. No.

Manufactured for:
Makhteshim Agan of North America, Inc.
4515 Falls of Neuse Rd., Suite 300
Raleigh, NC 27609

NET WEIGHT: POUNDS

KEEP OUT OF REACH OF CHILDREN CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, find someone to explain it to you in detail.)

FIRST AID

IF ON SKIN OR CLOTHING:

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes.
- Call a poison control center or doctor for treatment advice.

IF IN EYES:

- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.

IF INHALED:

- Call a poison control center or doctor for treatment advice.
- Move person to fresh air.
- If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.

IF SWALLOWED:

- Call a poison control center or doctor for further treatment advice.
- Call a poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to do so by a poison control center or doctor.
- Do not give anything by mouth to an unconscious person.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact PROSAR at 1-877-250-9291 for emergency medical treatment information.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS & DOMESTIC ANIMALS CAUTION

Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for Category A on an EPA chemical-resistance category selection chart.

Pilots, flaggers and groundbloom applicators must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks

Mixers, loaders, other applicators, and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks
- Chemical-resistant gloves made of any waterproof material such as polyethylene or polyvinylchloride.
- A NIOSH approved dust/mist filtering respirator with any N, R, P, or HE filter or with approval number prefix TC-21C.
- Chemical-resistant apron when mixing, loading, or cleaning equipment or spills.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROL STATEMENT

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

Pilots must use an enclosed cockpit that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(6)].

Flaggers supporting aerial applications must use an enclosed cab that meets the definition in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(5) for dermal protection.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

For terrestrial uses, do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters. Bromacil is known to leach through soil and has been found in ground water as a result of normal field use. Users are advised not to apply in areas where soils are permeable, particularly where ground water is used for drinking water. Consult with the pesticide state lead agency for information regarding soil permeability and aquifer vulnerability in your area.

GENERAL INFORMATION

Bromacil Plus Diuron is formulated as a dispersible granule to be mixed in water. It is to be sprayed in citrus for selective weed control and on non-crop areas for non-selective control of weeds. Bromacil Plus Diuron controls many annual weeds at the lower label rates and certain perennial weeds at the highest label rates.

Moisture is necessary to move Bromacil Plus Diuron into the root zone of susceptible weeds and brush. Application to moist soil or to soil where moisture from rainfall or sprinkler irrigation occurs within 14 days of application will provide the best control. Symptoms of control will be slow to appear because the product must first move into the root zone of the susceptible plants. The level and length of weed and brush control will be dependent upon the amount of herbicide applied, soil texture, amount of moisture applied and other soil and water management practices.

USE PRECAUTIONS AND RESTRICTIONS

If you plan to use Bromacil Plus Diuron for selective weed control in citrus or for non-selective weed control in non-crop areas, adhere to the following:

1. Do not use in any recreational areas or around homes.
2. To prevent injury or loss to desirable plants and trees:
 - a. Do not apply except as recommended by this label.

- b. Thoroughly clean all traces of Bromacil Plus Diuron from application equipment – tank, pump, hoses, etc.- immediately after use by thoroughly washing/rinsing with several changes of water after removing nozzle tips and screens (clean these parts separately).
 - c. Do not drain or clean equipment near desirable trees or other plants, or onto areas where roots of desirable trees and plants may extend, or onto areas where Bromacil Plus Diuron may be washed or moved into contact with roots of desirable trees and plants.
 - d. Do not use on lawns, walks, driveways, tennis courts, or similar areas.
 - e. Prevent drift of dry powder or spray to desirable plants.
 - f. Do not store and handle Bromacil Plus Diuron around fertilizers, insecticides, fungicides, or seeds.
 - g. Do not use in home fruit planting, or in citrus orchards interplanted to other trees or desirable plants.
3. Do not apply Bromacil Plus Diuron through any type of irrigation system.
 4. Do not graze cattle in treated areas.
 5. Only citrus may be planted into treated areas and only after one year of the last application.
 6. Other crops may not be planted until 2 years after the last application.

MIXING

Sprayers should be properly calibrated with clean water only and checked regularly during operation. It is important to accurately measure Bromacil Plus Diuron and to only mix enough spray mixture for the job at hand. The spray tank should not be over-filled. Excess spray mixture should be diluted and applied at labeled rates/uses. It should never be discharged at a single location in the field/grove or at the mixing/loading location. Do not store or mix near well site.

CROP ROTATION BIOASSAY

In sites where Bromacil Plus Diuron has been used, a field bioassay should be completed prior to planting any desired crop. In arid climates (10 inches of rainfall or less) or areas where drought conditions have prevailed for one or more years, a field bioassay should be completed prior to planting any desired crop. A successful field bioassay means growing to maturity a test strip of the crop(s) intended for production. The test strip should cross the entire field including knolls and low areas. The results from the bioassay may require the two-year crop rotation interval to be extended.

SPRAY PREPARATION

Mixing in water: Fill tank 1/2 full with water. Start agitation system, add Bromacil Plus Diuron and continue adding water. Add separately each additional component of any tank-mix while adding water. Continue agitation throughout.

Mixing in liquid fertilizer: A fertilizer solution may be used in the spray mixture. Small quantities should be tested for compatibility by the following procedures before full scale mixing:

1. Put 1 pint fertilizer solution in a quart jar.
2. Mix 2 teaspoons Bromacil Plus Diuron with 2 tablespoons of water: mix thoroughly and add to fertilizer solution.
3. Close jar and shake well.
4. If other herbicides are used in the mixture, premix 2 teaspoonfuls of dry materials or 1 teaspoonful of liquids with 2 tablespoons of water; add to Bromacil Plus Diuron fertilizer solution mixture.
5. Close jar and shake well.
6. Watch mixture for several seconds; check again in 30 minutes.
7. If mixture does not separate, foam, gel, or become lumpy, it may be used.

Provided the above procedure shows the mixture to be compatible, prepare the tank mixture as follows: add the fertilizer solution to the spray tank first; with agitator running, add the required amount of Bromacil Plus Diuron and mix thoroughly.

Mixing with other herbicides: Determine the tank mixture partner(s) compatibility with Bromacil Plus Diuron by following the directions above. For Step 1 above, use 1 pint of water instead of the liquid fertilizer. Provided the above procedure shows the mixture to be compatible, Bromacil Plus Diuron may be used in this tank mixture.

SPRAY TANK CLEAN OUT

Thoroughly clean all traces of Bromacil Plus Diuron, from application equipment immediately after use. Flush the tank, pump, hoses, and boom with several changes of water after removing nozzle tips and screens (clean these parts separately). Dispose of the equipment wash water by applying it to a use-site listed on this label.

RESISTANCE

When herbicides that affect the same biological site of action are used repeatedly over several years to control the same weed species in the same field, naturally-occurring resistant biotypes may survive a correctly applied herbicide treatment, propagate, and become dominant in that field. Adequate control of these resistant weed

biotypes cannot be expected. If weed control is unsatisfactory, it may be necessary to retreat the problem area using a product affecting a different site of action.

To better manage herbicide resistance through delaying the proliferation and possible dominance of herbicide resistant weed biotypes, it may be necessary to change cultural practices within and between crop seasons such as using a combination of tillage, retreatment, tank-mix partners and/or sequential herbicide applications that have a different site of action. Weed escapes that are allowed to go to seed will promote the spread of resistant biotypes.

It is advisable to keep accurate records of pesticides applied to individual fields to help obtain information on the spread and dispersal of resistant biotypes. Consult your agricultural dealer, consultant, applicator, and/or appropriate state agricultural extension service representative for specific alternative cultural practices or herbicide recommendations available in your area.

INTEGRATED PEST MANAGEMENT

This product may be used as part of an Integrated Pest Management (IPM) program that can include biological, cultural, and genetic practices aimed at preventing economic pest damage. IPM principles and practices include field scouting or other detection methods, correct target pest identification, population monitoring, and treating when target pest populations reach locally determined action thresholds. Consult your state cooperative extension service, professional consultants, or other qualified authorities to determine appropriate action treatment threshold levels for treating specific pest/crop systems in your area.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Bromacil Plus Diuron herbicide should only be used in accordance with recommendations on this label. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Use of this product in certain portions of California, Oregon, and Washington is subject to the January 22, 2004 Order for injunctive relief in Washington Toxics Coalition et al vs EPA. Col-132C (W.D.W.A.). For information, please refer to www.epa.gov/espp/wtc/.

AGRICULTURAL USES

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves made of any waterproof material
- Shoes plus socks

APPLICATION INFORMATION

Bromacil Plus Diuron should be applied as a band or broadcast treatment with a calibrated fixed-boom power sprayer with nozzles equipped with 50 mesh or larger screens. Shut off spray booms when starting, turning, slowing or stopping because higher than labeled use rates may be applied and injury to the crop or successive crops could occur. Use rates provided below for Bromacil Plus Diuron are expressed for broadcast treatments. If band treatments are applied, then proportionately less Bromacil Plus Diuron should be used. A minimum of 10 gallons spray volume per acre should be used to insure proper suspension of the product in the spray tank. Sufficient spray volume should be used to allow for uniform coverage of the sprayed area. Continuous agitation in the spray tank with mechanical or hydraulic means (do not use air agitation) is necessary to keep Bromacil Plus Diuron in suspension. If by-pass or return line agitation is used, it should terminate at the bottom of the tank to decrease foaming. When using Bromacil Plus Diuron alone or in tank mixture thoroughly re-agitate if allowed to settle.

Applications should be made at any time of the year provided rainfall or overhead irrigation is available to activate the herbicide, preferably just before or just after weeds have germinated. Best results are obtained when

Bromacil Plus Diuron is applied to bare ground. If weeds are already present, a tank-mixture of Bromacil Plus Diuron with a foliar active herbicide is recommended (See Tank Mixtures section of the label). Control dense populations of hard-to-kills weeds with other herbicides before making an application of Bromacil Plus Diuron.

WEEDS CONTROLLED (annuals)

Barnyardgrass	<i>Echinochloa crus-gali</i>
Brome, downy (cheatgrass)	<i>Bromus tectorum</i>
Chickweed, common	<i>Stellaria media</i>
Chickweed, mouseear	<i>Cerastium vulgatum</i>
Clovers, (annual)	<i>Trifolium spp.</i>
Filaree	<i>Erodium spp.</i>
Fleabane, flaxleaved (hairy)	<i>Conyza bonariensis</i>
Foxtail	<i>Setaria spp.</i>
Goatweed	<i>Scoparia dulcis</i>
Groundsel	<i>Senecio spp.</i>
Horseweed (marestail)	<i>Conyza canadensis</i>
Johnsongrass	<i>Sorghum halepense</i>
Junglerice	<i>Echinochloa colona</i>
Kochia	<i>Kochia scoparia</i>
Lambsquarter	<i>Chenopodium album</i>
Lettuce, wild	<i>Lactuca serriola</i>
Mustard, wild	<i>Brassica kaber</i>
Natalgrass (red top)	<i>Rhynchelytrum repens</i>
Nightshade (annual)	<i>Solanum spp.</i>
Pigweed	<i>Amaranthus spp.</i>
Pineappleweed	<i>Matricaria matricariodes</i>
Puncturevine, common	<i>Tribulus terrestris</i>
Purslane, common	<i>Portulaca oleracea</i>
Pusley, Florida	<i>Richardia scraba</i>
Ragweed, common	<i>Ambrosia artmisifolia</i>
Sandbur (sandspur)	<i>Cenchrus spp.</i>
Shepherdspurse	<i>Capsella bursa-pastoris</i>
Sowthistle, annual	<i>Sonchus oleraceus</i>
Spanishneedles	<i>Bidens pilosa</i>
Thistle, Russian	<i>Salsola australis</i>

WEEDS CONTROLLED (perennials) (at maximum rates and repeat treatments)

Balsamapple vine (seedling)	<i>Momordica charantia</i>
Bermudagrass	<i>Cynodon dactylon</i>
Drymary	<i>Drymaria spp.</i>
Guineagrass	<i>Panicum maximum</i>
Milkweed vine (strangler)	<i>Morrenia odorata</i>
Quackgrass	<i>Agropyron repens</i>
Vines (seedlings)	

NOTE: Best control of perennial weeds is obtained when the highest rates of Bromacil Plus Diuron is applied. Repeat applications are usually necessary to control perennials. To improve control of perennials, cultivate prior to treatment. After treatment with Bromacil Plus Diuron, avoid cultivation as long as weed control is holding. Working the soil after treatment could result in reduced weed control.

CITRUS

Bromacil Plus Diuron may be applied in citrus for control of the annual and perennial weeds listed above. Applications should be made at any time of the year provided rainfall or overhead irrigation is available to activate the herbicide, preferably just before or just after weeds have germinated. The degree and duration of control will vary with the amount of herbicide applied, soil texture, rainfall and other conditions. It is non-corrosive to equipment, non-flammable and non-volatile.

Apply Bromacil Plus Diuron as a band or broadcast treatment beneath and/or between trees. Avoid contact of foliage and fruit with spray or mist. Avoid overlapping and shut off spray boom while starting, turning, or stopping as injury to trees may results. Temporary yellowing of citrus leaves may occur following treatment.

- a. Do not use on soils with less than 1% organic matter.
- b. Do not use on poorly drained soils, gravelly soils or thinly covered or exposed subsoils.
- c. Do not treat trees planted in irrigation furrows.
- d. Do not treat diseased or stressed citrus trees.
- e. Do not use in citrus groves inter-planted with other desirable trees or plants or in areas where roots of desirable trees or plants may extend as injury to desirable trees or plants may result.
- f. Do not use in home citrus plantings.
- g. Do not apply at less than 60 day intervals when making multiple applications.
- h. Do not graze cattle in treated areas.

All traces of Bromacil Plus Diuron must be removed from the application equipment immediately after use. Remove nozzles and screens and clean separately. Several changes of water should be run through the application equipment.

CALIFORNIA, ARIZONA

Trees Established for at least Three Years:

Apply Bromacil Plus Diuron in late fall or early winter before winter annuals become established. Wait to apply treatment until after the soil has been settled by fall or early winter rains. Apply as needed, 4 to 5 pounds Bromacil Plus Diuron per acre on course soils (1-2% organic matter) and 5-6 pounds Bromacil Plus Diuron per acre on fine soils (2.5% or greater organic matter). Alternatively, apply 3-4 pounds per acre in the fall and 2-4 pounds in the spring. Do not exceed 6 pounds per acre per year. If problem weeds like groundsel or pucturevine are present at the location, use the highest rates allowed on this label according to soil type. These rates will also suppress low levels of bermudagrass and yellow nutsedge. Best results will occur if this treatment is applied annually.

FLORIDA

Apply Bromacil Plus Diuron as a band treatment only in Florida citrus groves. Do not apply trunk to trunk. The use of Bromacil Plus Diuron is prohibited for weed control in non-bedded citrus groves located on any permeable, better drained soil identified in the intended site of application. Permeable, better drained soils which occur in citrus producing areas of the state include soils unnamed and characteristic of quartzipsamments, and the following soil series classifications:

Adamsville	Archbold	Astatula
Bahiahonda	Broward	Canaveral
Candler	Cocoa	Dade
Florahome	Fort Meade	Gainseville
Lake	Lakewood	Neilhurst
Orlando	Orsino	Palm Beach
Paola	Satellite	St. Augustine
St. Lucie	Tavares	

APPLICATION INSTRUCTIONS

Since all use rates for Bromacil Plus Diuron are expressed for broadcast treatments, use proportionately less per acre for band treatments in Florida. Multiple applications may be required for control of some problem weeds.

Note: Do not apply more than 16 pounds per acre of Bromacil Plus Diuron per year. This corresponds to 6.4 pounds per treated acre of bromacil and 6.4 pounds per treated acre of diuron. These amounts represent the maximum allowable use rates for bromacil and diuron on citrus inclusive of all formulations of bromacil or diuron that might be used.

Trees Established Less Than One Year: For control of annual weeds, apply 2-4 pounds of Bromacil Plus Diuron per treated acre as needed to maintain weed control. Do not apply more than 6 pounds per treated acre during the any 6 month period nor more than 8 pounds per treated acre during the 1st year.

Trees Established One to Three Years: For control of annual weeds, apply 2-4 pounds of Bromacil Plus Diuron per treated acre. A second application may be made when needed to maintain weed control, but do not exceed 8 pounds per treated acre per year.

Trees Established Three or More Years: Apply 4-8 pounds per treated acre as needed to maintain weed control. Do not apply more than 16 pounds of Bromacil Plus Diuron per treated acre per year.

LOUISIANA

Trees Established for at least Three Years: On course soils, apply Bromacil Plus Diuron at 2 to 4 pounds per acre whereas on finer soils (2.5% or greater organic matter) apply Bromacil Plus Diuron at 4 to 6 pounds per acre. Use the highest labeled rates for suppression of perennials. Alternatively, make two applications per year

for weed control using 2 pounds per acre on coarse soils and 3 pounds per acre of finer soils. In either case, do not apply more than 6 pounds Bromacil Plus Diuron per treated acre per year.

TEXAS

Trees Established Less Than One Year: Apply 2-4 pounds of Bromacil plus diuron per acre as needed to maintain weed control. Do not apply at less than 60 day intervals. Do not apply more than 6 pounds per acre per year.

Trees Established One or Two Years: Apply 2-4 pounds of Bromacil plus diuron per acre. A second application may be made when needed to maintain weed control, but do not exceed 6 pounds per acre per year.

Trees Established Three or More Years: Make one to two applications per year as needed to maintain weed control. Use 2-4 pounds per acre on coarser soils (sands, loamy sands, sandy loams) and 4-6 pounds per acre on finer soils (silt loams, clay loams, or soils with organic matter of 2½% or more). Use the higher rate for maximum suppression of perennials. Do not use more than 6 pounds per acre per year.

TANK MIXTURES

To achieve a broader spectrum of weed control in citrus, Bromacil Plus Diuron may be tank mixed with other herbicides registered for use in citrus. Be sure to follow the use information and restrictions on the labels of the tank-mix herbicides. The label guidelines that are the most restrictive must be followed. Adjuvants used with herbicides in citrus may also be tank-mixed with Bromacil Plus Diuron.

Note: If the user has no experience with a particular tank-mixture, a jar-compatibility test should be performed to determine chemical compatibility of the products before they are mixed in the spray tank. See the SPRAY PREPARATION section of this label above for additional details.

NON-AGRICULTURAL USES

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Non-crop weed control is not within the scope of the Worker Protection Standard.

Do not enter or allow others to enter the treated area until sprays have dried.

USE RESTRICTIONS – STATE OF FLORIDA

In the state of Florida, the use of Bromacil Plus Diuron is prohibited in the counties of Hardee, Highland, Polk, Orange, and Lake. For Non-Agricultural Usage in all other areas of the state, do not apply more than 16 pounds per acre per year of Bromacil Plus Diuron. This corresponds to 6.4 pounds per treated acre of bromacil and 6.4 pounds per treated acre of diuron. These amounts represent the maximum allowable use rates for bromacil and diuron inclusive of all formulations of bromacil or diuron that might be used.

APPLICATION INFORMATION

For general weed control in uncultivated non-agricultural areas (for example, airports, highway, railroad and utility rights-of-way, sewage disposal areas), uncultivated non-crop producing areas (for example, farmyards, fuel storage areas, fence rows, barrier strips) and outdoor industrial sites (for example, lumberyards, pipeline and tank farms) apply Bromacil Plus Diuron using a properly calibrated fixed-boom power sprayer. Use rates provided below for Bromacil Plus Diuron are expressed for broadcast treatments. If band treatments are applied, then proportionately less Bromacil Plus Diuron should be used. A minimum of 10 gallons spray volume per acre should be used to insure proper suspension of the product in the spray tank. Sufficient spray volume should be used to allow for uniform coverage of the sprayed area. Continuous agitation in the spray tank with mechanical or hydraulic means (do not use air agitation) is necessary to keep Bromacil Plus Diuron in suspension. If by-pass or return line agitation is used, it should terminate at the bottom of the tank to decrease foaming. When using Bromacil Plus Diuron alone or in tank mixture thoroughly re-agitate if allowed to settle.

A broader spectrum of weeds may be controlled by tank-mixing Bromacil Plus Diuron with other herbicides. Also, higher rates of Bromacil Plus Diuron applied with residual herbicides will provide total vegetation control. Always add a suitable surfactant at 0.25% v/v when attempting to control emerged weeds.

Note: Applications may also be made using a handgun sprayer. Use a spray volume of at least 40 gallons per acre to insure uniform coverage. For small areas, a hand sprayer or sprinkling may be used.

NON-CROP WEED CONTROL

APPLICATION RATES AND TIMING

Bromacil Plus Diuron should be applied at the rates provided in the tables below according to weed type. Lower rates of Bromacil Plus Diuron provide short-term control of the weeds listed whereas higher rates provide more residual control. Always apply as a preemergence spray when weeds are actively germinating or growing. Since

moisture is required to move the herbicide into the root zone of the weeds, best preemergence control is obtained when Bromacil Plus Diuron is applied prior to a rainfall and weed germination. In areas of low and infrequent rainfall, such as the Western U.S., Bromacil Plus Diuron should be applied well before the Fall freeze or immediately after the Spring thaw to increase the chances of having adequate moisture to activate and disperse the herbicide in the soil. Do not treat frozen or saturated soils, or soils that are non-receptive to percolation.

Do not apply to sites which have roots of desirable plants growing into the treatment zone as plant injury or death may occur. Do not apply to hard or impervious soils, water saturated soils or to any surface that does not allow the herbicide to be moved into the soil horizon with moisture. Unusually heavy rainfall shortly after application may move the product off-target to the lowest surrounding point and cause plant injury or death.

If herbicide treated soil is disturbed by any physical or mechanical means, the herbicide barrier is disrupted and the likelihood of non-performance may increase. For best performance results, make sure the treatment area is stable after the application for the desired weed control period.

WEEDS CONTROLLED

Bromacil plus diuron effectively controls the following broadleaf weeds and grasses when applied at the rates shown.

Broadleaf Weeds: 6-8 pounds per acre

Clovers (annual)	<i>Trifolium spp.</i>
Fiddleneck	<i>Amsinckia intermedia</i>
Filaree	<i>Enulium spp.</i>
Knapweed, diffuse	<i>Centaurea diffusa</i>
Lambsquarter, common	<i>Chenopodium album</i>
Lettuce, prickly	<i>Lactuca serriola</i>
Mustards	<i>Brassica, spp.</i>
Pigweed	<i>Amaranthus spp.</i>
Ragweed	<i>Ambrosia spp.</i>
Sunflower, common	<i>Helianthus annuus</i>
Thistle, Russian	<i>Salsola iberica</i>

Broadleaf Weeds: 8-12 pounds per acre

Carrot, wild	<i>Daucus carota</i>
Dandelion, common	<i>Taraxacum officinale</i>
Dock, curly	<i>Rumex crispus</i>
Knapweed, spotted	<i>Centaurea maculosa</i>
Knotweed, prostrate	<i>Polygonum aviculare</i>
Kochia	<i>Kochia scoparia</i>
Marestail, common (horseweed)	<i>Conyza Canadensis</i>
Parsnip, wild	<i>Pastinaca sativa</i>
Plantain	<i>Plantago spp.</i>
Puncturevine	<i>Tribulus terrestris</i>
Spurge	<i>Euphorbia spp.</i>
Thistle, milk	<i>Silybum marianum</i>
Yarrow, common	<i>Achillea millefolium</i>

Broadleaf Weeds: 12-16 pounds per acre

Cinquefoil, common	<i>Potentilla Canadensis</i>
Goldenrod	<i>Solidago spp.</i>
Milkweed, common	<i>Asclepias syriaca</i>

Grasses: 6- 8 pounds per acre

Barley, foxtail	<i>Hordeum jubatum</i>
Brome	<i>Bromus spp.</i>
Cheat	<i>Bromus secalinus</i>
Cupgrass, Prairie	<i>Eriochloa contracta</i>
Foxtail	<i>Setaria spp.</i>
Oat, wild	<i>Avena fatua</i>
Ryegrass, Italian	<i>Lolium multiflorum</i>

Quackgrass	<i>Agropyron repens</i>
Wheatgrass, intermediate	<i>Agropyron intermedium</i>

Grasses: 8-12 pounds per acre

Bahiagrass	<i>Paspatum notatum</i>
Crabgrass	<i>Digitaria spp.</i>
Goosegrass	<i>Eleusine indica</i>
Rye	<i>Secale cereale</i>
Vaseygrass	<i>Paspalum urvillei</i>

Grasses: 12-16 pounds per acre

Bluegrass	<i>Poa spp.</i>
Dropseed, sand*	<i>Sporobolus cryptandrus</i>
Fescue	<i>Festuca spp.</i>
Saltgrass*	<i>Distichlis spp.</i>

*Note: Best control of Saltgrass and Sand Dropseed is achieved from a Spring application prior to plant green-up. For control of hard-to-kill perennials such as bermudagrass (*Cynodon dactylon*), bouncingbet (*Saporaria officinalis*), dogbane (*Apocynum spp.*), Johnsongrass (*Sorghum halepense*), and nutsedge (*Cyperus spp.*), apply 19-30 pounds per acre (except Florida).

For extended control of annual weeds and partial control of perennials such as bermudagrass and nutsedge apply 10-18 pounds* per acre. Use the higher Bromacil plus diuron rates on adsorptive soils (high in organic matter or carbon). Best results occur when application is made just before weed emergence or in the early stages of weed growth.

Retreating: Apply 4 to 6 pounds per acre when annual weeds and grasses reappear on sites where weed growth has been controlled.

Small areas: ¼ cupful of Bromacil plus diuron per 200 sq. ft. is approximately 15 pounds per acre.

TANK MIXTURES

To achieve a broader spectrum of weed control in non-crop areas, Bromacil Plus Diuron may be tank mixed with other herbicides registered for such use. Be sure to follow the use information and restrictions on the labels of the tank-mix herbicides. The label guidelines that are the most restrictive must be followed. Adjuvants used with herbicides in non-crop areas may also be tank-mixed with Bromacil Plus Diuron.

Note: If the user has no experience with a particular tank-mixture, a jar-compatibility test should be performed to determine chemical compatibility of the products before they are mixed in the spray tank. See the SPRAY PREPARATION section of this label above for additional details.

Be sure to thoroughly re-agitate the contents of the spray tank if allowed to settle out.

**SPECIAL USES
UNDER ASPHALT AND CONCRETE PAVEMENT**

APPLICATON INFORMATION

Bromacil plus diuron can be used to control weeds under asphalt and concrete pavement, such as that used in parking lots, highway shoulders, median strips, roadways, and other industrial sites.

Bromacil plus Diuron should only be used in an area that had been prepared according to good construction practices. Use sufficient water to insure uniform coverage, generally 100 gal. per acre. Agitate the tank continuously to keep Bromacil plus diuron in suspension.

APPLICATION TIMING AND RATES

Apply Bromacil Plus Diuron at 17 to 30 pounds per acre immediately before paving to avoid lateral movement of the herbicide as a result of soil movement due to rainfall or mechanical means. Use higher rates for control of problem weeds and for more residual control.

TANK MIXTURES

To control a broader spectrum of weeds, or for an extended period of weed control, a tank mixture of Bromacil Plus Diuron at 7 to 15 pounds per acre plus Dupont OUST®XP at 4 to 8 ounces per acre may be used.

IMPORTANT PRECAUTIONS: UNDER ASPHALT ONLY

- Do not use Bromacil Plus Diuron under pavement in residential properties such as driveways, or in recreational areas, including jogging or bike paths, tennis courts, or golf cart paths.
- Desirable plants may be injured if their roots extend into treated areas or if planted in treated areas.

SPRAY DRIFT MANAGEMENT

The interaction of many equipment and weather-related factors determines the potential for spray drift. The applicator is responsible for considering all these factors when making application decisions.

AVOIDING SPRAY DRIFT IS THE RESPONSIBILITY OF THE APPLICATOR.

IMPORTANCE OF DROPLET SIZE

The most effective way to reduce drift potential is to apply large droplets (>150-200 microns). The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. The presence of sensitive species nearby, the environmental conditions, and pest pressure may affect how an applicator balances drift control and coverage. **APPLYING LARGER DROPLETS REDUCES DRIFT POTENTIAL, BUT WILL NOT PREVENT DRIFT IF APPLICATIONS ARE MADE IMPROPERLY OR UNDER UNFAVORABLE ENVIRONMENTAL CONDITIONS! See Wind, Temperature and Humidity, and Surface Temperature Inversions sections of this label.**

Controlling Droplet Size – General Techniques

- **Volume** - Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.
- **Pressure** – Use the lower spray pressures recommended for the nozzle. Higher pressure reduces droplet size and does not improve canopy penetration. **WHEN HIGHER FLOW RATES ARE NEEDED, USE A HIGHER-CAPACITY NOZZLE INSTEAD OF INCREASING PRESSURE.**
- **Nozzle Type**- Use a nozzle type that is designed for the intended application. With most nozzle types narrower spray angles produce larger droplets. Consider using low-drift nozzles.

Controlling Droplet Size – Aircraft

- **Number of Nozzles** – Use the minimum number of nozzles with the highest flow rate that provide uniform coverage.
- **Nozzle Orientation** – Orienting nozzles so that the spray is emitted backwards, parallel to the airstream will produce larger droplets than other orientations.
- **Nozzle Type** – Solid stream nozzles (such as disc and core with swirl plate removed) oriented straight back produce larger droplets than other nozzle types.

BOOM LENGTH AND HEIGHT

- **Boom Length (aircraft)**- The boom length should not exceed ¾ of the wing length, using shorter booms decreases drift potential. For helicopters use a boom length and position that prevents droplets from entering the rotor vortices.
- **Boom Height (aircraft)**- Application more than 10 ft. above the canopy increases the potential for spray drift.
- **Boom Height (ground)**- Setting the boom at the lowest height which provides uniform coverage reduces the exposure of droplets to evaporation and wind. The boom should remain level with the crop and have minimal bounce.

WIND

Drift potential increases at wind speeds of less than 3 mph (due to variable direction and inversion potential) or more than 10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given wind speed. **AVOID APPLICATIONS DURING GUSTY OR WINDLESS CONDITIONS.**

NOTE: Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they effect spray drift.

TEMPERATURE AND HUMIDITY

When making applications in hot and dry conditions, set up equipment to produce larger droplets to reduce effects of evaporation.

SURFACE TEMPERATURE INVERSIONS

Drift potential is high during a surface temperature inversion. Surface inversions restrict vertical air mixing, which causes small suspended droplets to remain close to the ground and move laterally in a concentrated cloud. Surface inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates a surface inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

SHIELDED SPRAYERS

Shielding the boom or individual nozzles can reduce the effects of wind. However, it is the responsibility of the applicator to verify that the shields are preventing drift and not interfering with uniform deposition of the product.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store product in original container only. Store in a cool, dry place.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed, by State and local authorities, by burning. If burned, stay out of smoke.

LIMITATION OF WARRANTY AND LIABILITY

Read the entire Directions for Use, Conditions of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Makhteshim Agan of North America, Inc. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: To the extent consistent with applicable law, Makhteshim Agan of North America, Inc. makes no other warranties, express or implied, of merchantability or of fitness for a particular purpose or otherwise, that extend beyond the statements made on this label. No agent of Makhteshim Agan of North America, Inc. is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. To the extent consistent with applicable law, Makhteshim Agan of North America, Inc. disclaims any liability whatsoever for special, incidental or consequential damages resulting from the use or handling of this product.

LIMITATIONS OF LIABILITY: To the extent consistent with applicable law, the exclusive remedy of the user or buyer for any and all losses, injuries or damages resulting from the use or handling of this product, whether in contract, warranty, tort, negligence, strict liability or otherwise, shall not exceed the purchase price paid or at Makhteshim Agan of North America, Inc.'s election, the replacement of product.