

66222-185

09-04-2009

1/30



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

Ms, Anne Stout
Product Registration
Maktheshim Agan of North America
MANA, Inc.
4515 Falls of Neuse Rd, Suite 300
Raleigh, NC 27609

SEP 4 2009

SUBJECT: Application for Pesticide Notification (PRN 98-10)
Request Alternate Marketing Language (Comply w/ State of Arizona)
EPA Reg. No. 66222-185
Application Dated July 1, 2009

Dear Registrant:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 dated 07/01/09 for the above product. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the action(s) requested fall within the scope of PRN 98-10. The label submitted with the application has been stamped "Notification" and will be placed in our records.

If you have any questions, please call me directly at 703-305-6249 or Owen F. Beeder of my staff at 703-308-8899.

Sincerely,

Linda Arrington
Notifications & Minor Formulations Team Leader
Registration Division (7505P)
Office of Pesticide Programs

Please read instructions on reverse before completing form.

Form Approved. OMB No. 2070-0080. Approval expires 2-28-95



United States
Environmental Protection Agency
Washington, DC 20460

 Registration
 Amendment
 Other

OPP Identifier Number

Application for Pesticide - Section I

1. Company/Product Number 66222-185	2. EPA Product Manager Linda Arrington	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Quali-Pro Myclobutanil 20EW T&O	PM# Registration Support Branch	
5. Name and Address of Applicant (Include ZIP Code) Makhteshim Agan of North America, Inc. 4515 Falls of Neuse Rd, Ste 300 Raleigh NC 27609 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

Section - II

<input type="checkbox"/> Amendment - Explain below.	<input checked="" type="checkbox"/> Final printed labels in response to Agency letter dated _____	NOTIFICATION SEP - 4 2009
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.	
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.	

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

Notification of Alternate Marketing Language (state-specific statements):
This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula for this product. I understand that it is a violation of 18 USC Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under Sections 12 and 14 of FIFRA.

Section - III

1. Material This Product Will Be Packaged In:

Child-Resistant Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	2. Type of Container <input type="checkbox"/> Metal <input type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____
* Certification must be submitted		If "Yes" Unit Packaging wgt. No. per container	If "Yes" Package wgt. No. per container

3. Location of Net Contents Information
 Label Container

4. Size(s) Retail Container

5. Location of Label Directions

6. Manner in Which Label is Affixed to Product
 Lithograph
 Paper glued
 Stenciled Other _____

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)

Name Anne Stout	Title Registration Specialist	Telephone No. (Include Area Code) 901-861-4400
--------------------	----------------------------------	---

Certification
I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.

2. Signature 	3. Title Registration Specialist	6. Date Application Received (Stamped)
4. Typed Name Anne Stout	5. Date 7-1-09	



July 1, 2009

Registration Support Branch
Document Processing Desk (NOTIF)
U.S. Environmental Protection Agency
Registration Division (7504P)
Ariel Rios Building
1200 Pennsylvania Ave, NW
Washington, DC 20460

Re: Quali-Pro Myclobutanil 20EW T&O, EPA Reg. No. 66222-185
Notification of Alternate Marketing Language per PRN 98-10

To Whom it May Concern:

Makhteshim Agan of North America, Inc. is notifying the Agency of its intention to add alternate marketing language to the referenced product as allowed in PRN 98-10. We certify the following:

This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula for this product. I understand that it is a violation of 18 USC Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under Sections 12 and 14 of FIFRA.

In support of this submission, the following documents are attached:

- Application for registration (EPA Form 8570-1)
- One copy of product labeling with additions marked in yellow

Should you have any questions regarding this submission, please contact me at 901-861-4400 or by email at annies@manainc.com.


Sincerely,

Anne Stout
Registration Specialist

Enclosures

NOTIFICATION

SEP - 4 2009



Myclobutanil 20 EW T&O

A systemic, protectant, and curative fungicide for disease control in turfgrass, landscape ornamentals, greenhouse and nursery ornamentals, apples, stonefruit, and grapes

Contains myclobutanil, the active ingredient used in Eagle 20EW. Quali-Pro Myclobutanil 20EW T&O is not manufactured or distributed by Dow AgroSciences LLC.

Active Ingredient:	% By Wt.
Myclobutanil: alpha-butyl-alpha-(chlorophenyl)-1H-1,2,4-triazole-1-propanenitrile	19.7%
Other Ingredients:	80.3%
Total:	100.0%

Contains Petroleum Distillates. Contains 1.67 pounds of active ingredient per gallon.
EPA Reg. No. 68222-185

**SHAKE WELL BEFORE USING
KEEP OUT OF REACH OF CHILDREN
CAUTION**

For additional precautionary, handling, and use statements, see inside of this booklet.

Manufactured for: Makhteshim Agan of North America, Inc.
4515 Falls of Neuse Road, Suite 300
Raleigh, NC 27609

13018

FIRST AID	
IF IN EYES:	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15 – 20 minutes. Remove contact lenses, if present, after the first 5 minutes; then continue rinsing eye. Call a poison control center or doctor for treatment advice.
IF SWALLOWED:	<ul style="list-style-type: none"> Call a poison control center or doctor immediately for treatment advice. Do not give any liquid to the person. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none"> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
<p>Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact PROSAR at 1-877-250-9291 for emergency medical treatment information.</p> <p>NOTE TO PHYSICIAN: Contains petroleum distillate – vomiting may cause aspiration pneumonia.</p>	

EPA 052109/Notif 070109/Rev A

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION

Causes moderate eye irritation. Harmful if swallowed or absorbed through the skin. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made from barrier laminate
- Shoes plus socks

User Safety Requirements:

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove PPE immediately after handling this product.
- Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

PHYSICAL AND CHEMICAL HAZARDS

Flammable. Keep away from heat and open flame.

ENVIRONMENTAL HAZARDS

Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Do not apply when weather conditions favor drift or runoff from areas treated.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated such as plants, soil, or water is:

- Coveralls
- Chemical-resistant gloves made from barrier laminate
- Shoes plus socks

NONAGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are not within the scope of the Worker Protection Standard for agricultural pesticides, 40 CFR Part 170. The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Seed treatments and professional applications to lawn grasses, golf courses, industrial (office park), municipal, and residential lawns are not within the scope of the Worker Protection Standard.

Keep unprotected persons out of treated area until sprays have dried.

GENERAL INFORMATION

SHAKE WELL BEFORE USING.

State Specific Restrictions:

The state of Arizona has not approved this product for use on agricultural sites. Do not use this product on uses considered by the Arizona statutes to be agricultural uses.

Quali-Pro® Myclobutanil 20EW T&O fungicide is a systemic, protectant, and curative fungicide for the control of specific diseases listed on this label in established turfgrass (including but not limited to residential and commercial lawns, ornamental turf, grounds, or lawns around business and office complexes, and golf course fairways, roughs, tee boxes, and greens), landscape ornamentals, greenhouse and nursery ornamentals, apples, stone fruit, and grapes. Optimum disease control is achieved when the fungicide is applied in a regularly scheduled preventative spray program.

Carefully read and follow label directions including recommended use rates and restrictions. For proper application, determine the size of the area to be treated, the recommended label use rate, and the gallonage to be applied to the area. Prepare only the amount of spray solution required to treat the measured area. Careful calibration of spray equipment is recommended prior to use.

MIXING PROCEDURES

Be sure sprayer is clean and not contaminated with other materials prior to use. Fill the spray tank 1/4 to 1/2 full with clean water and start agitation. Be certain the agitation system is working properly and creates a rolling or rippling effect on the liquid surface. With the agitator running, add the required amount of Quali-Pro Myclobutanil 20EW T&O to the tank and finish filling tank. Always add Quali-Pro Myclobutanil 20EW T&O to the spray tank before adding other materials.

Compatibility: Quali-Pro Myclobutanil 20EW T&O is compatible with most commonly used fungicides, insecticides, growth regulators, micronutrients, and spray adjuvants. When preparing tank

mixtures, spray compatibility charts or your State Cooperative Extension Service specialist should be consulted prior to use.

When tank mixing this product with other pesticides, observe the more restrictive label limitations and precautions. No label dosage rates should be exceeded. This product cannot be mixed with any product containing a label prohibition against such mixing.

APPLICATION PROCEDURES

GROUND EQUIPMENT: Application equipment should be properly calibrated and provide uniform spray coverage.

HAND OR PRESSURIZED SPRAYERS: For best results when applying this product on a protectant schedule, ensure thorough coverage of all plant parts.

CHEMIGATION INSTRUCTIONS—SPRINKLER IRRIGATION: Quali-Pro Myclobutanil 20EW T&O must be applied on a regular protectant fungicide schedule, not an irrigation schedule. Apply this product only through solid set or hand-move sprinkler irrigation systems. Do not apply this product through any other type of irrigation system. Crop injury, lack of fungicidal effectiveness, or illegal pesticide residues in the crop can result from nonuniform distribution of treated water. If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers, or other experts.

Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place. Before applying Quali-Pro Myclobutanil 20EW T&O through sprinkler irrigation equipment, the chemigation system must meet the following specifications.

A person knowledgeable of the chemigation system and responsible for its operation or under the supervision of the responsible person shall shut the system down and make necessary adjustments should the need arise.

1. Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.
2. Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone (RPZ) back flow preventer or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the flow outlet end of the pipe fill and

- the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.
3. Systems not connected to a public water supply must contain a functional check valve, vacuum relief valve, and low-pressure drain appropriately located in the irrigation pipeline to prevent water source contamination from back flow.
 4. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
 5. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
 6. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops; or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.
 7. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
 8. Systems must use a metering pump such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
 9. The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from back flow.
 10. Do not apply when wind speed favors drift beyond the area intended for treatment.

Specific Instructions for Solid-Set and Hand Move Irrigation Equipment

1. Determine area covered by sprinkler.
2. Fill injector solution tank with water and adjust flow rate to use the contents over a 10- to 30-minute interval.
3. Determine the amount of Quali-Pro Myclobutanil 20EW T&O required for the area to be treated.
4. Add the required amount of Quali-Pro Myclobutanil 20EW T&O into the same quantity of water used to calibrate the injection equipment.
5. Maintain constant solution tank agitation during the injection period.
6. Operate system at normal pressures recommended by the manufacturer of the injection equipment and used for the time interval established during calibration.
7. Inject Quali-Pro Myclobutanil 20EW T&O at the end of an irrigation cycle or as a separate application to maximize foliar absorption and retention.
8. Stop injection equipment after treatment is completed. Continue to operate the system until the Quali-Pro Myclobutanil 20EW T&O solution has cleared the last sprinkler head.

10
30

**USE DIRECTIONS FOR TURFGRASS
GENERAL INFORMATION**

Quali-Pro Myclobutanil 20EW T&O should be used in conjunction with turf management practices that are optimal for plant health and disease control. Proper identification of the disease-causing organism is the key to selecting the correct fungicide. Diagnostic kits, extension experts, or other identification methods should be used when developing disease control strategies.

In nonresidential turfgrass (including but not limited to commercial lawns, ornamental turf, grounds or lawns around business and office complexes, and golf course fairways, roughs, tee boxes, and greens), optimum disease control is achieved when Quali-Pro Myclobutanil 20EW T&O is applied in a preventative disease control program at a rate of 1.0 to 2.4 fluid ounces per 1000 square feet. In residential turfgrass, optimum disease control is achieved when Quali-Pro Myclobutanil 20EW T&O is applied in a preventative disease control program at a rate of 1.2 fl. oz. per 1000 sq. ft.

See the following tables for specific application rates for various diseases. Quali-Pro Myclobutanil 20EW T&O should be applied in sufficient water to ensure thorough coverage. For foliar diseases, use approximately one gallon of water per 1000 square feet. Two to three gallons of spray solution per 1000 square feet should be used to control diseases causing root and crown rots. Under conditions favorable for high disease development, the spray interval between applications of Quali-Pro Myclobutanil 20EW T&O should be reduced. Under light to moderate disease pressure, apply Quali-Pro Myclobutanil 20EW T&O at the lower recommended use rate in rate range and/or the longer treatment interval.

When disease pressure is high or when used as a curative treatment, use a higher rate in the suggested rate range and a shorter treatment interval unless otherwise specified.

NONRESIDENTIAL TURFGRASS+			
RESTRICTIONS: Do not apply more than 13.8 fl oz per 1000 sq ft per year. For Nassau and Suffolk Counties of New York State, do not apply more than 3.43 fl oz per 1000 sq ft per year (1.95 lb active ingredient per acre).			
DISEASE	QUALI-PRO MYCLOBUTANIL 20EW T&O (fl oz/1000 sq ft)	APPLICATION INTERVAL	USE INSTRUCTIONS
Anthracnose Red thread Septoria leaf spot	1.2	14 - 21 days	Apply when conditions are favorable for disease development.
Brown patch	1.2	14 days	Begin applications when conditions are favorable for disease development, but before disease symptoms are apparent. If disease is present, mix Quali-Pro Myclobutanil 20EW T&O with an EPA-registered contact fungicide such as Fore* T/O fungicide. Under conditions of high temperature and humidity, use the shorter spray interval.
Copper spot Zonate leaf spot	1.2	14 days	Apply when conditions are favorable for disease development.
Crown rot Leaf spot Melting-out	1.2	14 days	Apply when conditions are favorable for disease development.
Dollar spot	1.0 - 2.4	14 - 28 days	Apply when conditions are favorable for disease development. Make no more than three consecutive applications for control of dollar spot before rotating to a registered fungicide with a different mode of action.

12
 30

NONRESIDENTIAL TURFGRASS+			
RESTRICTIONS: Do not apply more than 13.8 fl oz per 1000 sq ft per year. For Nassau and Suffolk Counties of New York State, do not apply more than 3.43 fl oz per 1000 sq ft per year (1.95 lb active ingredient per acre).			
DISEASE	QUAL-PRO MYCLOBUTANIL 20EW T&O (fl oz/1000 sq ft)	APPLICATION INTERVAL	USE INSTRUCTIONS
Fusarium blight	1.2 - 2.4	14 - 21 days	Apply when conditions are favorable for disease development.
Fusarium patch (pink snow mold)	1.2 - 2.4	Fall - Winter	Apply prior to snow cover.
Leaf smuts	1.2	14 days	Apply in the fall after grass enters dormancy and/or in the spring prior to the initiation of growth.
Necrotic ring spot	1.2 - 2.4	Spring: 28 days	Make applications on a preventative basis in early to mid-spring.
		Fall: 28 days	Make two applications beginning in August before the turf goes dormant. Apply 2.4 fl oz/1000 sq ft followed by a second application one month later.
Powdery mildew Rusts	1.2	14 - 28 days	Apply when conditions are favorable for disease development.
Spring dead spot	2.4	Fall: 28 days	Make one to two applications in the fall before turf dormancy. Make 2nd application one month later.

NONRESIDENTIAL TURFGRASS+			
RESTRICTIONS: Do not apply more than 13.8 fl oz per 1000 sq ft per year. For Nassau and Suffolk Counties of New York State, do not apply more than 3.43 fl oz per 1000 sq ft per year (1.95 lb active ingredient per acre).			
DISEASE	QUALI-PRO MYCLOBUTANIL 20EW T&O (fl oz/1000 sq ft)	APPLICATION INTERVAL	USE INSTRUCTIONS
Summer patch	1.2 - 2.4	14 - 28 days	Begin applications in the spring when conditions are favorable for disease development. Make 2 to 4 applications depending on recommendations from local turfgrass extension experts. Use at least two to three gallons of water per 1000 sq ft to increase spray penetration to crown and root areas of plant.
Take-All Patch	2.4	Spring/Fall: 28 days	Apply Quali-Pro Myclobutanil 20EW T&O to reduce the severity of take-all patch. Make 1 to 2 fall applications in September and October or when night temperatures drop to 55°F and 1 to 2 spring applications in April and May depending on local recommendations.
Zoysia large patch	2.4	Fall: 28 days	Make applications in fall before turf dormancy.

+Including but not limited to commercial lawns, ornamental turf, grounds or lawns around business and office complexes, and golf course fairways, roughs, tee boxes, and greens.

14
30

RESIDENTIAL TURFGRASS			
RESTRICTIONS: Do not apply more than 13.8 fl oz per 1000 sq ft per year. For Nassau and Suffolk Counties of New York State, do not apply more than 3.43 fl oz per 1000 sq ft per year (1.95 lb active ingredient per acre).			
DISEASE	QUALI-PRO MYCLOBUTANIL 20EW T&O (fl oz/1000 sq ft)	APPLICATION INTERVAL	USE INSTRUCTIONS
Anthracnose Red thread Septoria leaf spot	1.2	14 - 21 days	Apply when conditions are favorable for disease development.
Brown patch	1.2	14 days	Begin applications when conditions are favorable for disease development, but before disease symptoms are apparent. If disease is present, mix Quali-Pro Myclobutanil 20EW T&O with an EPA-registered contact fungicide such as Fore* T/O fungicide. Under conditions of high temperature and humidity, use the shorter spray interval.
Copper spot Zonate leaf spot	1.2	14 days	Apply when conditions are favorable for disease development.
Crown rot Leaf spot Melting-out	1.2	14 days	Apply when conditions are favorable for disease development.
Dollar spot	1.2	14 days	Apply when conditions are favorable for disease development. Make no more than three consecutive applications for control of dollar spot before rotating to a registered fungicide with a different mode of action.

15
30

RESIDENTIAL TURFGRASS			
RESTRICTIONS: Do not apply more than 13.8 fl oz per 1000 sq ft per year. For Nassau and Suffolk Counties of New York State, do not apply more than 3.43 fl oz per 1000 sq ft per year (1.95 lb active ingredient per acre).			
DISEASE	QUALI-PRO MYCLOBUTANIL 20EW T&O (fl oz/1000 sq ft)	APPLICATION INTERVAL	USE INSTRUCTIONS
Fusarium blight	1.2	14	Apply when conditions are favorable for disease development.
Fusarium patch (pink snow mold)	1.2	Fall - Winter	Apply prior to snow cover.
Leaf smuts	1.2	14 days	Apply in the fall after grass enters dormancy and/or in the spring prior to the initiation of growth.
Necrotic ring spot	1.2	Spring: 28 days	Make applications on a preventative basis in early to mid-spring.
		Fall: 28 days	Make two applications beginning in August before the turf goes dormant. Apply 2.4 fl oz/1000 sq ft followed by a second application one month later.
Powdery mildew Rusts	1.2	14 - 28 days	Apply when conditions are favorable for disease development.
Summer patch	1.2	14 - 28 days	Begin applications in the spring when conditions are favorable for disease development. Make 2 to 4 applications depending on recommendations from local turfgrass extension experts. Use at least two to three gallons of water per 1000 sq ft to increase spray penetration to crown and root areas of plant.

16
30

USE DIRECTIONS FOR LANDSCAPE, GREENHOUSE, AND NURSERY ORNAMENTALS

Quali-Pro Myclobutanil 20EW T&O is a locally systemic fungicide having protectant and curative properties that will translocate to new growth. For best control of labeled diseases, achieve thorough coverage of all plant parts on a protective application schedule. For dilute sprays (>100 gallons per acre) applied to ornamental plants in greenhouses, field-grown plantings, or in commercial and residential landscapes, apply Quali-Pro Myclobutanil 20EW T&O at the rate of 6 to 12 fluid ounces per 100 gallons of spray volume on a 10- to 14-day application schedule unless otherwise directed. Use the higher rate under conditions of high disease pressure and/or optimum conditions for infection.

For concentrate sprays (<100 gallons spray volume per acre) apply 8.0 fluid ounces per acre on a 10- to 14-day application schedule.

The addition of a nonphytotoxic spreading agent will improve coverage and fungicidal performance. Treated plants should be maintained in a vigorous growing condition. Plants under nutritional or water stress will not respond as well as plants that are well maintained. Overdosage of Quali-Pro Myclobutanil 20EW T&O can result in observable foliar greening, thickened leaves, and/or shortened internodes. If this condition is observed, reduce the fungicide use rate but do not extend the recommended application schedule.

Crop Tolerance

Plant tolerances are acceptable in the specific plants listed on this label. It is not possible to evaluate all ornamental plant species or varieties for tolerance to Quali-Pro Myclobutanil 20EW T&O. The user should test for possible phytotoxic responses by treating a limited number of plants at recommended use rates prior to initiating large-scale use. The effects of spraying Quali-Pro Myclobutanil 20EW T&O in combination with plant growth regulators are not fully understood at this time. If the use of a plant growth regulator is planned in an area being treated, the user should test for possible enhanced growth regulator effects by treating a small number of plants at the recommended use rates of all products prior to initiating large-scale use. Since the effectiveness of such products depends not just on plant species or cultivar but also weather and seasonal differences (e.g., day-light hours), it is recommended that tests be repeated on previously tested varieties as environmental factors change and that observations for growth regulator responses be made at regular intervals.

SPECIFIC USE DIRECTIONS FOR CHRYSANTHEMUM

Foliar Sprays: Best control is achieved by thorough coverage sprays applied to point of runoff on a protectant application schedule. Use Quali-Pro Myclobutanil 20EW T&O at a rate of 8 fluid ounces per 100 gallons of spray mixture. (Do not apply more than 19 fluid ounces of Quali-Pro Myclobutanil 20EW T&O (0.25 pounds active) per acre per application.) Application should be made on a 10- to 14-day schedule (not to exceed 21 days).

17
30

Prestick Dip Treatment: Chrysanthemum cuttings may be treated by a dip procedure prior to planting as follows: Prepare a dip suspension at a concentration equivalent to 8 fluid ounces of Quali-Pro Myclobutanil 20EW T&O per 100 gallons of water. Cuttings must be fully submerged in the dip suspension until wet throughout. (Cuttings should not remain submerged longer than 2 minutes.) If cuttings are dipped, this procedure will represent the first spray under the quarantine program. Used dip suspension should be disposed of if it becomes contaminated with soil, plant debris, or other foreign matter. Used dip suspension can be disposed of by spraying onto registered crops (but not onto previously dipped cuttings) after filtering or in a manner consistent with local, state, and federal guidelines.

NOTE: All infected plant material must be destroyed if your state is under quarantine directive.
NOTE: Not approved for use in Nassau and Suffolk Counties, New York.

RESTRICTIONS ON USE ON ORNAMENTALS

- Do not apply more than 20 fl oz of Quali-Pro Myclobutanil 20EW T&O (0.25 lb active ingredient) per acre per application. On a total volume per acre basis, do not apply more than 333 gallons of spray per acre at the 6 fl oz per 100 gallons rate or 167 gallons per acre at the 12 fl oz per 100 gallons rate per application.
- Do not apply more than 153 fl oz of Quali-Pro Myclobutanil 20EW T&O (2 lb active ingredient) per acre per year.
- Do not use treated plant materials for food or feed.

USE DIRECTIONS FOR ORNAMENTALS

CROP	DISEASE	USE INSTRUCTIONS	RESTRICTIONS
Abelia	Cercospora leaf spot Powdery mildew		
Acalypha (Copper-leaf)	Cercospora leaf spot Powdery mildew		
Achillea (Yarrow)	Powdery mildew Rust		
African violet	Powdery mildew		
Ageratum	Rust Powdery mildew		
Alder	Powdery mildew Rust		

CROP	DISEASE	USE INSTRUCTIONS	RESTRICTIONS
Almond, flowering	Blossom blight (<i>Monilinia</i> spp.)	Apply prebloom, 50% bloom, and at petal fall.	
Amelanchier (Juneberry, Shadbush)	Fabraea leaf spot Powdery mildew Rust		
Amorpha (False indigo)	Cercospora leaf spot Powdery mildew Rust		
Anemone	Rust		
Angelica	Cercospora leaf spot Rust		
Ash	Rust		
Aster	Rust Powdery mildew		
Azalea	Petal blight (<i>Ovulinia</i> spp.) Powdery mildew	Begin applications when flowers start to exhibit color.	
Barberry	Powdery mildew Rust		May cause temporary damage to "crimson pigmy" and other "atropurposis" varieties.
Begonia	Powdery mildew		
Bellflower	Cercospora leaf spot Powdery mildew Rust		
Birch	Rust		
Bittersweet	Powdery mildew		
Buckeye	Powdery mildew		
Buttonbush	Cercospora leaf blight Powdery mildew Rust		

19
30

CROP	DISEASE	USE INSTRUCTIONS	RESTRICTIONS
Calendula	Cercospora leaf spot		
California poppy	Powdery mildew		
Canna lily	Rust		
Carnation	Powdery mildew. Rust		
Catalpa	Cercospora leaf spot Powdery mildew		
Cherry, flowering	Leaf Spot Powdery mildew		
Chestnut, horse	Powdery mildew		
China aster	Rust		
Chokeberry	Rust Twig and Fruit Blight		
Christmas trees	Rust		
Chrysanthemum	Rust White rust Ascochyta blight		
Columbine	Rust		
Cornflower	Rust		
Cosmos	Powdery mildew		
Cottonwood	Powdery mildew		
Crabapple, flowering	Powdery mildew Rust Scab		
Crape myrtle	Powdery mildew		
Daffodil	Rust		
Dahlia	Powdery mildew		
Delphinium	Powdery mildew Rust		

20
 30

CROP	DISEASE	USE INSTRUCTIONS	RESTRICTIONS
Dogwood	Anthrachnose Powdery mildew Septoria leafspot		
Douglas fir	Needle rust	Apply 12 to 18 fl oz per acre starting early spring. Continue applications at 2- to 3-week intervals until the threat of infection has passed. Spray adjuvants must be added to spray solutions to obtain good spray coverage and disease control.	
Dianthus	Rust		
Elm	Powdery mildew		
Euonymus	Powdery mildew		
Fern	Rhizoctonia aerial blight		
Fleabane	Cercospora leaf spot Powdery mildew Rust		
Four o'clock	Rust		
Fuchsia	Rust		
Gaillardia	Powdery mildew Rust		
Gardenia	Powdery mildew Rust		
Geranium	Powdery mildew Rust		
Gerbera daisy	Powdery mildew		
Gourd, ornamental	Powdery mildew		

21
30

CROP	DISEASE	USE INSTRUCTIONS	RESTRICTIONS
Grape leaf ivy	Powdery mildew		
Hackberry	Cercospora leaf spot Powdery mildew		
Hawthorn	Fabraea leaf spot Powdery mildew Rust Scab		
Holly	Powdery mildew		
Hollyhock	Powdery mildew Rust		
Honeysuckle	Cercospora leaf spot Powdery mildew		
Hydrangea	Cercospora leaf spot		
Iris	Didymellina leaf spot Rust	Apply 12 fl oz per 100 gallons of spray solution.	
Juniper	Rust		
Leucothoe	Cercospora leaf spot		
Lilac	Powdery mildew		
Loblolly pine	Fusiform rust	Refer to Douglas fir.	
Locust	Powdery mildew		
Maple	Powdery mildew		Treated trees may not be used for syrup production. Do not apply to Abutilon (Flowering Maple).
Marigold	Cercospora leaf spot Rust		
Mock-orange	Powdery mildew Rust		
Moonflower	Rust		

CROP	DISEASE	USE INSTRUCTIONS	RESTRICTIONS
Mountain laurel	Cercospora leaf spot Ovulinia petal blight Powdery mildew	Refer to Azalea.	
Nephtytis	Cephalosporium leaf spot		
Ninebark	Rust		
Oak	Powdery mildew		
Pansy	Powdery mildew Rust		
Pear, flowering	Powdery mildew Rust Scab		
Petunia	Powdery mildew Rust		
Phlox	Cercospora leaf spot Powdery mildew Rust		
Photinia	Entomosporium leaf spot Powdery mildew Rust		
Poinsettia	Powdery mildew Poinsettia scab		
Poplar	Rust		
Potentilla	Rust		
Privet	Cercospora leaf spot Powdery mildew		
Pyracantha (Firethorn)	Fusicladium scab		
Quince, flowering	Blossom and Twig Blight Cercospora leaf spot Fabraea leaf spot Rust		

CROP	DISEASE	USE INSTRUCTIONS	RESTRICTIONS
Rhododendron	Cercospora leaf spot Ovulinia petal blight Powdery mildew	Refer to Azalea.	
Rose	Black spot Powdery mildew Rust	Apply on a 7- to 10-day schedule. In areas where black spot is not a problem, spray intervals may be increased to a maximum of 14 days.	
Russian olive	Cercospora leaf spot Rust		
Salvia	Powdery mildew Rust		
Sedum	Powdery mildew		
Slash pine	Fusiform rust	Refer to Douglas fir.	
Smoke-tree (Cotinus)	Cercospora leaf spot Rust		
Snapdragon	Powdery mildew Rust		
Spirea	Powdery mildew		
Sunflower	Cercospora leaf spot Powdery mildew Rust		Seeds from treated plants may not be used for food or feed.
Sycamore	Powdery mildew		
Trumpet creeper	Cercospora leaf blight Powdery mildew		
Viburnum	Powdery mildew Rust		
Walnut	Powdery mildew		Nuts from treated trees may not be used for food purposes.

24
30

CROP	DISEASE	USE INSTRUCTIONS	RESTRICTIONS
Willow	Powdery mildew		
Zinnia	Cercospora leaf spot Powdery mildew		

GENERAL USE DIRECTIONS FOR HOME ORCHARDS, VINEYARDS, OR FRUIT TREES

Best control of labeled diseases is achieved when Quali-Pro Myclobutanil 20EW T&O is applied on a 7- to 10-day application schedule. Application equipment spray nozzles should be adjusted to apply a uniform spray throughout the entire tree canopy.

Dilute (thorough coverage) applications are recommended and are based on the amount of spray solution required to thoroughly wet plants to the point of runoff. Refer to use directions for specific tree fruits and vines to determine actual use rate per 100 gallons of spray for control of labeled diseases. The following specific use directions are based on a dilute spray volume of 300 gallons per acre.

25
 7
 30

USE DIRECTIONS FOR APPLES

DISEASE	QUALI-PRO MYCLOBUTANIL 20EW T&O (fl oz/100 gal)*	USE INSTRUCTIONS	RESTRICTIONS
Powdery mildew (<i>Podosphaera</i> spp.)	4 - 6	Begin application at tight cluster and continue through the second cover spray. Additional sprays beyond second cover may be needed on susceptible varieties or under heavy disease pressure. Use high rate in rate range if powdery mildew was present in previous years.	Do not apply within 14 days of harvest. Do not apply more than 153 fl oz Quali-Pro Myclobutanil 20EW T&O (2 lb active ingredient) per acre per season.
Rusts (<i>Gymnosporangium</i> spp.)	4 - 6	Begin applications at pink stage and continue through the second cover spray.	
Scab (<i>Venturis</i> spp.)	4 - 6	Begin application at green tip or when environmental conditions become favorable for primary scab development. Apply Quali-Pro Myclobutanil 20EW T&O alone or tank mixed with a protectant fungicide on a 7- to 10-day schedule.	
Prebloom			
Bloom, Postbloom	4 - 6	For improved fruit scab and summer disease control, use Quali-Pro Myclobutanil 20EW T&O in a tank mixture with the recommended rate of a protectant fungicide registered for use on apples.	
Post-infection	6	Quali-Pro Myclobutanil 20EW T&O provides 96-hour post-infection control or curative activity. Apply as soon as possible after infection period. Follow with a standard preventative spray schedule.	

* Based on dilute spray volume of 300 gallons per acre.

26
30

USE DIRECTIONS FOR STONE FRUIT

Applications may be made up to the day of harvest.

APRICOTS			
DISEASE	QUALI-PRO MYCLOBUTANIL 20EW T&O (fl oz/100 gal)*	USE INSTRUCTIONS	RESTRICTIONS
Brown rot blossom blight (<i>Monilinia</i> spp.)	2 - 3	Begin application at early red bud stage before infection occurs. If conditions are favorable for disease development, apply again at full bloom and petal fall.	Do not apply more than 84 fl oz of Quali-Pro Myclobutanil 20EW T&O (1.1 lb active) per acre per season.
Brown rot (<i>Monilinia</i> spp.)		Apply 12 fl oz (0.16 lb active) per acre on a 7- to 14-day schedule. Applications should be made whenever environmental conditions favor disease development during the month prior to harvest.	
Powdery mildew (<i>Podosphaera</i> spp.)		Follow brown rot blossom blight schedule making additional applications at 10- to 14-day intervals until terminal growth ceases.	
Shothole (<i>Stigminta</i> spp.)		Follow brown rot blossom blight schedule making additional applications at 7- to 10-day intervals as long as needed.	
* Based on dilute spray volume of 300 gallons per acre.			

CHERRIES			
DISEASE	QUALI-PRO MYCLOBUTANIL 20EW T&O (fl oz/100 gal)*	USE INSTRUCTIONS	RESTRICTIONS
Brown rot blossom blight (<i>Monilinia</i> spp.)	2 - 3	Begin application at early popcorn stage before infection occurs. If conditions are favorable for disease development, apply again at full bloom and petal fall.	Do not apply more than 100 fl oz of Quali-Pro Myclobutanil 20EW T&O (1.3 lb active) per acre per season.
Brown rot (<i>Monilinia</i> spp.)		Refer to Apricots.	
Powdery mildew (<i>Podosphaera</i> and <i>Sphaerotheca</i> spp.)		Refer to Apricots.	
Leaf spot (<i>Blumeriella</i> spp.)		Follow the brown rot blossom blight treatment schedule and continue applications at 7- to 10-day intervals. Make additional applications after harvest.	

* Based on dilute spray volume of 300 gallons per acre.

28
30

NECTARINES			
DISEASE	QUALI-PRO MYCLOBUTANIL 20EW T&O (fl oz/100 gal)*	USE INSTRUCTIONS	RESTRICTIONS
Brown rot blossom blight (<i>Monilinia</i> spp.)	2-3	Begin application at early pink bud stage before infection occurs. If conditions are favorable for disease development, apply again at full bloom and petal fall.	Do not apply more than 100 fl oz of Quali-Pro Myclobutanil 20EW T&O (1.3 lb active) per acre per season.
Brown rot (<i>Monilinia</i> spp.)		Refer to Apricots.	
Powdery mildew (<i>Podosphaera</i> and <i>Sphaerotheca</i> spp.)		Refer to Apricots.	
Shothole (<i>Stigmata</i> spp.)		Follow brown rot blossom blight schedule making additional applications at 7- to 10-day intervals as long as needed.	

* Based on dilute spray volume of 300 gallons per acre.

29
30

PEACHES			
DISEASE	QUALI-PRO MYCLOBUTANIL 20EW T&O (fl oz/100 gal)*	USE INSTRUCTIONS	RESTRICTIONS
Brown rot blossom blight (<i>Monilinia</i> spp.)	2 - 3	Begin application at early pink bud stage before infection occurs. If conditions are favorable for disease development, apply again at full bloom and petal fall.	Do not apply more than 100 fl oz of Quali-Pro Myclobutanil 20EW T&O (1.3 lb active) per acre per season.
Brown rot (<i>Monilinia</i> spp.)		Refer to Apricots.	
Powdery mildew (<i>Podosphaera</i> spp.)		Refer to Apricots.	
Rust (<i>Tranzschelia</i> spp.)		Apply 12 fl oz (0.16 lb active) per acre. Begin application approximately 8 weeks after flowering if environmental conditions are favorable for disease development. For optimum disease control, do not apply on an application schedule exceeding 21 days.	

* Based on dilute spray volume of 300 gallons per acre.

30
30

PLUMS/PRUNES			
DISEASE	QUALI-PRO MYCLOBUTANIL 20EW T&O (fl oz/100 gal)*	USE INSTRUCTIONS	RESTRICTIONS
Brown rot blossom blight (<i>Monilinia</i> spp.)	2 - 3	Begin application at green tip before infection occurs. If conditions are favorable for disease development, apply again at full bloom and petal fall.	Do not apply more than 84 fl oz of Quali-Pro Myclobutanil 20EW T&O (1.1 lb active) per acre per season.
Rust (<i>Tranzschelia</i> spp.)		Refer to Peaches.	

* Based on dilute spray volume of 300 gallons per acre.