

66 222-210

8/13/2012

1/7



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

8/13/2012

Kristen B. Knox
Makhteshim Agan of North America, Inc.
3120 Highwoods Blvd., Suite 100
Raleigh, NC 27604

re: QualiPro Abamectin 0.15 EC, EPA Reg. # 66222-210
label amendment submitted 7/10/2012 (D# 468516)
accepted

Dear Ms. Knox:


This label amendment deletes oxidation potential text in the Physical or Chemical Hazards section which had been included by mistake on a label for the above product accepted on 7/6/2012. Both the original product chemistry review and the product chemistry study (MRID 457080-02) support the conclusion that no oxidation statement is required for this product.

The revised labeling reference above, submitted in connection with the registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is acceptable provided that you submit two copies of the revised final printed label for the record before you release the product for shipment.

If the above provision is not complied with the registration will be subject to cancellation in accordance with FIFRA Section 6(e). Your release for shipment of the product bearing the amended labeling constitutes acceptance of these conditions.

A copy of the label stamped "accepted" is enclosed for your records. If you have any questions please contact me at (703) 308-9423 or harris.thomas@epa.gov.

Sincerely yours,



John D. Hebert
Product Manager 7
Insecticide Rodenticide Branch
Registration Division (7505C)

enclosure

2/7

066222-00210.20120710.CLEAN Label Amendment to Correct Physical Chemical Hazard Statement Quali-Pro Abamectin 0.15

Quali-Pro[®] Abamectin 0.15 EC

Miticide/Insecticide

For Agricultural/Commercial Use to Control Leafminers and Mites, and to Suppress Aphids, Whiteflies, and Thrips on Ornamental Plants.

ACTIVE INGREDIENT:

Abamectin (CAS No. 65195-56-4 and 65195-55-3): 1.9%

INERT INGREDIENTS: 98.1%

Total: 100.0%

1 gallon contains 0.15 lb. abamectin

KEEP OUT OF REACH OF CHILDREN WARNING / AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

Manufactured for:
Makhteshim Agan of North America, Inc.
3120 Highwoods Blvd., Suite 100
Raleigh, NC 27604

EPA Reg. No. 66222-210

EPA Est. No. [REDACTED]

NET CONTENTS: QUART

FIRST AID

IF SWALLOWED:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Do not give any liquid to the person. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
IF IN EYES:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor immediately for treatment advice.
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of soap and water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
IF INHALED:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. • Call a poison control center or doctor for treatment advice

NOTE TO PHYSICIAN: Early signs of intoxication include mydriasis (dilated pupils), ataxia (unsteadiness), and muscle tremors. Toxicity follows accidental ingestion of Quali-Pro[®] Abamectin 0.15 EC and can be minimized by promptly administering activated charcoal. If toxicity from exposure has progressed to cause severe vomiting, the extent of resultant fluid and electrolyte imbalance should be gauged. Appropriate supportive parental fluid replacement therapy should be given, along with other required supportive measures (such as maintenance of blood pressure levels and respiratory functionality) as indicated by clinical signs, symptoms, and measurements. In severe cases, observations should continue for at least several days until clinical condition is stable and normal. Since abamectin is believed to enhance GABA activity in animals, it is probably wise to avoid drugs that enhance GABA activity (barbiturates, benzodiazepines, valproic acid) in patients with potentially toxic abamectin exposure.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact PROSAR at 1-877-250-9291 for emergency medical treatment information

ACCEPTED
8/13/2012
Under the Federal Insecticide, Fungicide, and Rodenticide Act, As amended, for the pesticide registered under EPA Reg. No. 66222-210

Quali-Pro Abamectin 0.15 EC; EPA #66222-210
Amendment to correct phys/chem. hazard
Clean Copy Label; July 10, 2012
Page 1 of 6

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
WARNING**

May be fatal if swallowed Causes substantial but temporary eye injury Do not get in eyes Harmful if absorbed through skin or inhaled Avoid contact with skin or clothing Avoid breathing spray mist Wash thoroughly with soap and water after handling and before eating drinking chewing gum or using tobacco Remove and wash contaminated clothing before reuse

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are listed below For more options follow the instructions for Category B on an EPA chemical-resistance category selection chart

Applicators and other handlers must wear

- Coveralls over short-sleeved shirt and short pants
- Chemical-resistant gloves such as barrier laminate or butyl rubber ≥ 14 mils
- Chemical resistant footwear plus socks
- Protective eyewear
- Chemical-resistant headgear for overhead exposure
- Chemical-resistant apron when cleaning equipment mixing or loading
- For shadehouse and greenhouse uses applicators and other handlers must wear a dust/mist filtering respirator (MSHA/NIOSH approval prefix number TC-21C or a NIOSH approved respirator with any R P or HE filter)

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate Do not reuse them Follow manufacturer's instructions for cleaning/maintaining PPE If no such instructions for washables use detergent and hot water Keep and wash PPE separately from other laundry

USER SAFETY RECOMMENDATIONS

Users Should

- Wash hands before eating drinking chewing gum using tobacco or using the toilet
- Remove clothing immediately if pesticide gets inside Then wash thoroughly and put on clean clothing

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and wildlife Do not apply directly to water to areas where surface water is present or to intertidal areas below the mean high water mark Do not contaminate water when disposing of equipment wash water or rinsate This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area Use of this product may pose a risk to threatened and endangered species of fish amphibians crustaceans (including fresh water shrimp) and insects All use of this product in the state of California should comply with the recommendations of the California Endangered Species Project Before using this product in California consult with your county agriculture commissioner to determine use limitations that apply in your area

This product may impact surface water quality due to runoff of rain water This is especially true for poorly draining soils and soils with shallow ground water This product is classified as having a medium potential for reaching both surface water and aquatic sediment via runoff for several weeks to months after application A level well-maintained vegetative buffer strip between areas to which this product is applied and surface water such as ponds streams and springs will reduce the potential loading of abamectin from runoff water and sediment Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours

PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat or open flame

Attention This product contains a chemical known to the State of California to cause birth defects or other reproductive harm

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Do not apply Quali-Pro® Abamectin 0.15 EC through any type of irrigation system. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls over short-sleeved shirt and short pants
- Chemical-resistant gloves, such as barrier laminate or butyl rubber ≥ 14 mils
- Chemical-resistant footwear plus socks
- Protective eyewear
- Chemical-resistant headgear for overhead exposure

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are not within the scope of the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not reenter treated areas until sprays have dried.

For Use on Greenhouse-, Shadehouse-, and Field-Grown Ornamentals, Foliage Plants, Christmas Trees, and other Woody Ornamentals.

Quali-Pro Abamectin 0.15 EC is an emulsifiable concentrate containing 0.15 lb. abamectin per gallon. When mixed with water according to the directions for use, Quali-Pro Abamectin 0.15 EC will control mites and leafminers, and suppress thrips, aphids, and whiteflies on the following ornamental plants: Greenhouse-, shadehouse- and field-grown ornamentals; foliage plants; Christmas trees and other woody ornamentals.

To control mites, apply Quali-Pro Abamectin 0.15 EC when mites initially appear, then repeat as needed to maintain control. To control leafminers, apply Quali-Pro Abamectin 0.15 EC and repeat every 7 days or as needed to maintain control.

To suppress whiteflies, thrips, and aphids, apply Quali-Pro Abamectin 0.15 EC when the young/immature stages of these pests are first seen, then repeat once per week for 2–3 weeks. Then rotate to other registered insecticides having a different mode of action for a minimum of 2–3 weeks before reverting to Quali-Pro Abamectin 0.15 EC. Please see the **RESISTANCE MANAGEMENT** section of this label for additional information regarding rotating products.

Note: For Quali-Pro Abamectin 0.15 EC to kill aphids, whiteflies, and thrips, it must come in direct contact with the pests.

To obtain uniform coverage, mix Quali-Pro Abamectin 0.15 EC with sufficient water and apply as a foliar spray. Residual control of pests can be improved on field-grown woody ornamentals, Christmas trees, and landscape plants by adding a Horticultural Spray Oil (not dormant oil) at 2–4 qts per 100 gals finished spray (equivalent to 0.5–1.0% of spray volume). To maintain control, repeat application of Quali-Pro Abamectin 0.15 EC as necessary, but not sooner than 7 days after previous application. Some plants are sensitive to oils. Therefore, unless prior experience has shown plant safety, the user should apply an Quali-Pro Abamectin 0.15 EC plus Horticultural Spray Oil mixture on a small number of plants and observe them for two weeks. If no plant damage is observed,

then spray the remaining plants Excessively warm or cold temperatures following treatment in combination with oils could increase the chances of plant damage Read and follow label directions for both Quali-Pro Abamectin 0 15 EC and any horticultural spray oil product to be used and do not exceed the maximum rates detailed on either label

Do not apply Quali-Pro Abamectin 0 15 EC in quantities less than 8 fl oz per acre or more than 16 fl oz per acre per application Use a sufficient quantity of water to obtain thorough coverage on foliage Review the **RESISTANCE MANAGEMENT** and **RESTRICTIONS** sections (below) for additional guidance

Greenhouse-, Shadehouse-, and Field-Grown Ornamentals, Foliage Plants, Christmas Trees, and other Woody Ornamentals

Pests	Amount of Quali-Pro Abamectin 0 15 EC fl oz per 100 gals water	Notes
Mites European Red Mite Two-spotted Spider Mite Carmine Spider Mite Southern Red Mite Spruce Spider Mite Tarsonemid Mites Cyclamen & Broad Mites Eriophyid Mites Rust and Bud Mites	4 ^a	For tarsonemid mites multiple applications to newly developing tissue may be needed to maintain effective control
Boxwood Leafminer	8	To control mining larvae apply when adults begin to lay eggs onto new foliage
<i>Liriomyza</i> Leafminers	8 ^b	Repeat at intervals of 7 days or as needed to maintain effective control
Aphids Thrips & Whiteflies	8	Spray must contact young immatures

^a For example apply in 200–400 gals of water per acre If water volume is less than 200 gals use a minimum of 8 fl oz per acre If water volume of 400 gals or more per acre is required for sufficient plant coverage apply at the maximum rate of 16 fl oz per acre Another example if a volume of 650 gals of water is necessary use 2.5 fl oz per 100 gals

^b For example apply in 100–200 gals of water per acre If water volume is less than 100 gals use at least 8 fl oz per acre If water volume of 200 gals or more per acre is required for sufficient plant coverage apply the maximum rate of 16 fl oz per acre Another example if 400 gals of water is necessary apply 4 fl oz per 100 gals

Use Restrictions

- Do not apply by air
- Do not use Quali-Pro Abamectin 0 15 EC for suppressing thrips whiteflies and aphids on gerbera chrysanthemums and roses which are primary hosts of *Liriomyza* leafminers and mites controlled by Quali-Pro Abamectin 0 15 EC Additional applications of Quali-Pro Abamectin 0 15 EC on these plants will increase the selection pressure on mites and *Liriomyza* leafminers which could produce greater pest tolerance (resistance) to Quali-Pro Abamectin 0 15 EC
- Do not use Quali-Pro Abamectin 0 15 EC in citrus nurseries
- Do not use Quali-Pro Abamectin 0 15 EC on ferns or Shasta Daisy since phytotoxicity may occur on certain species of ferns (e.g. *Adiantum* spp.) and Shasta Daisy (*L. eucanthemum* spp.)
- Do not apply through any type of irrigation system

Quali Pro Abamectin 0 15 EC is non-phytotoxic on a wide range of ornamental plants However since all combinations or sequences of pesticide sprays including surfactants and adjuvants have not been tested it is recommended that a small area be sprayed first to make certain that no phytotoxicity occurs

Resistance Management

To minimize potential development of pest resistance to abamectin use of resistance management strategies is imperative especially when applying Quali-Pro Abamectin 0.15 EC to ornamental plants that are hosts of pests on this label that have multiple generations per crop or per year. Examples of resistance management may include but are not limited to the following: use of various cultural practices, nonchemical alternatives (such as beneficial arthropods), rotating susceptible and nonsusceptible plants, rotating pesticides with different modes of action, avoiding use of the same product on successive pest generations, and using labeled rates at specified spray intervals. Consult your State Cooperative Extension Service to obtain additional information regarding the implementation of these or other resistance management strategies.

For managing susceptibility in *Liriomyza* leafminers, apply Quali-Pro Abamectin 0.15 EC no more than three times, then apply a registered alternative product with a different mode of action for no more than three times before rotating back to Quali-Pro Abamectin 0.15 EC or another alternative product. Product rotation intervals should be based on the generation time of *Liriomyza* leafminers to avoid applying Quali-Pro Abamectin 0.15 EC to successive generations.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited. Do not reuse empty container.

PESTICIDE STORAGE Store in a tightly closed container in a cool, dry place.

PESTICIDE DISPOSAL Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING

Nonrefillable Container (five gallons or less) Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Dispose of in an incinerator or landfill approved for pesticide containers.

Nonrefillable Container (greater than five gallons) Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Dispose of in an incinerator or landfill approved for pesticide containers.

Refillable Container Refillable container. Refill this container with abamectin only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup procedures and disposal of wastes.

IMPORTANT Read the information below before using this product. If the terms are not acceptable, you should return the unopened product container immediately for a complete refund.

LIMITATION OF WARRANTY AND LIABILITY

Read the entire directions for use, conditions of warranties, and limitations of liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following **CONDITIONS, DISCLAIMER OF WARRANTIES, and LIMITATIONS OF LIABILITY**.

CONDITIONS The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other

materials or the manner of use or application all of which are beyond the control of Makhteshim Agan of North America Inc All such risks shall be assumed by the user or buyer

DISCLAIMER OF WARRANTIES To the extent consistent with applicable law Makhteshim Agan of North America Inc makes no other warranties express or implied of merchantability or of fitness for a particular purpose or otherwise that extend beyond the statements made on this label No agent of Makhteshim Agan of North America Inc is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein To the extent consistent with applicable law Makhteshim Agan of North America Inc disclaims any liability whatsoever for special incidental or consequential damages resulting from the use or handling of this product

LIMITATIONS OF LIABILITY To the extent consistent with applicable law the exclusive remedy of the user or buyer for any and all losses injuries or damages resulting from the use or handling of this product whether in contract warranty tort negligence strict liability or otherwise shall not exceed the purchase price paid or at Makhteshim Agan of North America Inc s election the replacement of product

Quali-Pro is a registered trademark of Makhteshim Agan of North America Inc

Quali Pro Abamectin 0.15 EC (66222-210) (EPA Approved 10-20-04) (EPA Transferred 01-04-10) (Notif to EPA 01-29-10)
(Amendment to EPA 05-04-12)