

PM 04
66330-6
82-11-99

CAPTAN METHOXYCHLOR 75-3 SEED PROTECTANT
(INSECTICIDE-FUNGICIDE)

ACTIVE INGREDIENTS:	By. Wt.
*Captan.....	73.3%
Related Derivatives.....	1.7%
**Methoxychlor, Technical.....	3.0%
INERT INGREDIENTS:	22.0%
* N-Trichloromethylthio-4-cyclohexene-1,2-dicarboximide	
** Equivalent to 2.64% 2,2-bis (p-methoxyphenyl)-1,1,1-trichloroethane and 0.36% of other isomers and related compounds	

KEEP OUT OF REACH OF CHILDREN

DANGER

**SEE SIDE PANEL FOR ADDITIONAL
PRECAUTIONARY STATEMENTS**

STATEMENT OF PRACTICAL TREATMENT

If swallowed, contact physician, poison control center, or emergency clinic before inducing vomiting. Give water or milk to drink and induce vomiting by touching back of throat with finger, or by administering syrup of ipecac. Do not induce vomiting or give anything by mouth to an unconscious person. If medical advice cannot be obtained, take person and a copy of the label to the nearest emergency treatment facility. In case of eye contact, immediately flush eyes with plenty of clear water and get medical attention. For skin contact, wash skin with plenty of soap and water. If inhaled, remove person from exposure area.

**FOR 24-HOUR EMERGENCY MEDICAL ASSISTANCE CALL (800) F-A-S-T-M-E-D
(327-8633)1-800-228-5635 OR 1-612-221-3999 ext. 174.**

**FOR CHEMICAL EMERGENCY: Spill, leak, fire, exposure, or accident call
CHEMTREC 1-800-424-9300.**

EPA Reg. No. 40182-31266330-6

EPA Est. No. 40182-
Net Weight:

Made in U.S.A
ICI Agricultural Products **TOMEN AGRO, Inc.**
ICI Americas, Inc. 100 First Street, Suite 1700
Wilmington DE 19897 San Francisco, CA 94105

ACCEPTED

FEB 11 1999

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 66330-6

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

DANGER

Causes irreversible eye damage. Harmful if swallowed or inhaled. May cause allergic skin reactions. Do not get in eyes. Wear goggles or face shield when handling. Avoid contact with skin and clothing. Avoid inhalation of dust. Remove and separately launder clothing before reuse.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not clean equipment or dispose of equipment washwaters in a manner that will contaminate water resources. In the event that treated seeds are spilled and cannot be salvaged, completely cover them with soil.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

Mixers/loaders when mixing and loading this product must wear a long-sleeved shirt and long pants or coverall that covers all parts of the body except the head, hands and feet. Wear chemical-resistant gloves, goggles or a face shield and a chemically-resistant apron.

A dust mask must be worn when mixing and loading this product.

AFTER WORK: Take off clothes and shoes. Shower using soap and water. Do not wear contaminated clothing. Wash protective clothing and equipment with soap and water after each use. Personal clothing worn during use must be laundered separately from household articles. Clothing and protective equipment heavily contaminated or drenched with CAPTAN must be destroyed according to state and local regulations.

CAPTAN Methoxychlor 75-3 Seed Protectant is a special formulation containing CAPTAN and Methoxychlor for seed application. CAPTAN Methoxychlor 75-3 Seed Protectant now contains a binder which helps to adhere the treatant to the seed so that there is minimum loss during processing and planting. Has no adverse effect on seed germination or planting rate. CAPTAN Methoxychlor 75-3 Seed Protectant is used in preventing seed rot and damp off diseases on germinating seeds and seedlings and smut on onion and hulled sorghum seeds. It also reduces losses from Cadelle, Rice Weevil, Confused Flour Beetle, Angoumois Grain Moth and Indian Meal Moth during preplanting storage of clean seed.

DIRECTIONS: Apply as a slurry Treatment. The recommended amount of product should be mixed with the required amount of water for the slurry treater machine to be used. Do not allow slurry mixtures to stand overnight in the machine. Clean out equipment thoroughly between treating operations.

GLADIOLUS CORMS: For prevention of Blub Rot – Use 1½ oz. in 1 gal. water (1 lb. to 10 gals.). Soak bulbs in mixture 20 to 30 minutes.

ONION SEED PELLETING: Use 1 lb. CAPTAN Methoxychlor 75-3 Seed Protectant per 1lb. of seed.

Do not feed treated seed to animals. Do not use treated seed for human consumption. Seeds treated with CAPTAN Methoxychlor 75-3 Seed Protectant, sold or shipped as feed or foodstuffs may be considered adulterated under state and federal laws. Treated seed must be labeled as follows: "TREATED SEED. DO NOT USE FOR FOOD, FEED OR OIL".

THE FOLLOWING TABLE gives amount of CAPTAN Methoxychlor 75-3 Seed Protectant to use on various seeds. Do not use on baby lima beans.

Seed	Dosage Rate	Lbs. Seed Treated/lb. of Product	
Corn - Field	1¼ oz./100 lbs. ⅔ oz./bu.	1280	
Corn - Sweet	2⅔ oz./100 lbs. 1½ oz./bu.	600	
Wheat	For seed and Seedling Rots-	1¼ oz./100 lbs. ⅔ oz./bu.	1280
Rye	Not for	1¾ oz./100 lbs.	910
Oats	Smut	2 oz./100 lbs. ⅔ oz./bu.	800
Barley		1½ oz./100 lbs. ⅔ oz./bu.	1070
Flax		2½ oz./100 lbs. 1½ oz./bu.	640
Beans (Except Baby Limas)		1¾ oz./100 lbs.	910
Peas		1 oz./bu.	
Cucumbers, Cantaloupe		2 oz./100 lbs.	800
Squash, Pumpkin, Watermelon		1⅓ oz./100 lbs.	1200
Beets, Swiss Chard		8 oz./100 lbs.	200
Sugar Beets		8 oz./100 lbs.	200
Clover		5½ oz./100 lbs.	290
Alfalfa		3⅓ oz./bu.	
Soybeans		1⅔ oz./100 lbs. 1 oz./bu.	960
Cotton - Acid Delinted		2 oz./100 lbs.	800
Cotton - Machine Delinted Fuzzy (not delinted)		3 oz./100 lbs.	530
Blue Grass		5½ oz./100 lbs.	290
Cabbage, Cauliflower, Broccoli, Brussels Sprouts		1 oz./100 lbs.	1600
Spinach		4 oz./100 lbs.	400
Sorghum hulled (Kernal Smut)		3 oz./100 lbs.	530
Peppers		2 oz./100 lbs.	800
Cowpeas		2 oz./100 lbs. 1¼ oz./bu.	800

STORAGE AND DISPOSAL

PROHIBITIONS: Do not contaminate water, food or feed by storage, disposal or cleaning of equipment. Open dumping is prohibited.

STORAGE: Keep pesticide in original container. Do not store or transport near feed or food. Store in a cool, dry place. Protect from excessive heat. For help with any spill, leak, fire or exposure involving this material, call day or night 1-800-424-9300.

PESTICIDE DISPOSAL: This product is acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Completely empty liner by shaking and taping sides and bottom to loosen clinging particles. Empty residue into application equipment. Do not reuse container. Dispose of liner in a sanitary landfill or by incineration if allowed by State and local authorities. If drum is contaminated and cannot be reused, dispose of in the same manner.

~~**NOTICE TO BUYER AND USER:** Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal use conditions, or under conditions not reasonably foreseeable to Seller, and Buyer and User assume the risk of any such use. SELLER DISCLAIMS ALL OTHER WARRANTIES, EXPRESSED OR IMPLIED, INCLUDING ANY WARRANTY OF FITNESS OR MERCHANTABILITY. SELLER SHALL NOT BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT AND SELLER'S SOLE LIABILITY AND BUYER'S AND USER'S EXCLUSIVE REMEDY SHALL BE LIMITED TO THE REFUND OF THE PURCHASE PRICE.~~

NOTICE TO BUYER AND USER:

CONDITIONS OF SALE:

1. TOMEN AGRO, Inc. warrants that this product conforms to the chemical description on the label and is reasonable fit for the purposes stated on the label when used in accordance with the directions under normal conditions of use.
2. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal use conditions, or under conditions not reasonably foreseeable to Tomen. TOMEN DISCLAIMS ALL OTHER WARRANTIES, EXPRESSED OR IMPLIED, INCLUDING ANY WARRANTY OF FITNESS OR MERCHANTABILITY. SELLER SHALL NOT BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT

DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT AND SELLER'S SOLE LIABILITY AND BUYER'S AND USER'S EXCLUSIVE REMEDY SHALL BE LIMITED TO THE REFUND OF THE PURCHASE PRICE. TOMEN DOES NOT AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE ANY OTHER WARRANTY, GUARANTEE OR REPRESENTATION CONCERNING THIS PRODUCT.

3. Critical and unforeseeable factors beyond Tomen's control prevent it from eliminating all risks in connection with the use of this product. Such risks include, but are not limited to, damage to plants and crops to which the product is applied, lack of complete control, and damage caused by drift to other plants or crops. Such risks occur even though the product is reasonably fit for the uses stated on the label and even though label directions are followed. Except as stated in 1 above, Buyer and User acknowledge and assume all risks and liability resulting from handling, storage, and use of this product.

RQ Hazardous substance, solid, N.O.S., NA 9188,
(Captan; Methoxychlor)

ORM-E