



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, D.C. 20460

OFFICE OF  
PREVENTION, PESTICIDES AND  
TOXIC SUBSTANCES

Bill Washburn  
Regulatory Manager  
Arysta LifeScience North America LLC  
15401 Weston Parkway, Suite 150  
Cary, NC 27513

JAN 29 2010

Subject: Label Notification(s) for Pesticide Registration Notices 2007-4 and 98-10

1. **Update of company name**
2. **Added emergency telephone number(s)**
3. **Updated warranty statement**

Dear Mr. Washburn:

The Agency is in receipt of your Application(s) for Pesticide Notification under Pesticide Registration Notices (PRN) 2007-4 and 98-10 dated October 28, 2009 for:

**EPA Registration 66330-35**

**Elevate 50 WDG Fungicide**

The Registration Division (RD) has conducted a review of this request for applicability under PR Notices 2007-4 and 98-10 and finds that the label changes requested falls within the scope of PR Notices 2007-4 and 98-10. The label has been date-stamped "Notification" and will be placed in our records.

Please be reminded that 40 CFR Part 156.140(a)(4) requires that a batch code, lot number, or other code identifying the batch of the pesticide distributed and sold be placed on nonrefillable containers. The code may appear either on the label (and can be added by non-notification/PR Notice 98-10) or durably marked on the container itself.

If you have any questions, please contact me directly at 703-305-6249 or Nicole Williams of my staff at 703-308-5551.

Sincerely,





Linda Arrington  
Notifications & Minor Formulations Team Leader  
Registration Division (7505P)  
Office of Pesticide Programs

2028

Please read instructions on reverse before completing form.

Form Approved, OMB No. 2070-0060, Approval expires 05-31-98

 <b>EPA</b> United States Environmental Protection Agency Washington, DC 20460		<input type="checkbox"/> Registration <input type="checkbox"/> Amendment <input checked="" type="checkbox"/> Other:	OPP Identifier Number
<b>Application for Pesticide - Section I</b>			
1. Company/Product Number <b>66330-35</b>		2. EPA Product Manager <b>Mary waller</b>	
4. Company/Product (Name) <b>ELEVATE 50 WDG FUNGICIDE</b>		3. Proposed Classification <input type="checkbox"/> None <input type="checkbox"/> Restricted	
5. Name and Address of Applicant (Include ZIP Code)  <b>Arysta LifeScience North America, LLC</b> <b>15401 Weston Parkway, Suite 150</b> <b>Cary, NC 27513</b> <input type="checkbox"/> Check if this is a new address		6. <b>Expedited Review.</b> In accordance with FIFRA Section 3(c)(3) (b)(I), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	
<b>Section - II</b>			
<input type="checkbox"/> Amendment – Explain below. <input type="checkbox"/> Resubmission in response to Agency letter dated _____ <input checked="" type="checkbox"/> Notification - Explain below.		<b>NOTIFICATION</b> Final printed labels in response to Agency letter dated <b>JAN 29 2010</b> <input type="checkbox"/> "Me Too" Application <input type="checkbox"/> Other - Explain below	
<b>Explanation:</b> Use additional page(s) if necessary. (For Section I and Section II.) <b>Notification of label change per PR Notice 2007-4.</b> This notification is consistent with guidance in PR Notice 2007-4 and the requirements of EPA's regulations at 40 CFR §§ 156.10, 156.140, 156.144, 156.146, and 156.156. No other changes have been made to the labeling or the Confidential Statement of Formula for this product. I understand that it is a violation of 18 USC Sec 1001 to willfully make any false statement to EPA. I further understand that if the amendment is not consistent with the requirements of 40 CFR §§ 156.10, 156.140, 156.144, 156.146, and 156.156, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA. <b>Notification of label change per PR Notice 98-10.</b> This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 USC Sec 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA. <b>Update company name and emergency telephone numbers.</b>			
<b>Section - III</b>			
1. Material This Product Will Be Packaged In:			
Child-Resistant Packaging <input type="checkbox"/> Yes* <input checked="" type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No If "Yes" Unit Packaging wgt.      No. per container	Water Soluble Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No If "Yes" Package wgt.      No. per container	2. Type of Container <input type="checkbox"/> Metal <input checked="" type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify)
* <b>Certification must be submitted</b>			
3. Location of Net Contents Information <input checked="" type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container <b>5 lb., 10 lb, 40 lb.</b>	
		5. Location of Label Directions <input checked="" type="checkbox"/> On Label <input type="checkbox"/> On labeling accompanying product	
6. Manner in Which Label is Affixed to Product		<input type="checkbox"/> Lithograph <input checked="" type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled <input type="checkbox"/> Other _____	
<b>Section - IV</b>			
1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application)			
Name <b>Bill Washburn</b>		Title <b>Regulatory Manager</b> Telephone No. (Include Area Code) <b>901-432-5118</b>	
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.			6. Date Application Received <b>(Stamped)</b>
2. Signature 		3. Title <b>Regulatory Manager</b>	
4. Typed Name <b>Bill Washburn</b>		5. Date <b>October 28, 2009</b>	

30 of 28



Arysta LifeScience

October 28, 2009

Section IV  
refill label

Ms. Mary Waller, PM 21  
Document Processing Desk (NOTIF)  
Office of Pesticide Programs – 750PC  
U.S. Environmental Protection Agency  
One Potomac Yard, Room S-4900  
2777 South Crystal Drive  
Arlington, VA 22202

**Subject: Elevate 50 WDG Fungicide  
EPA Reg. No. 66330-35**

***Notification of Label Change per PR Notice 2007-4  
and PR Notice 98-10***

Dear Ms. Waller:

Please find the following enclosed:

- Application for Pesticide Registration (Other) dated 10/28/09.
- One highlighted copy of subject label, showing all changes.
- One clean copy of the subject label.

Notification of label change per PR Notice 2007-4. This notification is consistent with guidance in PR Notice 2007-4 and the requirements of EPA's regulations at 40 CFR §§ 156.10, 156.140, 156.144, 156.146, and 156.156. No other changes have been made to the labeling or the Confidential Statement of Formula for this product. I understand that it is a violation of 18 USC Sec 1001 to willfully make any false statement to EPA. I further understand that if the amendment is not consistent with the requirements of 40 CFR §§ 156.10, 156.140, 156.144, 156.146, and 156.156, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 USC Sec 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

We are using PR Notice 98-10 to update the name of the company, emergency telephone numbers and warranty. Arysta LifeScience North America Corporation was changed to Arysta LifeScience North America, LLC.

40828

Please acknowledge acceptance of this notification. Should you have any questions or comments, please do not hesitate to contact me at 901-432-5118 or by e-mail at [bill.washburn@arystalifescience.com](mailto:bill.washburn@arystalifescience.com)

Sincerely,

Bill Washburn

Bill Washburn  
Regulatory Manager

Bill Washburn  
Regulatory Manager  
Arysta LifeScience  
40828

50 of 28

[Alternate language appear in brackets]  
[Individual Sub-label language denoted by e, d, or j]

## ELEVATE® 50 WDG Fungicide

GROUP 17 FUNGICIDE

[ 50% Water Dispersible Granule for Control of Botrytis and Monilinia Diseases  
and Suppression of Powdery Mildew in Various Crops ]<sub>e</sub>

[ 50% Water Dispersible Granule for Control of Botrytis and Monilinia Diseases in  
Various Post-Harvest Fruit Treatments ]<sub>j</sub>

[ FOR AGRICULTURAL USE ONLY ]<sub>e,j</sub>

[ 50% Water Dispersible Granule for Control of Botrytis in Various Greenhouse  
Crops, Botrytis Diseases in Ornamentals, Forestry Conifers and Non-Bearing  
Fruit Trees and Vines and for the Suppression of Powdery Mildew in  
Ornamentals ]<sub>d</sub>

[ FOR AGRICULTURAL AND ORNAMENTAL HORTICULTURAL USE ONLY ]<sub>d</sub>

Not for Residential Use **NOTIFICATION**

JAN 29 2010

### INGREDIENTS

#### ACTIVE INGREDIENT:

Fenhexamid\* ..... 50.0%

OTHER INGREDIENTS: ..... 50.0%

Total..... 100.0%

\* N-(2,3-dichloro-4-hydroxyphenyl)-1-methyl-cyclohexane carboxamide

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

See back panel for additional precautionary statements.

For Product Use Information Call: 1-866-761-9397

Produced for:

Arysta LifeScience North America, LLC Corporation  
15401 Weston Parkway, Suite 150  
Cary, NC 27513

EPA Reg. No. 66330-35

EPA Est. No.

NET WEIGHT: \_\_\_\_ POUNDS ( \_\_\_\_ KILOGRAMS)

[Alternate language appear in brackets]  
 [Individual Sub-label language denoted by e, d, or j]

<b>FIRST AID</b>	
<b>If swallowed</b>	<ul style="list-style-type: none"> <li>• Call poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told so by the poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
<b>If in eyes</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with warm water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If on skin or clothing</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
Have the product container or label with you when calling a poison control center or doctor, or going for treatment.	
<b>FOR 24-HOUR EMERGENCY MEDICAL ASSISTANCE:</b> Call PROSAR at 1-866-303-6952 or 1-651-632-8946 if calling from outside of the U.S.	
<b>FOR 24-HOUR CHEMICAL EMERGENCY</b> Spill, leak, fire, exposure, or accident, call CHEMTREC at 1-800-434-9300 or 1-703-527-3887 if calling from outside of the U.S.	

## **PRECAUTIONARY STATEMENTS**

### **HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**CAUTION.** Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes and clothing. Wash thoroughly with soap and water after handling.

### **PERSONAL PROTECTIVE EQUIPMENT (PPE)**

Applicators and other handlers must wear long-sleeved shirt and long pants, socks, shoes and chemical-resistant gloves made of any waterproof material. Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables are available, use detergent and hot water. Keep and wash PPE separately from other laundry.

### **ENGINEERING CONTROLS**

When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

[Alternate language appear in brackets]  
[Individual Sub-label language denoted by e, d, or j]

**IMPORTANT:** When reduced PPE is worn because a closed system is being used, handlers must be provided all PPE specified above for "applicators and other handlers" and have such PPE immediately available for use in an emergency, such as a spill or equipment break-down.

#### USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

#### ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates. Drift and runoff may be hazardous to aquatic organisms in neighboring areas. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters or rinsate.

#### DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not apply this product in any way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

#### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions on this label about personal protective equipment (PPE), notification to workers, and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry into treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is: coveralls, socks, shoes, chemical resistant gloves made of waterproof material.

Carefully read and understand the Directions for Use and restrictions before applying this product.

[Alternate language appear in brackets]  
[Individual Sub-label language denoted by e, d, or j]

## GENERAL INFORMATION

[ ELEVATE® 50 WDG Fungicide is a hydroxylanilide protectant fungicide used for the control of Gray mold/Fruit Rot (*Botrytis cinerea*) on various fruits and vegetables. In addition, ELEVATE 50 WDG also controls Botrytis blossom and shoot blight on pistachios and Brown rot, Blossom blight and Twig blight (*Monilinia* spp.) on stone fruit and almonds. When ELEVATE 50 WDG is used to control Botrytis diseases, suppression of powdery mildew (*Uncinula necator*) on grapes may be achieved. ELEVATE 50 WDG should be used in a program with other fungicides that are labeled for the control of powdery mildew. ELEVATE 50 WDG should be used in a program with other products to provide season-long Botrytis protection. Apply as directed in the "APPLICATION" section using sufficient water to obtain thorough coverage of plants. Under severe disease conditions, use maximum rates and shorter application intervals as specified on the label. Dense foliage or excessive growth will often prevent adequate coverage; adjust spray volumes accordingly. ]<sup>e</sup>

[ ELEVATE® 50 WDG Fungicide is a hydroxylanilide protectant fungicide used for the control of Gray mold (*Botrytis cinerea*) on various greenhouse vegetables and for control of Gray mold or Botrytis blights (*Botrytis cinerea*, *Botrytis elliptica* and *Botrytis tulipae*) on ornamental crops, forestry conifers and non-bearing fruit trees and vines. When ELEVATE 50 WDG is used to control Botrytis diseases, suppression of powdery mildew (*Uncinula necator*) on ornamental crops may be achieved. ELEVATE 50 WDG should be used in a program with other fungicides that are labeled for the control of powdery mildew. ELEVATE 50 WDG should be used in a program with other products to provide season-long Botrytis protection. Apply as directed in the "APPLICATION" section using sufficient water to obtain thorough coverage of plants. Under severe disease conditions, use maximum rates and shorter application intervals as specified on the label. Dense foliage or excessive growth will often prevent adequate coverage; adjust spray volumes accordingly. ]<sup>d</sup>

[ ELEVATE™ 50 WDG Fungicide is a hydroxylanilide protectant fungicide used for the control of Gray mold/Fruit rot (*Botrytis cinerea*) and Brown rot (*Monilinia* spp.). ]<sup>j</sup>

## RESISTANCE MANAGEMENT

ELEVATE 50 WDG is a group 17 fungicide. While there is no indication of cross-resistance between the active ingredient in this product and other fungicides, no more than two consecutive applications may be made with ELEVATE 50 WDG or any product containing its active ingredient, fenhexamid. After the second application, use an alternative fungicide effective in controlling the labeled diseases for two applications before reapplying the active ingredient in this product. Consult your local crop advisor for appropriate rotation recommendations. Resistance management strategies such as blocking program, alternation or tank mixture with other fungicides used for Botrytis and Monilinia control are recommended. To avoid the development of resistant strains, DO NOT apply ELEVATE 50 WDG at rates lower than specified on this label.



90628

[Alternate language appear in brackets]  
[Individual Sub-label language denoted by e, d, or j]

## COMPATIBILITY OF MIXTURES

This product is believed to be compatible with most commonly used fungicides, insecticides, micronutrients, growth regulators and spray adjuvants. Consult specific product labels for additional information. It is always advisable to conduct a tank spray compatibility test before mixing this product with other products. To determine the physical compatibility of this product with other products, use a jar test. [Using a quart jar, add the proportionate amounts of the products to approximately one quart of water with agitation. Add dry formulations first, then flowables, then emulsifiable concentrates last. After thorough mixing, allow this mixture to stand for 5 minutes. If the combination remains mixed or can be readily remixed, it is physically compatible]. Once compatibility has been proven, use the same procedure for adding required ingredients to the spray tank. It is suggested that combinations be used on a small number of plants before treating large areas.

## [PREPARATION OF SPRAY SOLUTION

**Read COMPATIBILITY OF MIXTURES section prior to any tank mixing with this product.** Begin with clean equipment. Add sufficient clean water to the spray tank for one-half of the mix load. Start tank agitation. Add and properly suspend the necessary amount of product before adding any other products. Agitate to ensure thorough mixing while adding the remaining required water. A high quality wetting agent or other spray adjuvant, approved for use on your crop, may be added to spray solutions according to the manufacturer's use instructions. ELEVATE 50 WDG should be properly suspended and diluted prior to the addition of any adjuvant. When an adjuvant is to be used with this product, Arysta recommends the use of a Chemical Producers and Distributors Association (CPDA) certified adjuvant. Consult the adjuvant label or manufacturer for crop tolerance and safety information when used with this product. Maintain agitation during mixing. DO NOT allow the spray mixture to stand without agitation. Apply with properly calibrated spray equipment. ]e, d

## APPLICATION

Prepare only the amount of spray solution needed to treat the area to be sprayed. Maintain agitation during application. Applications should be made immediately after the spray solution is prepared. If there is any delay in application, continue mix agitation. DO NOT allow the spray mixture to stand without agitation. If agitation is stopped for any reason, the spray solution must be thoroughly re-mixed prior to any further use. [Rainfall or overhead irrigation within 2 hours of application may necessitate re-treatment.]e, d

[Proper calibration and operation of spray equipment is essential for successful spraying. Consult the spray equipment manufacturer's operators manual, spray nozzle catalogs, and/or your crop advisor for more information.]e, d

[DO NOT apply this product through any type of irrigation system.] e, d

[ For agricultural applications, apply using ground equipment only (such as boom type, air blast, hand gun or mist blower sprayer). For food crops, thorough coverage of all flowers and fruiting parts is essential for control of blossom and fruit infections caused by *Botrytis* and *Monilinia* spp. ]e

[Alternate language appear in brackets]  
[Individual Sub-label language denoted by e, d, or j]

[ For agricultural applications, apply using ground equipment only (such as boom type, air blast, hand gun or mist blower sprayer). For food crops, thorough coverage of all flowers and fruiting parts is essential for control of blossom and fruit infections caused by *Botrytis*. ]<sub>d</sub>

[ For ornamental crops, thorough coverage is essential for disease control. Apply in sufficient water to obtain complete coverage of flowers, foliage and stems. Applications may be made with high volume, low volume or ultra low volume (non-thermal foggers, misters, etc.) ground equipment only. Follow the spray equipment manufacturer's directions to determine the amount of spray solution required to obtain thorough coverage. Consult the spray equipment manufacturer's operator manual, spray nozzle catalogs and/or your crop advisor for more information. DO NOT apply this product using thermal application equipment. ]<sub>d</sub>

[Do not plant food crops in the treated field other than those for which the product is labeled within 30 days following the last application.]<sub>e,d</sub>

<sub>e</sub>

ALMONDS			
Disease	Application Rate		Application Timing
Blossom blight, Twig blight ( <i>Monilinia</i> spp.)	Stand-alone	1.5 lbs/A (0.75 lb AI/A)	<ul style="list-style-type: none"><li>• Pink bud.</li><li>• Full bloom.</li><li>• Petal fall.</li><li>• Up to 28 days after petal fall.</li><li>• Apply in a minimum of 20 gallons of water by air and 50 gallons by ground.</li></ul>
	Tank-mix	1.0 – 1.5 lbs/A (0.5 – 0.75 lb AI/A)	
<b><u>Restrictions</u></b> <ul style="list-style-type: none"><li>• Apply in sufficient water to obtain complete coverage of flowers, fruit and twigs.</li><li>• Do not make more than 2 consecutive applications. Refer to the Resistance Management section for rotation guidelines.</li><li>• Under conditions conducive to low or moderate disease pressure only, ELEVATE 50 WDG may be applied in a tank-mix at a rate of 1.0 lb (0.5 lb AI) per acre with the specified rate of another fungicide which is registered for use on almonds to control <i>Monilinia</i> spp.</li><li>• Do not apply more than 6.0 lbs of ELEVATE 50 WDG (3.0 lbs AI) per acre per season.</li></ul>			

[Alternate language appear in brackets]  
[Individual Sub-label language denoted by e, d, or j]

e

<b>ASPARAGUS</b>		
<b>Disease</b>	<b>Application Rate</b>	<b>Application Timing</b>
Gray mold ( <i>Botrytis cinerea</i> )	1.5 lbs/A (0.75 lb AI/A)	<ul style="list-style-type: none"> <li>• Apply at Fern stage only. The ferns must be mowed down or allowed to senesce between the last application and harvest of the asparagus spears.</li> <li>• Make up to 4 foliar applications.</li> <li>• Make uniform applications of ELEVATE 50 WDG in a minimum of 40 gallons of water per acre.</li> <li>• Repeat at 7–14 day intervals if conditions favor disease development.</li> <li>• Apply in a minimum of 50 gallons of water by ground.</li> </ul>
<b><u>Restrictions</u></b> <ul style="list-style-type: none"> <li>• PHI for all states excluding California: Do not harvest the spears within 180 days of the last application.</li> <li>• PHI for California: Do not harvest the spears within 90 days of the last application.</li> <li>• Do not make more than 2 consecutive applications. Refer to the Resistance Management section for rotation guidelines.</li> <li>• Do not apply more than 6.0 lbs of ELEVATE 50 WDG (3.0 lbs AI) per acre per season.</li> </ul>		

e

<b>BUSHBERRIES</b> <b>(blueberries, gooseberries, currants and huckleberries)</b> <b>JUNEBERRIES, LINGONBERRIES and SALAL</b>		
<b>Disease</b>	<b>Application Rate</b>	<b>Application Timing</b>
Gray mold ( <i>Botrytis cinerea</i> )	1.5 lbs/A (0.75 lb AI/A)	<ul style="list-style-type: none"> <li>• Begin applications at 10% bloom.</li> <li>• Continue through harvest.</li> <li>• Applications should be made every 7 days or when conditions favor disease development.</li> <li>• Apply in a minimum of 50 gallons of water by ground.</li> </ul>
<b><u>Restrictions</u></b> <ul style="list-style-type: none"> <li>• PHI = 0 day.</li> <li>• Do not make more than 2 consecutive applications. Refer to the Resistance Management section for rotation guidelines.</li> <li>• Do not apply more than 6.0 lbs of ELEVATE 50 WDG (3.0 lbs AI) per acre per season.</li> </ul>		

[Alternate language appear in brackets]  
[Individual Sub-label language denoted by e, d, or j]

e

<b>CANE BERRIES</b> (red and black raspberries, blackberries and loganberries )		
<b>Disease</b>	<b>Application Rate</b>	<b>Application Timing</b>
Gray mold ( <i>Botrytis cinerea</i> )	1.5 lbs/A (0.75 lb AI/A)	<ul style="list-style-type: none"> <li>• Begin at 10% bloom.</li> <li>• Continue through harvest.</li> <li>• Applications should be made every 7 days or when conditions favor disease development.</li> <li>• Apply in a minimum of 50 gallons of water by ground.</li> </ul>
<b><u>Restrictions</u></b> <ul style="list-style-type: none"> <li>• PHI = 0 day.</li> <li>• Do not make more than 2 consecutive applications. Refer to the Resistance Management section for rotation guidelines.</li> <li>• Do not apply more than 6.0 lbs of ELEVATE 50 WDG (3.0 lb AI) per acre per season.</li> </ul>		

e

<b>GINSENG (EXCEPT CALIFORNIA)</b>		
<b>Disease</b>	<b>Application Rate</b>	<b>Application Timing</b>
Botrytis leaf and fruit blight ( <i>Botrytis cinerea</i> )	1.5 lb/A (0.75 lb AI/A)	<ul style="list-style-type: none"> <li>• Begin applications when conditions favor disease development.</li> <li>• Repeat at 7-14 day intervals if conditions continue to favor disease development (especially cool and wet weather or high humidity).</li> <li>• Make up to 4 foliar applications.</li> <li>• Make a uniform application in a minimum of 50 to 150 gallons of water per acre.</li> <li>• May be applied up to day of harvest however, wait for the fungicide solution to dry on the foliage prior to harvest.</li> </ul>
<b><u>Restrictions</u></b> <ul style="list-style-type: none"> <li>• PHI = 0 day.</li> <li>• Do not make more than 2 consecutive applications. Refer to the Resistance Management section for rotation guidelines.</li> <li>• Do not apply more than 6.0 lbs of ELEVATE 50 WDG (3.0 lbs AI) per acre per season.</li> </ul>		

[Alternate language appear in brackets]  
[Individual Sub-label language denoted by e, d, or j]

e

GRAPES		
Disease	Application Rate	Application Timing
Botrytis bunch rot ( <i>Botrytis cinerea</i> )	1.0 lb/A (0.5 lb AI/A)	<ul style="list-style-type: none"><li>• Early bloom.</li><li>• Bunch pre-closure.</li><li>• Veraison.</li><li>• Up to day of harvest.</li><li>• Apply in a minimum of 50 gallons of water by ground.</li></ul>
Powdery mildew suppression ( <i>Uncinula necator</i> )		
<b><u>Restrictions</u></b> <ul style="list-style-type: none"><li>• PHI = 0 day.</li><li>• ELEVATE 50 WDG has been shown to enhance the efficacy of other fungicides when used in a powdery mildew spray program.</li><li>• Do not make more than 2 consecutive applications. Refer to the Resistance Management section for rotation guidelines.</li><li>• Do not apply more than 3.0 lbs of ELEVATE 50 WDG (1.5 lbs AI) per acre per season.</li></ul>		

e

PEAR and ORIENTAL PEAR		
Disease	Application Rate	Application Timing
Fruit Rot ( <i>Botrytis cinerea</i> )	1.0 to 1.5 lbs/A (0.5 to 0.75 lb AI/A)	<ul style="list-style-type: none"> <li>• Make first application within 14 days of harvest.</li> <li>• Apply in a 7-day interval up to the day of harvest.</li> <li>• Apply in a minimum of 20 gallons of water by air and 50 gallons by ground.</li> </ul>
<b>Restrictions</b> <ul style="list-style-type: none"> <li>• PHI = 0 day.</li> <li>• Do not make more than 2 consecutive applications. If multiple harvests are planned on the same variety (i.e., color picking or selective harvesting) over an extended period, rotate between different classes of registered fungicides. Refer to the Resistance Management section for rotation guidelines.</li> <li>• Do not apply more than 6.0 lbs of ELEVATE 50 WDG (3.0 lb AI) per acre per season.</li> </ul>		

e

14 of 28

[Alternate language appear in brackets]  
 [Individual Sub-label language denoted by e, d, or j]

PISTACHIOS		
Disease	Application Rate	Application Timing
Botrytis blossom and shoot blight ( <i>Botrytis cinerea</i> )	1.0 - 1.5 lbs/A (0.5 – 0.75 lb AI/A)	<ul style="list-style-type: none"> <li>• 5-10% bloom.</li> <li>• Full bloom.</li> <li>• Then every 7-10 days or when conditions favor disease development.</li> <li>• Up to 28 days after petal fall.</li> <li>• Apply in a minimum of 20 gallons of water by air and 50 gallons by ground.</li> </ul>
<b><u>Restrictions</u></b> <ul style="list-style-type: none"> <li>• Apply in sufficient water to obtain complete coverage of flowers, fruit and twigs.</li> <li>• Do not make more than 2 consecutive applications. Refer to the Resistance Management section for rotation guidelines.</li> <li>• Under conditions conducive to low or moderate disease pressure only, ELEVATE 50 WDG may be applied at a rate of 1.0 lb (0.5 lb AI) per acre.</li> <li>• Do not apply more than 6.0 lbs of ELEVATE 50 WDG (3.0 lbs AI/) per acre per season.</li> </ul>		

[Alternate language appear in brackets]  
[Individual Sub-label language denoted by e, d, or j]

e

STONE FRUIT (apricot, cherries, nectarines, peaches, plumcot, plums and fresh prunes)			
Field Application			
Disease	Application Rate		Application Timing
Brown rot, Blossom blight, Twig blight ( <i>Monilinia</i> spp.)	Stand-alone	1.5 lbs/A (0.75 lb AI/A)	<ul style="list-style-type: none"> <li>• Early bloom.</li> <li>• Late bloom.</li> <li>• 2 weeks after petal fall.</li> <li>• Up to day of harvest.</li> <li>• Apply in a minimum of 20 gallons of water by air and 50 gallons by ground.</li> </ul>
	Tank-mix	1.0-1.5 lbs/A (0.5 - 0.75 lb AI/A)	
Gray mold on cherries ( <i>Botrytis cinerea</i> )		1.0 - 1.5 lbs/A (0.5 - 0.75 lb AI/A)	<ul style="list-style-type: none"> <li>• Early bloom.</li> <li>• Then every 7-10 days or when conditions favor disease development.</li> <li>• Up to day of harvest.</li> <li>• Apply in a minimum of 20 gallons of water by air and 50 gallons by ground.</li> </ul>
<b>Restrictions</b> <ul style="list-style-type: none"> <li>• PHI = 0 day.</li> <li>• Do not make more than 2 consecutive applications. Refer to the Resistance Management section for rotation guidelines.</li> <li>• Under conditions conducive to low or moderate <i>Monilinia</i> disease pressure only, ELEVATE 50 WDG may be applied in a tank-mix at a rate of 1.0 lb (0.5 lb AI) per acre with the specified rate of another fungicide which is registered for use on stone fruit to control <i>Monilinia</i> spp.</li> <li>• Under conditions conducive to low or moderate <i>Botrytis</i> disease pressure only, ELEVATE 50 WDG may be applied at a rate of 1.0 lb (0.5 lb AI) per acre to control <i>Botrytis cinerea</i> on cherries only.</li> <li>• Do not apply more than a maximum of 6.0 lbs of ELEVATE 50 WDG (3.0 lbs AI) per acre per season.</li> </ul>			

16 of 28

[Alternate language appear in brackets]  
 [Individual Sub-label language denoted by e, d, or j]

e

STRAWBERRIES			
Disease	Application Rate		Application Timing
Gray mold ( <i>Botrytis cinerea</i> )	Stand-alone	1.5 lbs/A (0.75 lb AI/A)	<ul style="list-style-type: none"><li>• Begin applications at early bloom.</li><li>• Repeat at 7-14 day intervals if conditions continue to favor disease development.</li><li>• May be applied up to and including the day of harvest.</li><li>• Apply in a minimum of 50 gallons of water by ground.</li></ul>
	Tank-mix	1.0-1.5 lbs/A (0.5-0.75 lb AI/A)	

**Restrictions**

- PHI = 0 day.
- Under conditions conducive to low or moderate disease pressure only, ELEVATE 50 WDG may be applied in a tank-mix at a rate of 1.0 lb (0.5 lb AI) per acre with the specified rate of another fungicide which is registered for use on strawberries to control *Botrytis cinerea*.
- Do not make more than 2 consecutive applications. Refer to the Resistance Management section for rotation guidelines.
- Do not apply more than 6.0 lbs of ELEVATE 50 WDG (3.0 lbs AI) per acre per season.
- When plastic mulch is used, do not apply within 16 feet (5 meters) of naturally vegetated or aquatic areas.



170828

[Alternate language appear in brackets]  
[Individual Sub-label language denoted by e, d, or j]

d

CILANTRO			
Transplant Production and Greenhouse Production (EXCEPT CALIFORNIA)			
Transplant Production			
Disease	Application Rate		Application Timing
Gray mold ( <i>Botrytis cinerea</i> )	Stand-alone	1.5 lbs/A (0.75 lb AI/A)	<ul style="list-style-type: none"><li>• Begin applications when conditions favor disease development.</li><li>• Repeat application after 7 days if conditions favor disease development.</li></ul>
	Tank-mix	1.0 - 1.5 lbs/A (0.5 - 0.75 lb AI/A)	
Greenhouse Production			
Disease	Application Rate		Application Timing
Gray mold ( <i>Botrytis cinerea</i> )	Stand-alone	1.5 lbs/A (0.75 lb AI/A)	<ul style="list-style-type: none"><li>• Begin applications when conditions favor disease development.</li><li>• Repeat application after 7 days if conditions favor disease development.</li><li>• Do not apply within 3 days of harvest.</li></ul>
	Tank-mix	1.0 - 1.5 lbs/A (0.5 - 0.75 lb AI/A)	
<b><u>Restrictions</u></b>			
<ul style="list-style-type: none"><li>• Do not apply in field.</li><li>• Do not make more than 2 consecutive applications. Refer to the Resistance Management section for rotation guidelines.</li><li>• Under conditions conducive to low or moderate disease pressure only, ELEVATE 50 WDG may be applied in a tank-mix at a rate of 1.0 lb (0.5 lb AI) per acre with the recommended rate of another fungicide which is registered for use on cilantro to control <i>Botrytis cinerea</i>.</li></ul>			
<b><u>Transplant Production</u></b>			
<ul style="list-style-type: none"><li>• Apply in a full coverage protective spray.</li><li>• Do not apply more than 3.0 lbs of ELEVATE 50 WDG (1.5 lb AI) per acre to transplants.</li></ul>			
<b><u>Greenhouse Production</u></b>			
<ul style="list-style-type: none"><li>• PHI = 3 days.</li><li>• Do not apply on transplanted cilantro prior to bloom.</li><li>• Do not make more than 3.0 lbs of ELEVATE 50 WDG (1.5 lb AI) per acre per crop.</li></ul>			

18 of 28

[Alternate language appear in brackets]  
[Individual Sub-label language denoted by e, d, or j]

d

CUCUMBER			
Transplant and Greenhouse Production (EXCEPT CALIFORNIA)			
Transplant Production			
Disease	Application Rate		Application Timing
Gray mold ( <i>Botrytis cinerea</i> )	Stand-alone	1.5 lbs/A (0.75 lb AI/A)	<ul style="list-style-type: none"><li>• Begin applications when conditions favor disease development.</li><li>• Repeat at 7-10 day intervals if conditions continue to favor disease.</li></ul>
	Tank-mixed	1.0 - 1.5 lbs/A (0.5 - 0.75 lb AI/A)	
Greenhouse Production			
Disease	Application Rate		Application Timing
Gray mold ( <i>Botrytis cinerea</i> )	Stand-alone	1.5 lbs/A (0.75 lb AI/A)	<ul style="list-style-type: none"><li>• Begin applications when conditions favor disease development.</li><li>• Repeat at 7-10 day intervals if conditions continue to favor disease.</li><li>• May be applied up to and including the day of harvest.</li></ul>
	Tank-mixed	1.0 - 1.5 lbs/A (0.5 - 0.75 lb AI/A)	
<b><u>Restrictions</u></b>			
<ul style="list-style-type: none"><li>• Do not apply in field.</li><li>• Do not make more than 2 consecutive applications. Refer to the Resistance Management section for rotation guidelines.</li><li>• Under conditions conducive to low or moderate disease pressure only, ELEVATE 50 WDG may be applied in a tank-mix at a rate of 1.0 lb (0.5 lb AI) per acre with the recommended rate of another fungicide which is registered for use on cucumber to control <i>Botrytis cinerea</i>.</li></ul>			
<b><u>Transplant Production</u></b>			
<ul style="list-style-type: none"><li>• Apply in a full coverage protective spray.</li><li>• Do not apply more than 6.0 lbs of ELEVATE 50 WDG (3.0 lb AI) per acre to transplants.</li></ul>			
<b><u>Greenhouse Production</u></b>			
<ul style="list-style-type: none"><li>• PHI = 0 day.</li><li>• Do not apply more than 6.0 lbs of ELEVATE 50 WDG (3.0 lb AI) per acre per season.</li></ul>			

[Alternate language appear in brackets]  
[Individual Sub-label language denoted by e, d, or j]

d

FRUITING VEGETABLES (except Tomatoes) (EGGPLANT, GROUND CHERRY, PEPINO, PEPPER [Bell Pepper, Chili Pepper, Cooking Pepper, Pimento, Sweet Pepper], and TOMATILLO) Transplant and Greenhouse Production (EXCEPT CALIFORNIA)			
Transplant Production			
Disease	Application Rate		Application Timing
Gray mold ( <i>Botrytis cinerea</i> )	Stand-alone	1.5 lbs/A (0.75 lb AI/A)	<ul style="list-style-type: none"><li>• Begin applications when conditions favor disease development.</li><li>• Repeat at 7-10 day intervals if conditions continue to favor disease.</li></ul>
	Tank-mixed	1.0 - 1.5 lbs/A (0.5 - 0.75 lb AI/A)	
Greenhouse Production			
Disease	Application Rate		Application Timing
Gray mold ( <i>Botrytis cinerea</i> )	Stand-alone	1.5 lbs/A (0.75 lb AI/A)	<ul style="list-style-type: none"><li>• Begin applications when conditions favor disease development.</li><li>• Repeat at 7-10 day intervals if conditions continue to favor disease.</li><li>• May be applied up to and including the day of harvest.</li></ul>
	Tank-mixed	1.0 - 1.5 lbs/A (0.5 - 0.75 lb AI/A)	
<b><u>Restrictions</u></b> <ul style="list-style-type: none"><li>• Do not apply in field.</li><li>• Do not make more than 2 consecutive applications. Refer to the Resistance Management section for rotation guidelines.</li><li>• Under conditions conducive to low or moderate disease pressure only, ELEVATE 50 WDG may be applied in a tank-mix at a rate of 1.0 lb (0.5 lb AI) per acre with the recommended rate of another fungicide which is registered for use on fruiting vegetables to control <i>Botrytis cinerea</i>.</li></ul>			
<b><u>Transplant Production</u></b> <ul style="list-style-type: none"><li>• Apply in a full coverage protective spray.</li><li>• Do not apply more than 6.0 lbs of ELEVATE 50 WDG (3.0 lb AI) per acre to transplants.</li></ul>			
<b><u>Greenhouse Production</u></b> <ul style="list-style-type: none"><li>• PHI = 0 day.</li><li>• Do not apply more than 6.0 lbs of ELEVATE 50 WDG (3.0 lb AI) per acre per season.</li></ul>			

[Alternate language appear in brackets]  
[Individual Sub-label language denoted by e, d, or j]

d

TOMATOES			
Transplant and Greenhouse Production			
Transplant Production			
Disease	Application Rate		Application Timing
Gray mold ( <i>Botrytis cinerea</i> )	Stand-alone	1.5 lbs/A (0.75 lb AI/A)	<ul style="list-style-type: none"><li>• Begin applications when conditions favor disease development.</li><li>• Repeat at 7-10 day intervals if conditions continue to favor disease.</li></ul>
	Tank-mixed	1.0 - 1.5 lbs/A (0.5 - 0.75 lb AI/A)	
Greenhouse Production			
Disease	Application Rate		Application Timing
Gray mold ( <i>Botrytis cinerea</i> )	Stand-alone	1.5 lbs/A (0.75 lb AI/A)	<ul style="list-style-type: none"><li>• Begin applications when conditions favor disease development.</li><li>• Repeat at 7-10 day intervals if conditions continue to favor disease.</li><li>• May be applied up to and including the day of harvest.</li></ul>
	Tank-mixed	1.0 - 1.5 lbs/A (0.5 - 0.75 lb AI/A)	
<b><u>Restrictions</u></b>			
<ul style="list-style-type: none"><li>• Do not apply in field.</li><li>• Do not make more than 2 consecutive applications. Refer to the Resistance Management section for rotation guidelines.</li><li>• Under conditions conducive to low or moderate disease pressure only, ELEVATE 50 WDG may be applied in a tank-mix at a rate of 1.0 lb (0.5 lb AI) per acre with the recommended rate of another fungicide which is registered for use on fruiting vegetables to control <i>Botrytis cinerea</i>.</li></ul>			
<b><u>Transplant Production</u></b>			
<ul style="list-style-type: none"><li>• Apply in a full coverage protective spray.</li><li>• Do not apply more than 6.0 lbs of ELEVATE 50 WDG (3.0 lb AI) per acre to transplants.</li></ul>			
<b><u>Greenhouse Production</u></b>			
<ul style="list-style-type: none"><li>• PHI = 0 day.</li><li>• Do not apply more than 6.0 lbs of ELEVATE 50 WDG (3.0 lb AI) per acre per season.</li></ul>			

[Alternate language appear in brackets]  
[Individual Sub-label language denoted by e, d, or j]

d

LEAFY GREENS (EXCEPT SPINACH)			
(Amaranth; arugula; chervil; chrysanthemum, edible-leaved and garland; corn salad; cress, garden and upland; dandelion; dock; endive; lettuce; orach; parsley; purslane, garden and winter; radicchio [red chicory])			
Transplant and Greenhouse Production (EXCEPT CALIFORNIA)			
Transplant Production			
Disease	Application Rate		Application Timing
Gray mold ( <i>Botrytis cinerea</i> )	Stand-alone	1.5 lbs/A (0.75 lb AI/A)	<ul style="list-style-type: none"><li>• Begin applications when conditions favor disease development.</li><li>• Repeat application after 7 days if conditions continue to favor disease.</li></ul>
	Tank-mix	1.0 - 1.5 lbs/A (0.5 - 0.75 lb AI/A)	
Greenhouse Production			
Disease	Application Rate		Application Timing
Gray mold ( <i>Botrytis cinerea</i> )	Stand-alone	1.5 lbs/A (0.75 lb AI/A)	<ul style="list-style-type: none"><li>• Begin applications when conditions favor disease development.</li><li>• Repeat application after 7 days if conditions continue to favor disease.</li><li>• Do not apply within 3 days of harvest.</li></ul>
	Tank-mix	1.0 - 1.5 lbs/A (0.5 - 0.75 lb AI/A)	
<b>Restrictions</b> <ul style="list-style-type: none"><li>• Do not apply in field.</li><li>• Do not make more than 2 consecutive applications. Refer to the Resistance Management section for rotation guidelines.</li><li>• Under conditions conducive to low or moderate disease pressure only, ELEVATE 50 WDG may be applied in a tank-mix at a rate of 1.0 lb (0.5 lb AI) per acre with the recommended rate of another fungicide which is registered for use on leafy greens to control <i>Botrytis cinerea</i>.</li></ul>			
<b>Transplant Production</b> <ul style="list-style-type: none"><li>• Apply in a full coverage protective spray.</li><li>• Do not apply more than 3.0 lbs of ELEVATE 50 WDG (1.5 lb AI) per acre to transplants.</li></ul>			
<b>Greenhouse Production</b> <ul style="list-style-type: none"><li>• PHI = 3 days.</li><li>• Do not apply more than 3.0 lbs of ELEVATE 50 WDG (1.5 lb AI) per acre per crop.</li></ul>			

[Alternate language appear in brackets]  
[Individual Sub-label language denoted by e, d, or j]

d

# **OUTDOOR-, GREENHOUSE- AND NURSERY-GROWN ORNAMENTALS, FORESTRY CONIFERS AND NON-BEARING FRUIT TREES AND VINES**

This product, when applied as a foliar spray, has been shown to effectively protect plants from *Botrytis cinerea*, *Botrytis elliptica* and *Botrytis tulipae* and stop sporulation of the fungus if an infection has occurred. Also, when this product is used to control Botrytis, suppression of powdery mildew on the treated crop may be achieved. These directions are based upon the results of product testing programs.

This product has been shown to be safe and effective in a limited number of research trials on certain varieties or cultivars of the plant types listed. Users should conduct small scale tests under local growing conditions prior to wide-scale use.

African Violet	Iris	Rose
Crabapple, Flowering <sup>1</sup>	Lily (Asiatic/Oriental)	Spathiphyllum
Daffodil	Lisianthus	Snapdragon
Douglas Fir	Pansy	Tulip
Fuchsia	Petunia	Western Hemlock
Geranium	Plum, Flowering <sup>1</sup>	
Gerbera Daisy	Poinsettia <sup>2</sup>	
Impatiens	Ranunculus	

<sup>1</sup> Not for food or feed purposes.

<sup>2</sup> Due to the sensitivity of mature Poinsettia bracts and varying local environmental conditions, users should conduct preliminary tests as suggested above.

d

High Volume Sprays		
Application Rate	Disease Pressure	Spray Interval
0.75 lb product per 100 gal.	Conditions favor disease development; no disease present.	7 to 14 Days
1.0 to 1.5 lb product per 100 gal.	Conditions favor disease development; disease present.	7 Days

d

Low Volume / Ultra Low Volume Sprays		
Application Rate	Disease Pressure	Spray Interval
0.75 lb product per 100 gal.	Conditions favor disease development; no disease present.	7 to 14 Days
1.0 to 1.5 lb product per 100 gal.	Conditions favor disease development; disease present.	7 Days

For small application equipment, the scoop attached to the container is provided for ease in measuring the amount of this product to use per 1 gallon of water. Apply 1 to 2 level scoops per 1 gallon of water. One scoop equals 0.12 ounces or 3.4 grams or approximately one rounded teaspoon of this product.

[Alternate language appear in brackets]  
 [Individual Sub-label language denoted by e, d, or j]

d

### POST-HARVEST APPLICATIONS

For use on cut flowers, rose budwood, and bare-root nursery stock when applied as a spray. Apply ELEVATE 50 WDG to prevent *Botrytis* diseases (Gray mold) in cold storage or transit. Apply after harvest and grading prior to cold storage. Apply 1.0 - 1.5 pounds of ELEVATE 50 WDG per 100 gallons of spray.

d

### LIMITATIONS

Do not make more than two consecutive applications of this product. After the second application, use an alternative fungicide effective in controlling *Botrytis* spp. for two applications before reapplying the active ingredient in this product. Consult your local Horticultural Advisor for the most appropriate alternative products.

d

### Ornamentals and Other Nursery Crops Grown Outdoors

DO NOT apply more than 6.0 pounds of product per acre per season (3.0 lb AI/A/season).

### Ornamentals and Other Nursery Crops Grown in Greenhouses

DO NOT apply more than 0.14 pounds product per 1000 square feet per crop cycle (equal to 2.2 oz. product per 1000 square feet per crop cycle).

KIWI Post-Harvest Applications			
Disease	Application Type	Application Rate	Application Timing
Fruit Rot ( <i>Botrytis cinerea</i> )	Packing-line spray	1.5 lbs (0.75 lb AI)/200,000 lbs fruit in 8–20 gallons of water.	Post-harvest
	Fruit dip treatment	1.5 lbs (0.75 lb ai) in 100 gallons of water. Fruit should remain in dip solution for 20 to 30 seconds.	
<b><u>Post-Harvest Restrictions:</u></b>			
<ul style="list-style-type: none"><li>• Do not make more than one application to fruit.</li><li>• Use only as a post harvest application.</li><li>• Leftover dip or spray mixtures containing ELEVATE 50 WDG may be used as a foliar spray on other registered crops, observing all restrictions such as maximum pounds applied per application per season.</li><li>• When calculating application rates, if analytical services are not available to determine the exact quantity of ELEVATE 50 WDG remaining in the mixture, assume that the tank still contains the initial concentration of ELEVATE 50 WDG per gallon (1.5 lbs per volume).</li><li>• If the dip or spray mixture contains other pesticides in addition to ELEVATE 50 WDG, refer to the product label(s) for information regarding disposal. Apply only on a crop for which all products are registered and follow all restrictions such as maximum pounds applied per application and season.</li></ul>			

240728

[Alternate language appear in brackets]  
 [Individual Sub-label language denoted by e, d, or j]

PEAR and ORIENTAL PEAR Post-Harvest Applications			
Disease	Application Type	Application Rate	Application Timing
Fruit Rot ( <i>Botrytis cinerea</i> )	Low-Volume Packing-line spray	1.0 to 1.5 lbs (0.75 lb AI) per 200,000 lbs fruit in 8–20 gallons of an appropriate water, wax/oil emulsion, or aqueous dilution of a wax/oil emulsion.	Post-harvest
	High-Volume Packing-line spray	1.0 to 1.5 lbs (0.75 lb AI) per 200,000 lbs fruit in 100 gallons of an appropriate water, wax/oil emulsion, or aqueous dilution of a wax/oil emulsion.	
	Fruit dip treatment	1.0 to 1.5 lbs (0.75 lb AI) per 200,000 lbs fruit in 100 gallons of an appropriate water, wax/oil emulsion, or aqueous dilution of a wax/oil emulsion. Fruit should remain in dip solution for 20 to 30 seconds.	
<b><u>Post-Harvest Restrictions:</u></b> <ul style="list-style-type: none"><li>• Apply only as post-harvest spray to fruit in a packing line or dip fruit in the solution for 30 seconds.</li><li>• Do not make more than one post-harvest application.</li><li>• For low volume (concentrated) application use a control droplet-type application (CDA) system or similar low volume system.</li><li>• Do not apply more than 1.5 lbs of ELEVATE 50 WDG (0.75 lb AI) as a post-harvest treatment.</li><li>• Leftover dip or spray mixtures containing ELEVATE 50 WDG may be used as a foliar spray on other registered crops, observing all restrictions such as maximum pounds applied per application per season.</li><li>• When calculating application rates, if analytical services are not available to determine the exact quantity of ELEVATE 50 WDG remaining in the mixture, assume that the tank still contains the initial concentration of ELEVATE 50 WDG per gallon (1.5 lb per volume).</li><li>• If the dip or spray mixture contains other pesticides in addition to ELEVATE 50 WDG, refer to the product label(s) for information regarding disposal. Apply only on a crop for which all products are registered and follow all restrictions such as maximum pounds applied per application per season.</li></ul>			



[Alternate language appear in brackets]  
 [Individual Sub-label language denoted by e, d, or j]

POMEGRANATE Post-Harvest Applications			
Disease	Application Type	Application Rate	Application Timing
Botrytis Fruit Rot	Fruit dip treatment	1.5 lbs (0.75 lb ai) in 100 gallons of water. Fruit should remain in dip solution for 20 to 30 seconds.	Post harvest
<b>Post-Harvest Restrictions:</b> <ul style="list-style-type: none"> <li>Do not make more than one application to fruit.</li> <li>Use only as a post harvest application.</li> <li>Leftover dip containing ELEVATE 50 WDG may be used as a foliar spray on other registered crops, observing all restrictions such as maximum pounds applied per application per season.</li> <li>When calculating application rates, if analytical services are not available to determine the exact quantity of ELEVATE 50 WDG remaining in the mixture, assume that the tank still contains the initial concentration of ELEVATE 50 WDG per gallon (1.5 lbs per 100 gallons).</li> <li>If the dip contains other pesticides in addition to ELEVATE 50 WDG, refer to the product label(s) for information regarding disposal. Apply only on a crop for which all products are registered and follow all restrictions such as maximum pounds applied per application and season.</li> </ul>			

STONE FRUIT except Cherries ( <u>apricots, nectarines, peaches, plumcot, plums and fresh prunes</u> ) Post-Harvest Applications			
Disease	Application Rate		Application Timing
Brown rot ( <i>Monilinia</i> spp.) on apricot, nectarines, peaches and plumcots	High volume Packing-line spray	1.5 lb (0.75 lb AI) per 100 gallons of water, wax/oil emulsion or aqueous dilution of a wax/oil emulsion to 200,000 lbs of fruit.	Post-harvest
	Low volume Packing-line spray	1.5 lb (0.75 lb AI) in 8 - 25 gallons of water, wax/oil emulsion or aqueous dilution of a wax/oil emulsion to 200,000 lbs of fruit.	

260f28

[Alternate language appear in brackets]  
 [Individual Sub-label language denoted by e, d, or j]

Post-Harvest Application – Cherries		
Disease	Application Rate	
Brown rot ( <i>Monilinia</i> spp.) and Gray mold ( <i>Botrytis cinerea</i> )	High-volume Packing-line spray	1.5 lb (0.75 lb ai) per 100 gallons of water, wax/oil emulsion or aqueous dilution of a wax/oil emulsion to 25,000 lb of fruit.
	Fruit dip treatment	1.5 lb (0.75 lb ai) per 100 gallons of water, wax/oil emulsion or aqueous dilution of a wax/oil emulsion to 25,000 lb of fruit. Fruit should remain in dip solution for 30 seconds.
<b>Post-Harvest Restrictions:</b> <ul style="list-style-type: none"> <li>• Apply only as post-harvest spray to fruit in a packing line or dip fruit in the solution for 30 seconds.</li> <li>• Do not make more than one post-harvest application.</li> <li>• For low volume (concentrated) application use a control droplet-type application (CDA) system or similar low volume system.</li> <li>• Do not apply more than 1.5 lbs of ELEVATE 50 WDG (0.75 lb AI) as a post-harvest treatment.</li> <li>• Leftover dip or spray mixtures containing ELEVATE 50 WDG may be used as a foliar spray on other registered crops, observing all restrictions such as maximum pounds applied per application per season.</li> <li>• When calculating application rates, if analytical services are not available to determine the exact quantity of ELEVATE 50 WDG remaining in the mixture, assume that the tank still contains the initial concentration of ELEVATE 50 WDG per gallon (1.5 lb per volume).</li> <li>• If the dip or spray mixture contains other pesticides in addition to ELEVATE 50 WDG, refer to the product label(s) for information regarding disposal. Apply only on a crop for which all products are registered and follow all restrictions such as maximum pounds applied per application per season.</li> </ul>		

[Alternate language appear in brackets]  
 [Individual Sub-label language denoted by e, d, or j]

### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**PESTICIDE STORAGE:** Store in a secure dry place away from excessive heat. Do not store near food or feed. Store in original container only. To close package, fold package or replace and tighten cap to form an airtight seal.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

#### CONTAINER DISPOSAL:

**Bags:** Non-refillable container. Do not reuse or refill this container. Completely empty bag into application equipment. Then dispose of bag in a sanitary landfill or by incineration, or (if allowed by State and local authorities) by burning. If burned, stay out of smoke. Offer for recycling if available.

**Plastic containers:** Non-refillable container. Do not reuse or refill this container. Offer for recycling, if available, or puncture and dispose of in a sanitary landfill, or (if allowed by State and local authorities) by burning. If burned stay out of smoke. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or (if allowed by State and local authorities) by burning. If burned, stay out of smoke.

**Fiber drums:** Refillable container. Refill this container with this pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. Completely empty liner by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Then dispose of liner in a sanitary landfill (or incinerate if allowed by State and local authorities). If drum is contaminated and cannot be reused, dispose of in the same manner. To clean the container before final disposal, empty any remaining contents from container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this procedure two more times.

[Alternate language appear in brackets]  
[Individual Sub-label language denoted by e, d, or j]

#### CONDITIONS OF SALE

1. Arysta LifeScience North America, ~~LLC~~ ~~Corporation~~ ("Arysta") warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in accordance with the Directions for Use under normal conditions of use.
2. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal use conditions, or under conditions not reasonably foreseeable to Arysta. **ARYSTA DISCLAIMS ALL OTHER WARRANTIES, EXPRESSED OR IMPLIED, INCLUDING ANY WARRANTY OF FITNESS FOR A PARTICULAR PURPOSE OR MERCHANTABILITY. TO THE EXTENT PERMITTED BY LAW, SELLER SHALL NOT BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM OR IN CONNECTION WITH THE MANUFACTURE, SALE, DELIVERY, USE, HANDLING OR STORAGE OF THIS PRODUCT. TO THE EXTENT PERMITTED BY LAW, SELLER'S LIABILITY SHALL BE LIMITED TO THE REFUND OF THE PURCHASE PRICE. ARYSTA DOES NOT AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE ANY OTHER WARRANTY, GUARANTEE OR REPRESENTATION CONCERNING THIS PRODUCT.**
3. Critical and unforeseeable factors beyond Arysta's control prevent Arysta from eliminating all risks in connection with the use of this product. Such risks include, but are not limited to, damage to plants and crops to which the product is applied, lack of complete control, and damage caused by drift to other plants or crops. Such risks occur even though the product is reasonably fit for the use stated on the label. Except as stated in 1 above, to the extent permitted by law, by purchasing, accepting and using this product, the buyer and user acknowledge and assume all risks and liabilities resulting from handling, storage, and use of this product.
4. For species and their varieties or cultivars not specifically listed on the label, the user assumes all risk from phytotoxicity or unacceptable residue or disease control.

Fenhexamid was invented and is produced by Bayer AG, Germany.  
ELEVATE is a trademark of Arysta LifeScience North America Corporation.