

**EVEREST®**  
**70% WATER DISPERSIBLE GRANULAR HERBICIDE**

**FOR PREPLANT, PREEMERGENCE AND POSTEMERGENCE CONTROL OF  
WILD OAT, GREEN FOXTAIL AND OTHER GRASS AND BROADLEAF WEEDS  
IN SPRING, DURUM AND WINTER WHEAT.**

Active Ingredient	By wt.
Flucarbazone-sodium*, 4,5-Dihydro-3-methoxy-4-methyl-5-oxo-N- [[2-(trifluoromethoxy)phenyl]sulfonyl]-1H- 1,2,4-triazole-1-carboxamide, sodium salt .....	70.0%
Inert Ingredients .....	30.0%
<b>Total</b> .....	<b>100.0%</b>

\*66% Flucarbazone acid equivalent

®EVEREST is a registered trademark of Arysta LifeScience North America Corporation.

Read entire label before use.

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

See back panel for additional precautionary statements.

**ARYSTA LIFESCIENCE NORTH AMERICA CORPORATION**

15401 Weston Parkway, Suite 150  
Cary, North Carolina 27513

EPA Registration No. 66330-49  
EPA Est. No.

NET WEIGHT \_\_\_\_ POUNDS ( \_\_\_\_ KILOGRAMS)

FIRST AID	
If on skin or clothing	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15 to 20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
Have an EVEREST container or label with you when calling a poison control center or doctor.	
<b>IN CASE OF EMERGENCY CALL DAY OR NIGHT: (800) 228-5635 EXT. 174.</b>	
<b>Note To Physician:</b> No specific antidote is available. Treat the patient symptomatically.	

**ACCEPTED**

5-22-06

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.

66330-49

## PRECAUTIONARY STATEMENTS

### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**CAUTION:** Harmful if absorbed through skin. Avoid contact with skin, eyes or clothing.

### PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves (Category A) made of materials such as butyl rubber  $\geq 14$  mils, natural rubber  $\geq 14$  mils, neoprene rubber  $\geq 14$  mils, or nitrile rubber  $\geq 14$  mils
- Shoes plus socks

Follow manufacturer's instructions for cleaning / maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

### ENGINEERING CONTROL STATEMENT

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR §170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

#### USER SAFETY RECOMMENDATIONS:

User should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

### ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water when disposing of equipment washwaters.

Do not allow sprays to drift onto adjacent desirable plants.

Important: Read these entire DIRECTIONS FOR USE and CONDITIONS OF SALE before using EVEREST.

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

#### **AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours following application.

Exception: If the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls, chemical-resistant gloves (Category A) made of materials such as butyl rubber  $\geq 14$  mils, natural rubber  $\geq 14$  mils, neoprene rubber  $\geq 14$  mils, or nitrile rubber  $\geq 14$  mils, shoes plus socks.

2. Add the appropriate rate of EVEREST directly to the spray tank.
3. Add the broadleaf weed herbicide.
4. Add the surfactant.
5. Add micronutrients (if needed).
6. Fill the spray tank to the required level.
7. Maintain sufficient agitation during both mixing and application of EVEREST

4/10

### GENERAL INFORMATION

EVEREST is a selective herbicide for the control of wild oat, green foxtail, Italian ryegrass, windgrass, cheat, Japanese brome and numerous broadleaf weeds, including redroot pigweed, wild mustard and shepherd's purse, in spring, durum and winter wheat. EVEREST also suppresses additional grass and broadleaf weeds, including yellow foxtail, downy brome and wild buckwheat.

EVEREST is absorbed by foliage and roots of susceptible weeds, which cease growth soon after application. Complete weed emergence is not necessary for control due to the soil residual activity provided by EVEREST. However, maximum weed control may not be seen for one to two weeks, though susceptible weeds will stop growing and will no longer be competitive. For broader spectrum activity, EVEREST may be tank mixed with a broadleaf herbicide listed on this label. See "TANK MIXES" section for recommended products.

EVEREST is an acetolactate synthase (ALS) inhibitor, and will therefore control weed biotypes which have developed target site resistance to certain classes of herbicides, including ACCase inhibitors, dinitroanilines and triallates. See "RESISTANCE MANAGEMENT" section for additional information.

Read the entire DIRECTIONS FOR USE before using EVEREST.

### GENERAL USE RESTRICTIONS

1. For use only in wheat. Treated wheat fields may be grazed at any time.
2. Do not mix, load or clean spray equipment within 33 feet of well-heads or aquatic systems, including marshes, ponds, ditches, streams, lakes, etc. Do not apply within 50 feet of well-heads or the above mentioned aquatic systems.
3. Do not apply postemergence when rain is expected within the next hour.
4. Do not allow this chemical to drift onto other crops.
5. Observe minimum interval to harvest of 60 days after treatment.
6. Do not apply this product through any type of irrigation system.
7. Do not use flood irrigation to apply or incorporate EVEREST.

### MIXING INSTRUCTIONS

Ensure the spray tank is clean. In-line strainers and nozzle screens should be clean and 50 mesh or coarser.

1. Fill the spray tank  $\frac{1}{4}$  to  $\frac{1}{2}$  full with clean water and begin agitation or bypass.
2. Add the appropriate rate of EVEREST directly to the spray tank.
3. Add the broadleaf weed herbicide.
4. Add the surfactant.
5. Add micronutrients (if needed).
6. Fill the spray tank to the required level.
7. Maintain sufficient agitation during both mixing and application of EVEREST.

5/10

**POSTEMERGENCE USE DIRECTIONS**  
**FOR SPRING, DURUM AND WINTER WHEAT**

**APPLICATION PROCEDURES**

Best weed control is observed when environmental conditions support vigorous growth of crop and weeds. Research has demonstrated that optimum wheat yield is obtained by early removal of grassy weeds. Apply EVEREST to spring wheat prior to jointing, when the majority of plants have from one leaf to a maximum of 4 leaves on the main stem plus two tillers. For winter wheat, apply either in the fall or spring when the majority of plants have one leaf to full tillering, but prior to jointing.

EVEREST should not be applied after jointing begins to avoid the risk of crop injury.

Do not apply more than 0.018 lbs. acid equivalent (a.e.)/acre flucarbazone (0.6 ounce/acre of EVEREST) per growing season.

Do not make more than one postemergence application of EVEREST per growing season.

**GROUND APPLICATION**

Apply in a spray volume of 5 to 10 gallons/acre (or 50 to 100 liters/hectare) at 30 to 50 psi to ensure proper weed coverage. Flat fan nozzles of 80 or 110 degrees are recommended for optimum coverage. Do not use floodjet or control droplet application equipment. Nozzles may be oriented 45 degrees forward to enhance crop penetration and to give better weed coverage.

**AERIAL APPLICATION**

Apply in water using a minimum spray volume of 3 gallons/acre (or 30 liters/hectare). For best results, use a minimum of 5 gallons/acre (or 50 liters/hectare) under dry conditions or heavy weed infestations. Use nozzles that provide 200 to 350 micron size droplets for best results and to insure uniform spray coverage. Aerial applications with EVEREST should be made with low drift nozzles at a maximum height of 10 feet above the crop and at a maximum pressure of 40 psi. Do not apply aerially when wind speed is greater than 10 mph. Do not allow spray to drift onto adjacent crops, as injury or loss may occur.

See the *Aerial Drift Reduction Advisory Information* section of this label for additional information on how to reduce drift during aerial application.

6/10

**ENDANGERED SPECIES PROTECTION**

To avoid adverse effects on endangered dicot plant species, the following measures will be required where endangered plant species occur in the counties listed in the table below:

State	County	State	County	State	County
Idaho	Idaho	Oregon	Benton	Washington	Asotin
	Lewis		Clackamas		Chelan
Nez Perce	Lane		Cowlitz		
Minnesota	Brown		Linn		Lewis
	Cottonwood		Marion	Lincoln	
	Goodhue		Polk	Spokane	
Montana	Jackson	Union	Wyoming	Laramie	
	Renville	Wallowa			
	Flathead	Washington			
	Lake	Yamhill			

For ground applications, the applicator must:

1. Apply when there is sustained wind away from native plant communities, OR
2. Use low-pressure nozzles according to manufacturer's specifications that produce only coarse or very coarse droplets, OR
3. Leave a 50 foot untreated buffer between the treatment and native plant communities

For aerial applications, the applicator must:

1. Apply only when there is sustained wind away from native plant communities, OR
2. Leave a 350 foot untreated buffer between the treatment and native plant communities

**USE RATES AND TIMING OF APPLICATION**

Timing of Application to Wheat	
Crop	Growth Stage
Spring Wheat Including durum	Apply prior to jointing, from 1 leaf to a maximum of 4 leaves on the main stem plus 2 tillers.
Winter Wheat Fall application	Minimum of 1 leaf in the fall.
Winter Wheat Spring application	Apply in spring as soon as wheat growth resumes. 1 leaf minimum to full tillering. Apply before jointing begins.

7/10

Recommended Rates and Timing of Application for Grass Weeds			
Grass Weeds	Rate	Growth Stage	Remarks
Wild Oat ( <i>Avena fatua</i> )	0.4 - 0.6 oz/A	1 leaf to 6 total leaves <sup>1</sup>	Control – Low to moderate infestations – 0.4 oz/A High infestations or when tank mixed with dicamba <sup>3</sup> – 0.6 oz/A
Volunteer Tame Oat ( <i>Avena sativa</i> )	0.4 - 0.6 oz/A	1 leaf to 6 total leaves <sup>1</sup>	Control
Green Foxtail ( <i>Setaria viridis</i> )	0.3 - 0.6 oz/A	1 leaf to 6 total leaves <sup>1</sup>	Control
Windgrass ( <i>Apera spica-venti</i> and <i>Apera interrupta</i> )	0.4 - 0.6 oz/A	1 leaf to 6 total leaves <sup>1</sup>	Control
Italian Ryegrass ( <i>Lolium multiflorum</i> )	0.6 oz/A	1 leaf to tillering <sup>2</sup>	Control <sup>4</sup> or Suppression
Cheat (True Cheat) ( <i>Bromus secalinus</i> )	0.6 oz/A	Apply when actively growing	Fall: Control Spring: Control <sup>4</sup> or Suppression
Japanese Brome ( <i>Bromus japonicus</i> )	0.6 oz/A	Apply when actively growing	Fall: Control Spring: Control <sup>4</sup> or Suppression
Downy Brome ( <i>Bromus tectorum</i> )	0.6 oz/A	Apply when actively growing	Suppression <sup>5</sup>
Yellow Foxtail ( <i>Setaria glauca</i> )	0.6 oz/A	1 leaf to 6 total leaves <sup>1</sup>	Suppression

<sup>1</sup> 1 leaf to 4 leaves on main stem plus 2 tillers

<sup>2</sup> 1 leaf to 4 leaves on main stem until end of tillering

<sup>3</sup> If EVEREST is applied in a tank mix combination with a dicamba-containing broadleaf herbicide, wild oat control may be reduced.

<sup>4</sup> Control is achieved using 1 quart of non-ionic surfactant per 100 gallons of spray solution (0.25 %v/v) + either liquid nitrogen fertilizer (2 qt/A and up to 50% of spray solution volume) OR ammonium sulfate fertilizer (nitrogen rate equivalent to 1.5 lb/A) **in winter wheat only**. Applications of liquid nitrogen fertilizer may result in temporary leaf burn or discoloration.

<sup>5</sup> Suppression is achieved using 1 quart of non-ionic surfactant per 100 gallons of spray solution (0.25 %v/v) + either liquid nitrogen fertilizer (2 qt/A and up to 50% of spray solution volume) OR ammonium sulfate fertilizer (nitrogen rate equivalent to 1.5 lb/A) **in winter wheat only**. Applications of liquid nitrogen fertilizer may result in temporary leaf burn or discoloration.

8/10

In addition to grass weeds, EVEREST alone will control or suppress the following broadleaf weeds. Applications should be made before broadleaf weeds are two inches in height.

Recommended Rates for Broadleaf Weeds		
Broadleaf Weeds	Rate	Remarks
Redroot Pigweed ( <i>Amaranthus retroflexus</i> )	0.3 - 0.6 oz/A	Control
Wild Mustard ( <i>Brassica kaber</i> )		Control
Black Mustard ( <i>Brassica nigra</i> )	0.4 - 0.6 oz/A	Control
Blue Mustard ( <i>Chorispora tenella</i> )		Control
Field Pennycress ( <i>Thlaspi arvense</i> )		Control
Shepherd's Purse ( <i>Capsella bursa-pastoris</i> )		Control
Tansy Mustard ( <i>Descurania pinnata</i> )		Control
Tumble Mustard ( <i>Sisymbrium altissimum</i> )		Control
Wild Turnip ( <i>Brassica rapa ssp. Sylvestris</i> )		Control
Volunteer Canola (conventional) ( <i>Brassica rapa ssp. Canola</i> )		Control
Flixweed ( <i>Descurania sophia</i> )	0.6 oz/A	Control
Small Seeded False Flax ( <i>Camelina microcarpa</i> )		Control
Burr Buttercup ( <i>Ranunculus testiculatus</i> )		Suppression
Common Waterhemp ( <i>Amaranthus tamariscinus</i> )		Suppression
Tall Wormseed Wildflower ( <i>Erysimum cheiranthoides</i> )		Suppression
Wild Buckwheat ( <i>Polygonum convolvulus</i> )		Suppression

Wheat exposed to water logged or saturated soils or temperature extremes such as heat or drought, low fertility, freezing weather or plant disease immediately prior to or after application could result in unacceptable injury symptoms. Weed control may also be reduced by these same conditions.

9/16

**ADJUVANT USE RATES**

EVEREST 70 WDG as a standalone or tank mix treatment may be mixed with adjuvants according to the following recommendations:

<b>Recommended Adjuvant Use Rates For Spring and Durum Wheat</b>	
EVEREST alone or with amine water soluble herbicides	Use 1 quart of non-ionic surfactant per 100 gallons (0.25% v/v).
EVEREST with ester or EC base herbicides	Do not add surfactant.
EVEREST with sulfonylurea herbicides + 2,4-D or dicamba <sup>1</sup>	Use 1 pint of non-ionic surfactant per 100 gallons (0.125% v/v). <ul style="list-style-type: none"> <li>Do not add surfactant if mixing with an ester or EC base 2,4-D</li> </ul>
<b>Recommended Adjuvant Use Rates For Winter Wheat</b>	
EVEREST alone or with any recommended herbicide tank mix	Use 0.5 – 1 quart of non-ionic surfactant per 100 gallons (0.125 – 0.25% v/v). <ul style="list-style-type: none"> <li>For improved performance on susceptible weeds, the following may be used with non-ionic surfactant:                             <ul style="list-style-type: none"> <li>liquid nitrogen fertilizer (2 qt/A and up to 50% of total spray solution<sup>2,3</sup>)</li> <li>OR</li> <li>ammonium sulfate fertilizer (nitrogen rate equivalent to 1.5 lb/A)</li> </ul> </li> </ul>

<sup>1</sup> If EVEREST is applied in a tank mix combination with a dicamba-containing broadleaf herbicide, wild oat control may be reduced.

<sup>2</sup> For fall applications to winter wheat, use liquid nitrogen fertilizer at a rate of 2 qt/A. For spring applications, use 2 qt/A and up to 50% of total spray solution.

<sup>3</sup> Applications of liquid nitrogen fertilizer may result in temporary leaf burn or discoloration.

10/10

**TANK MIXES**

For broader spectrum control of broadleaf weeds, EVEREST may be mixed with the broadleaf herbicides listed in the following table. Also, a non-ionic surfactant may be included in the spray solution. See "ADJUVANT USE RATES" section.

With all tank mix partners, read and follow the use directions, rates, precautions, timing, recropping restrictions, grazing interval restrictions and recommendations on broadleaf herbicide and surfactant labels. The tank mix must be used in accordance with the more restrictive label limitations and precautions for all pesticides used.

EVEREST Tank Mix Partners	Rate Per Acre
2,4-D Amine (4 lbs/gal)	0.5 to 1.5 pt
2,4-D Lo Volatile Ester (4 lbs/gal)	0.5 to 1 pt
2,4-D Lo Volatile Ester (6 lbs/gal)	0.33 to 0.67 pt
Agasco B-4	1.1 – 1.5 pt
Aim	0.33 to 1.24 oz
Aim EW	0.5 fl oz
Bromoxynil (2 lbs/gal)	1 to 2 pt
Broxomynil + MCPA (2 + 2 lbs/gal)	1 to 2 pt
Bronate Advanced	12.8 fl oz
Curtail	2 to 2.67 pt
Curtail M	1.75 pt
Dicamba (4 lbs/gal) <sup>1</sup>	2 to 4 fl oz
Double-Up B+D	0.75 – 1 pt
MCPA Amine or Ester <sup>2</sup> (3.7 lbs/gal)	0.5 to 1 pt
Starane	0.5 to 0.67 pt
Stinger	0.25 to 0.33 pt
Weco Max	16 oz
WideMatch	1 to 1.33 pt

<sup>1</sup> If EVEREST is applied in a tank mix combination with a dicamba-containing broadleaf herbicide, wild oat control may be reduced.

<sup>2</sup> Do not apply EVEREST in combination with MCPA/MCPA Ester (MCPE) within 72 hours of frost.

11/10

If one of the sulfonylurea herbicides in the following table is included with EVEREST for broadleaf control, 2,4-D or dicamba<sup>1</sup> is required in spring and durum wheat at the rate range listed in the table below. The addition of 2,4-D or dicamba<sup>1</sup> is not required in winter wheat. For adjuvant recommendations, see "ADJUVANT USE RATES" section.

Recommended Rates For EVEREST + Sulfonylurea Tank Mixes		
Sulfonylurea Tank Mix Partner	Rate Per Acre	In Spring and Durum Wheat, Add 2,4-D Or Dicamba <sup>1</sup> At The Following Rate Per Acre
Affinity TankMix	0.6 oz	2,4-D Amine or LV Ester (4 lbs/gal): 0.25 – 0.75 pt 2,4-D LV Ester (6 lbs/gal): 0.17 – 0.5 pt Dicamba <sup>1</sup> (4 lbs/gal): 2 – 4 fl oz
Affinity BroadSpec	0.4 – 0.6 oz	
Ally	0.1 oz	
Ally Extra	0.2 – 0.4 oz	
Amber	0.28 - 0.47 oz	
Express	0.17 to 0.33 oz	
Finesse	0.2 to 0.4 oz	
Harmony Extra	0.3 to 0.6 oz	
Harmony GT	0.3 to 0.6 oz	
Peak	0.38 to 0.5 oz	

<sup>1</sup> If EVEREST is applied in a tank mix combination with a dicamba-containing broadleaf herbicide, wild oat control may be reduced.

**PREPLANT, PREEMERGENCE AND PRE + POST USE DIRECTIONS  
FOR SPRING AND WINTER WHEAT**

**NOTICE TO PREPLANT AND PREEMERGENCE USERS:** Plant tolerance to this product for preplant and preemergence use has been found to be acceptable in research trials for the varieties tested. However, due to the large number of varieties and due to variable growing conditions, it is impossible to test every variety for tolerance to this product. Do not apply this product preplant or preemergence on more than 40 acres of wheat.

**APPLICATION PROCEDURES  
GROUND APPLICATION**

Apply in a spray volume of 5 - 10 gallons/acre (50 - 100 liters/hectare) at 30 psi to ensure proper coverage. If activating rainfall is not received within 7 - 10 days of application, performance may be reduced.

**AERIAL APPLICATION**

Apply in water using a minimum spray volume of 3 gallons/acre (or 30 liters/hectare). For best results, use a minimum of 5 gallons/acre (or 50 liters/hectare). Use nozzles that provide 200 to 350 micron size droplets for best results and to insure uniform spray coverage. Aerial applications with EVEREST should be made with low drift nozzles at a maximum height of 10 feet above the crop and at a maximum pressure of 40 psi. Do not apply aerially when wind speed is greater than 10 mph. Do not allow spray to drift onto adjacent crops, as injury or loss may occur. If activating rainfall is not received within 7 - 10 days of application, performance may be reduced.

See the "Aerial Drift Reduction Advisory Information" section of this label for additional information on how to reduce drift during aerial application.

12/10

**ADDITIONAL APPLICATION INFORMATION**

For postemergence applications following a preplant or preemergence application, observe use directions under "POSTEMERGENCE USE DIRECTIONS IN SPRING, DURUM AND WINTER WHEAT". Do not apply more than 0.018 lb acid equivalent (a.e.)/acre flucarbazone (0.6 ounce/acre of EVEREST) per growing season.

Do not apply preplant or preemergence to durum wheat.

**USE RATES AND TIMING OF APPLICATION**

For the control of wild oat and green foxtail, apply as follows:

Application Timing	Rate	Remarks
Preplant (apply up to 1 week prior to planting)	0.4 - 0.6 oz/A	For light infestations of wild oat, use 0.4 oz/A. For moderate to high infestations or in soils with high OM and/or low pH, use 0.6 oz/A for improved residual control.
Postplant/Preemergence	0.4 - 0.6 oz/A	
Preplant or Postplant/Preemergence + Postemergence <sup>1</sup>	0.2 oz/A + 0.4 oz/A <sup>1</sup> OR 0.3 oz/A + 0.3 oz/A <sup>1</sup> OR 0.4 oz/A + 0.2 oz/A <sup>1</sup>	Do not apply more than 0.6 oz/A of EVEREST per growing season.

<sup>1</sup> Make postemergence application with 0.25 – 0.75 pt/A of 2,4-D Amine or Ester (4 lbs/gal) or with 2 - 4 fl oz/A of Dicamba (4 lbs/gal).

**ADDITIONAL INFORMATION**

**SPRAYER CLEAN-UP**

Clean sprayer using the following procedures:

1. Drain the tank and thoroughly rinse spray tank, boom and hoses with clean water especially all visible deposits.
2. Fill the tank with water and add household ammonia to make a 1% v/v solution (1 gal/100 gal). Flush the hoses, boom and nozzles with the cleaning solution. Circulate for at least 15 minutes. Flush hoses, boom and nozzles once more and then drain the tank.
3. Clean nozzles and screens in a separate container using the 1% v/v solution of ammonia and water.
4. Repeat Step 2.
5. Rinse tank and flush boom and hoses with clean water.

Do not clean sprayer near desirable vegetation, wells or other water sources:

1. Dispose of all rinsate in accordance with pertinent regulations.
2. Check tank mix partner label for any additional clean-up procedures.

13/10

**RESISTANCE MANAGEMENT**

EVEREST is an acetolactate synthase (ALS) inhibiting herbicide. Any weed population may contain or develop plants naturally resistant to a herbicidal mode of action. Resistant biotypes may eventually dominate the weed population if herbicides with an identical mode of action are used repeatedly in the same field and weed control may fail. Where possible, rotate the use of EVEREST with herbicides that have a different mode of action.

Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist. The use of EVEREST should conform to resistance management strategies established for the use area. Consult your agricultural advisor for resistance management strategies and recommended pest management practices for your area.

**CROP ROTATION RESTRICTIONS**

<b>0 Days</b>		
Spring and Winter Wheat		
<b>4 Months</b>		
Durum Wheat		
<b>9 Months</b>		
Barley	Canola	Potatoes
Soybeans	Sugar beets	
<b>11 Months</b>		
Field peas		
<b>24 Months</b>		
Lentils	Mustard	

As EVEREST is degraded by soil microbes, environmental conditions that decrease microbial activity must be considered when making rotational cropping decisions. These environmental conditions include prolonged drought and/or cold temperatures within and following the cropping season, as well as soils with both low OM (less than 2%) and high pH (greater than 7.5). If these conditions exist, a soil bioassay may be necessary to ensure rotational crop safety.

**AERIAL DRIFT REDUCTION ADVISORY INFORMATION**

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment-and-weather-related factors determines the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions.

The following drift management requirements must be followed to avoid off-target drift movement from aerial applications to agricultural field crops. These requirements do not apply to forestry applications, public health uses or to applications using dry formulations.

1. The distance of the outer most nozzles on the boom must not exceed ¾ the length of the wingspan or rotor.
2. Nozzles must always point backward, parallel with the air stream and never be pointed downwards more than 45 degrees.

14/10

When applying EVEREST in a tank mix with other herbicides (e.g. 2,4-D, bromoxynil, dicamba, MCPA, sulfonyleurea herbicides) in eastern Washington, observe all applicable Washington State Department of Agriculture herbicide rules.

The applicator should be familiar with and take into account the information covered in the Aerial Drift Reduction Advisory Information.

### **Information On Droplet Size**

The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions (see Wind, Temperature and Humidity, and Temperature Inversions).

### **Controlling Droplet Size**

- Volume – Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.
- Pressure – Do not exceed the nozzle manufacturer's recommended pressures. For many nozzle types, lower pressure produces larger droplets. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.
- Number of Nozzles – Use the minimum number of nozzles that provide uniform coverage.
- Nozzle Orientation – Orienting nozzles so that the spray is released parallel to the airstream produces larger droplets than other orientations and is the recommended practice. Significant deflection from horizontal will reduce droplet size and increase drift potential.
- Nozzle Type – Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce the largest droplets and the lowest drift.

### **Boom Length**

For some use patterns, reducing the effective boom length to less than  $\frac{3}{4}$  of the wingspan or rotor length may further reduce drift without reducing swath width.

### **Application Height**

Applications should not be made at a height greater than 10 feet above the top of the largest plants unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind.

15/14

### Swath Adjustment

When applications are made with a crosswind, the swath will be displaced downwind. Therefore, on the up and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the aircraft upwind. Swath adjustment distance should increase, with increasing drift potential (higher wind, smaller drops, etc.)

### Wind

Drift potential is lowest between wind speeds of 2 to 10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given speed. Application should be avoided below 2 mph due to variable wind direction and high inversion potential. NOTE: Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

### Temperature And Humidity

When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

### Temperature Inversions

Applications should not occur during a temperature inversion because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue in the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**Pesticide Storage:** Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store in original container and out of reach of children, preferably in a locked storage area.

Handle and open container in a manner as to prevent spillage. If the container is leaking or material spilled for any reason or cause, carefully sweep material into a pile. Refer to Precautionary Statements on label for hazards associated with the handling of this material. Do not walk through spilled material. Dispose of pesticide as directed below. In spill or leak incidents, keep unauthorized people away. For help with any spill, leak, fire or exposure involving this material, call day or night CHEMTREC (703) 527-3887 or (800) 424-9300.

**Pesticide Disposal:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**Container Disposal:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

16/16

### CONDITIONS OF SALE

1. Arysta LifeScience North America Corporation (“Arysta”) warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label only when used in accordance with the directions under normal conditions of use.
2. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal use conditions, or under conditions not reasonably foreseeable to Arysta. **ARYSTA DISCLAIMS ALL OTHER WARRANTIES, EXPRESSED OR IMPLIED, INCLUDING ANY WARRANTY OF FITNESS FOR A PARTICULAR PURPOSE OR MERCHANTABILITY. TO THE EXTENT PERMITTED BY LAW, SELLER SHALL NOT BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM OR IN CONNECTION WITH THE MANUFACTURE, SALE, DELIVERY, USE, HANDLING OR STORAGE OF THIS PRODUCT. TO THE EXTENT PERMITTED BY LAW, SELLER’S LIABILITY SHALL BE LIMITED TO THE REFUND OF THE PURCHASE PRICE. ARYSTA DOES NOT AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE ANY OTHER WARRANTY, GUARANTEE OR REPRESENTATION CONCERNING THIS PRODUCT.**
3. Critical and unforeseeable factors beyond the control of Arysta prevent Arysta from eliminating all risks in connection with the use of this product. Such risks include, but are not limited to, damage to plants and crops to which product is applied, lack of complete control, and damage caused by drift to other plants or crops. Such risks occur even though the product is reasonably fit for the use stated on the label and even though label directions are followed. Except as stated in 1 above, to the extent permitted by law, by purchasing, accepting and using this product, the buyer and user acknowledge and assume all risks and liabilities resulting from handling, storage, and use of this product.

Aim is a Registered Trademark of FMC Corp.

Amber and Peak are Registered Trademarks of Syngenta.

Buctril, Bronate are Registered Trademarks of Bayer CropScience.

Curtail, Stinger, Starane and WideMatch are Registered Trademarks of Dow AgroSciences LLC.

Ally, Ally Extra, Express, Finesse, Harmony Extra, Harmony GT, Affinity BroadSpec and Affinity TankMix are Registered Trademarks of E.I. DuPont de Nemours & Co.

Double-Up is a Registered Trademark of Helena Holding Co.

B-4 is a Trademark of Agsco, Inc.

Weco Max is a Trademark of Wilbur-Ellis Co.