

66330-52

02/28/2007

1/14



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, D.C. 20460

OFFICE OF  
PREVENTION, PESTICIDES  
AND TOXIC SUBSTANCES

Doina Bujor  
Arysta LifeScience North America Corporation  
15401 Weston Parkway, Suite 150  
Cary, NC 27513

FEB 28 2007

Dear Ms Bujor:

Subject: Labeling Amendment; Addition of Potatoes & Grapes and Minor Changes  
CELERO 16 WSG  
EPA Registration No. 66330-52

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable with the following provisions:

1. All references to emerald ash borer must be deleted from the label. To have this pest added on at a future date, acceptable efficacy data must be submitted to the Agency.
2. Within the CONDITIONS OF SALE, the phrase "*to the extent permitted by law*" must be rephrased "*to the extent permitted by applicable law*".

A stamped copy is enclosed for your records. Please submit one (1) final printed copy for the referenced label, incorporating the above changes, before releasing the product for shipment.

Sincerely,

A handwritten signature in black ink, appearing to read "Kable Bo Davis", written over a horizontal line.

Kable Bo Davis  
Entomologist  
Insecticide-Rodenticide Brach  
Registration Division (7505C)

Enclosure

2/14

Alternate Brand Name for CELERO™ 16 WSG:

- BELAY™ 16 WSG

**CELERO™ 16 WSG**  
**INSECTICIDE**

**GROUP 4A INSECTICIDE**

FOR SYSTEMIC CONTROL OF INSECTS ON ORNAMENTAL PLANTS GROWN IN FIELD NURSERY PLANTINGS, CONTAINER NURSERIES, FOREST NURSERIES, GREENHOUSES, LATH AND SHADE HOUSES, INTERIORSCAPES AND OTHER SITES WHERE PLANTS ARE GROWN FOR ORNAMENTAL, AESTHETIC AND CLIMATE MODIFICATION PURPOSES AND FOR USE ON GRAPES, POTATO AND TOBACCO

<u>INGREDIENTS</u>	<u>By wt.</u>
Active Ingredient	
*Clothianidin.....	16.0%
Other Ingredients.....	84.0%
Total.....	100.0%
*(E)-1-(2-chloro-1,3-thiazol-5-ylmethyl)-3-methyl-2-nitroguanidine.	

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

See back panel for additional precautionary statements.

For Product Information Call: 1-866-761-9397

ARYSTA LIFESCIENCE NORTH AMERICA CORPORATION  
15401 Weston Parkway, Suite 150  
Cary, NC 27513

EPA Reg. No. 66330-52  
EPA Est. No. \_\_\_\_\_

NET WEIGHT \_\_\_\_\_ OUNCES

**ACCEPTED**  
**with COMMENTS**  
**In EPA Letter Dated:**

**FEB 28 2007**

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
as amended, for the pesticide  
registered under EPA Reg. No.  
66330-52

<b>FIRST AID</b>	
<b>If in eyes</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If swallowed</b>	<ul style="list-style-type: none"> <li>• Call poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by the poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
<b>If on skin or clothing</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If inhaled</b>	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>
Have the product container or label with you when calling a poison control center or doctor, or going for treatment.	
<b>HOTLINE NUMBERS:</b>  <b>FOR 24-HOUR MEDICAL EMERGENCY MEDICAL ASSISTANCE:</b> Call PROSAR at 1-800-303-6952 or 1-651-632-8946 if calling from outside of the U.S.  <b>FOR 24-HOUR CHEMICAL EMERGENCY ASSISTANCE:</b> Spill, leak, fire, exposure, or accident, call CHEMTREC at 1-800-424-9300 or 1-703-527-3887 if calling from outside of the U.S.	

## **PRECAUTIONARY STATEMENTS**

### **HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**CAUTION.** Causes moderate eye irritation. Harmful if swallowed, absorbed through skin or inhaled. Avoid contact with eyes, skin or clothing. Avoid breathing spray mist. Remove contaminated clothing and wash clothing before reuse. Wash thoroughly with soap and water after handling.

### **PERSONAL PROTECTIVE EQUIPMENT (PPE)**

Some of the materials that are chemical-resistant to this product are listed below. If you want more options, follow the instruction for category A on an EPA chemical-resistance category selection chart.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Socks

- Shoes, and
- Chemical-resistant gloves made of any waterproof material such as polyethylene or polyvinyl chloride (PVC).

Follow the manufacturer's instructions for cleaning/maintaining PPE. If no instructions for washables are available, use detergent and hot water. Keep and wash PPE separately from other laundry.

### **ENGINEERING CONTROLS**

When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

**IMPORTANT:** When reduced PPE is worn because a closed system is being used, handlers must be provided all PPE specified above for "applicators and other handlers" and have such PPE immediately available for use in an emergency, such as a spill or equipment break-down.

### **USER SAFETY RECOMMENDATIONS**

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

### **ENVIRONMENTAL HAZARDS**

This product is toxic to aquatic invertebrates. Do not apply when weather conditions favor drift from treated areas. Drift and runoff may be hazardous to aquatic organisms in neighboring areas. Do not apply where runoff is likely to occur. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters or rinsate.

This product is toxic to bees exposed to treatment and for more than 5 days following treatment. Do not apply this product to blooming, pollen-shedding or nectar-producing parts of plants if bees may forage on the plants during this time period, unless the application is made in response to a public health emergency declared by appropriate state or federal authorities.

### **DRIFT MANAGEMENT**

Do not allow this product to drift onto neighboring crops or non-crop areas or use in a manner or at a time other than in accordance with label directions because animal, plant or crop injury, illegal residues or other undesirable results may occur.

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment and weather-related factors determine the potential for spray drift. The applicator is responsible for considering all these factors when making decisions. Where states have more stringent regulations, they should be observed.

**DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Carefully read and understand the Directions for Use and restrictions before applying this product.

Do not apply this product in any way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

**AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions on this label about personal protective equipment (PPE), notification to workers and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry into treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is: coveralls, socks, shoes, chemical-resistant gloves made of any waterproof material.

**NON- AGRICULTURAL USE REQUIREMENTS**

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Keep children and pets off treated area until the spray has dried.

**GENERAL INFORMATION**

CELERO™ 16 WSG Insecticide is a systemic insecticide for broad spectrum control of insect pests infesting ornamentals, grapes, potato and tobacco. When applied as directed, CELERO 16 WSG provides excellent and long residual control of insects listed on this label.

**\* DO NOT USE ON FERNS GROWN FOR CUT FOLIAGE**

#### **NOTICE TO USER**

Plant tolerance to this product has been found to be acceptable in research trials for the general plant species listed on the label. However, due to the large number of species and their associated varieties or cultivars, and due to variable growing conditions, it is impossible to test every plant and variety or cultivar for tolerance to this product. Neither the Manufacturer nor the Seller has determined whether or not this product can be safely used on all cultivars. The user should determine if this product can be used safely prior to commercial use. In a smaller test area, test the recommended rates on plants for phytotoxicity prior to general use.

#### **RESISTANCE MANAGEMENT**

Certain insects have demonstrated a tendency to develop resistance to insect control products. For these insects, avoid treating consecutive generations with the same product or products from the same class (Group 4A Insecticides). CELERO 16 WSG is a member of the neonicotinoid insecticides and should be rotated with products that belong to different classes. Always consult with your local agricultural specialist for guidance and information on area resistance management programs. In the absence of a locally coordinated program, the following IPM strategy will help to ensure the prolonged usefulness of CELERO 16 WSG and conventional insecticides. In order to reduce selection pressure, which can lead to the development of resistance, the following strategies are recommended.

1. Incorporate IPM principles into your insect control program.
2. Ensure thorough spray coverage of all foliage.
3. Always apply CELERO 16 WSG at the recommended rates (don't use less than labeled rates of any insect control product when applied alone or in tank mixtures) and apply at the proper timing according to label recommendations.
4. Use CELERO 16 WSG in alternation with other insecticides having different modes of action.

#### **PLANT TOLERANCE**

Neither the manufacturer nor the seller has determined whether or not CELERO 16 WSG can be used safely on all ornamentals. See the list of ornamental species, varieties and cultivars that CELERO 16 WSG has been tested on. Since your growing conditions may be different, it is highly recommended that a small number of plants under your conditions be sprayed to make certain that no phytotoxicity occurs, prior to any large scale application to plants under commercial production. The end user assumes all risks arising from application of CELERO 16 WSG in a manner inconsistent with its labeling. **DO NOT USE ON FERNS GROWN FOR CUT FOLIAGE.**

#### **COMPATIBILITY OF MIXTURES**

To obtain broad-spectrum pest control, CELERO 16 WSG can be tank-mixed with other pesticides. However, all plant species and their varieties and cultivars have not been tested with possible tank mix combinations, sequential pesticide treatments and adjuvants and surfactants. The user should conduct a spray mix compatibility and

phytotoxicity trial under local conditions to ensure compatibility prior to any large-scale use.

### **PREPARATION OF SPRAY SOLUTION**

Begin with clean equipment. Add sufficient clean water to the spray tank for one-half of the mix load. Start tank agitation. Add and properly suspend the necessary amount of CELERO 16 WSG before adding any other products. Agitate to ensure thorough mixing while adding the rest of the water. A high quality wetting agent or other spray adjuvant, approved for use on your crop, may be added to spray solutions according to the manufacturer's use instructions. CELERO 16 WSG should be properly suspended and diluted prior to the addition of any adjuvant. Consult the adjuvant label or manufacturer for crop tolerance and safety information when used with this product. Maintain agitation during mixing and application. DO NOT allow the spray mixture to stand without agitation. Apply with properly calibrated spray equipment. If agitation is stopped for any reason, the spray solution must be thoroughly re-mixed prior to any further use.

### **APPLICATION THROUGH IRRIGATION SYSTEMS**

CELERO 16 WSG can be applied through irrigation systems at rates recommended on the label. Meter the product into the irrigation water during the first part of the irrigation cycle. Prepare a solution of CELERO 16 WSG in water using the appropriate dilution for the injection system. The product may be diluted before injection to obtain the appropriate concentration for the injection system. Make sure the injection solution is properly mixed during the injection process.

Apply CELERO 16 WSG only through micro irrigation (individual spaghetti tubes), drip irrigation, overhead irrigation, ebb and flood, or hand-held or motorized calibrated irrigation equipment.

Do not apply this product through any other type of irrigation system. Crop injury or lack of effectiveness can result from non-uniform distribution of treated water.

If you have any questions about calibration, contact your State Extension Service specialist, equipment manufacturers or other experts in this area.

Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.

A person knowledgeable of the chemigation system and responsible for its operation, or a person who is under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

### **SAFETY DEVICES FOR IRRIGATION SYSTEMS CONNECTED TO PUBLIC WATER SUPPLIES:**

If the source of water for your irrigation system is a public water supply, follow the instructions below:

1. Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.
2. Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or over-flow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.
3. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
4. The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
5. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.
6. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. Do not apply when wind speed favors drift beyond the area intended for treatment.

**SAFETY DEVICES FOR IRRIGATION SYSTEMS NOT CONNECTED TO A PUBLIC WATER SUPPLY:**

1. The system must contain a functional check valve, vacuum relief valve and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
2. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.



## APPLICATION TO ORNAMENTALS

CELERO 16 WSG can be used for systemic insect control in ornamentals grown in\*:

- Field nursery plantings
- Container nurseries
- Forest nurseries
- Greenhouses
- Lath and shade houses
- Interiorscapes
- Other sites where plants for ornamental purposes are grown for aesthetic or climate modification uses.

CELERO 16 WSG can be applied as a foliar, drench, surface application or through an irrigation system.

CELERO 16 WSG can be used for insect control on these plants:

Annuals
Evergreens
Flowering Plants
Foliage Plants (except for ferns grown for cut foliage)
Ground Covers
Non-Bearing Fruits and Nuts (no harvestable fruit within 12 months of the last application)
Perennials
Shrubs
Trees

## FOLIAR APPLICATIONS

CELERO 16 WSG should be applied when threshold populations are observed. Do not wait until large insect populations are established. Apply CELERO 16 WSG in sufficient water to ensure uniform and thorough coverage of foliage. Thorough coverage is required for optimum control. To achieve adequate coverage, use proper spray pressures, nozzles, nozzle spacing, and water volume per acre. Use lower rates for light infestations, and/or smaller plants and higher rates for heavy infestations and/or larger plants.

## RATES OF APPLICATION FOR FOLIAR APPLICATION

FOLIAR APPLICATION	
PEST	DOSAGE (oz/100 gallons of water)
Aphids	2 - 4
Azalea Lace Bug	2 - 4

10/14

Flea Beetles	2 - 4
Japanese Beetle (adults)	3 - 4
Leaf Hoppers (including Glassy Winged Sharpshooter)	2 - 4
Mealybugs	4
Scales, Soft	3 - 4
Whiteflies	4

**Note:** Do not apply more than a total of 40 oz of CELERO 16 WSG (0.4 lbs active ingredient) per acre per season.

#### FOR CONTROL OF INSECTS AND MITES

If mites are present or expected when making foliar applications of CELERO 16 WSG for insect control, apply a tank mix of CELERO 16 WSG and SHUTTLE 15 SC at labeled rates. CELERO 16 WSG is also compatible with most other miticides registered for use in ornamentals. When using tank mixtures, always follow the most restrictive label statements.

#### DRENCH, SURFACE, AND IRRIGATION SYSTEM APPLICATIONS

When applying CELERO 16 WSG as a drench or irrigation system application, use enough solution to wet the growing media in the pots or the root zone of the plants. When applying CELERO 16 WSG as a surface application, water in with enough irrigation to move the insecticide into the root zone of the plants.

#### RATES OF APPLICATION FOR DRENCH, SURFACE, AND IRRIGATION SYSTEM APPLICATIONS

DRENCH, SURFACE, AND IRRIGATION SYSTEM APPLICATIONS	
PEST	DOSAGE (oz/100 gallons of water)*
Aphids	2 - 4
Azalea Lace Bug	2 - 4
Emerald Ash Borer	2 - 4
Flea Beetles	2 - 4
Fungus Gnats	2 - 4
Japanese Beetles (adults)	3 - 4
Leaf Hoppers (including Glassy Winged Sharpshooter)	2 - 4
Leaf Miners	3 - 4
Mealybugs	4

Root Weevils (including Black Vine Weevil)	3 - 4
Scales, Soft	3 - 4
Whiteflies	4
White Grubs (Japanese Beetle Green June Beetle European Chafer Northern Masked Chafer Southern Masked Chafer Oriental Beetle Asiatic Garden Beetle)	2 - 4

**Note:** Do not apply more than a total of 40 oz of CELERO 16 WSG (0.4 lbs active ingredient) per acre per season.

\* Apply sufficient drench or irrigation solution to thoroughly wet the media or root zone of the plants.

The specified amount of product should be mixed with a sufficient volume of water to evenly disperse the solution into the pots and to thoroughly wet the planting medium, without resulting in overflow or leaching and the loss of product. When watering treated pots, avoid watering to the point of overflowing or leaching.

If desired, less than full drench volumes can be applied to pots provided the solution is watered in with overhead irrigation. To determine the proper concentration, calculate the amount of product needed by determining how much water is needed to wet the pots of the ornamentals to be treated. Then add the desired amount of product to the amount of water to be used and add a proportionate amount of the solution to each pot and then water the solution into the growing media or root zone of the plants.

**Ornamentals that have been evaluated and found to be tolerant of CELERO 16 WSG applied at recommended rates and methods of application.\***

African Daisy	Lantana
Argyranthemum	Lobelia
Ageratum	Mandevilla
Aster	Marigold
Bacopa	New Guinea Impatiens
Begonia	Osteospermum
Calibrachoa	Nicotiana
Chrysanthemum	Pansy
Cleyera	Petunia
Cleome	Poinsettia, pre and post bract formation
Coleus	Primula
Dahlia	Red Bud
Daylily	Rose

12/14

Diascia Easter Lily Eunomymus Fuschia Geranium, Zonal Geranium, Ivy Gerbera Holly Impatiens Indian Hawthorne Ivy	Miniature Rose Salvia Scaevola Shasta Daisy Snapdragon Stock Torenia Verbena Vinca Zinnia
<p>*Since your growing conditions may be different than the conditions under which the ornamentals were tested for phytotoxicity, it is highly recommended that a small number of plants growing under your conditions be sprayed to make certain that no phytotoxicity occurs prior to any large scale application to plants under commercial production. The end user assumes all risks arising from application of CELERO 16 WSG in a manner inconsistent with its labeling.</p>	
<p><b>GENERAL RESTRICTIONS AND LIMITATIONS</b></p> <ul style="list-style-type: none"> <li>• DO NOT USE ON FERNS GROWN FOR CUT FOLIAGE</li> <li>• Do not apply more than a total of 40 oz of CELERO 16 WSG (0.4 lbs active ingredient) on ornamental plants per acre per season.</li> </ul>	

GRAPES (Drip irrigation, chemigation application)		
METHOD	PEST	APPLICATION RATE
Drip irrigation (chemigation)	Grape mealybug Grape leafhopper Grape berry moth	10 – 20 oz/A (1.6 – 3.2 oz ai/A)
	Japanese beetle	20 oz/A (3.2 oz ai/A)
<p><b>GENERAL RESTRICTIONS AND LIMITATIONS</b></p> <ul style="list-style-type: none"> <li>• Apply 1-2 applications per seasons.</li> <li>• Do not apply more than a total of 20 oz of CELERO 16 WSG (3.2 oz active ingredient) per acre per season.</li> <li>• PHI = 30 days.</li> </ul>		

POTATO (In-furrow or Side-dress application)		
METHOD	PEST	APPLICATION RATE
In-furrow or Side-dress	Colorado potato beetle Aphids	12 – 18 oz/A (1.9 – 2.9 oz ai/A)

13/14

	Leafhoppers	
<b>GENERAL RESTRICTIONS AND LIMITATIONS</b> <ul style="list-style-type: none"> <li>• 1 application per season.</li> <li>• Apply at planting.</li> </ul>		

#### TOBACCO (SOIL APPLICATION)

CELERO 16 WSG must be mixed with water prior to application. Keep the CELERO 16 WSG solution agitated or mix regularly to avoid settling in the spray or transplant tank. Apply CELERO 16 WSG at the specified dosage to the soil in transplant water or by drench to trays or flats prior to transplanting. Plants in flats or trays should be watered by overhead irrigation to wash CELERO 16 WSG from the foliage into the potting media. Failure to wash CELERO 16 WSG from foliage may result in a reduction of control.

<b>TOBACCO - SOIL APPLICATION</b> <b>(in transplant water application or tray or flat drench application)</b>	
<b>PEST</b>	<b>APPLICATION RATE</b>
Flea Beetles	5 to 10 oz of product per acre (0.8-1.6 oz ai/A)
Aphids	10 oz of product per acre (1.6 oz ai/A)
<b>SPECIAL INSTRUCTIONS AND LIMITATIONS</b> <ul style="list-style-type: none"> <li>• Soil applications may sometimes cause yellowing on the leaves present at the time of transplanting. This effect does not have any impact on plant growth and yields.</li> <li>• Do not apply more than a total of 0.2 lb active ingredient of CELERO 16 WSG per acre per season.</li> </ul>	

#### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**PESTICIDE STORAGE:** Store in a dry place away from excessive heat. Do not store near food or feed. Store in original container only. To close package, replace and tighten cap to form an airtight seal.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** Triple rinse (or equivalent), then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or (if allowed by State and local authorities) by burning. If burned, stay out of smoke.

14/14

#### CONDITIONS OF SALE

1. Arysta LifeScience North America Corporation ("Arysta") warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in accordance with the directions under normal conditions of use.
2. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal use conditions, or under conditions not reasonably foreseeable to Arysta. **ARYSTA DISCLAIMS ALL OTHER WARRANTIES, EXPRESSED OR IMPLIED, INCLUDING ANY WARRANTY OF FITNESS FOR A PARTICULAR PURPOSE OR MERCHANTABILITY. TO THE EXTENT PERMITTED BY LAW, SELLER SHALL NOT BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM OR IN CONNECTION WITH THE MANUFACTURE, SALE, DELIVERY, USE, HANDLING OR STORAGE OF THIS PRODUCT. TO THE EXTENT PERMITTED BY LAW, SELLER'S LIABILITY SHALL BE LIMITED TO THE REFUND OF THE PURCHASE PRICE. ARYSTA DOES NOT AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE ANY OTHER WARRANTY, GUARANTEE OR REPRESENTATION CONCERNING THIS PRODUCT.**
3. Critical and unforeseeable factors beyond Arysta's control prevent Arysta LifeScience from eliminating all risks in connection with the use of this product. Such risks include, but are not limited to, damage to plants and crops to which the product is applied, lack of complete control, and damage caused by drift to other plants or crops. Such risks occur even though the product is reasonably fit for the use stated on the label and even though label directions are followed. Except as stated in 1 above, to the extent permitted by law, by purchasing, accepting and using this product, the buyer and user acknowledge and assume all risks and liabilities resulting from handling, storage, and use of this product.

CELERO is a trademark of Arysta LifeScience North America Corporation.