

66330-56

6/5/2014

1/30

David G. Bolin, Ph.D.
Regulatory Affairs Manager
Arysta Life Science North America, LLC
15401 Weston Parkway, Suite 150
Cary, NC 27513

JUN 05 2014

Subject: Arysta Life Science North America, LLC; Endorse Water Dispersible Granules, EPA Registration No. 66330-56; Amendment to Add Disease Listings and Expand Uses; D# 489082, Application Dated 3/20/14

Dear. Mr. Bolin:

The amendment referred to above, submitted in connection with registration under FIFRA section 3(c)(7)(A), is acceptable provided that you:

1. Submit and/or cite all data required for registration/reregistration of your product under FIFRA section 3(c)(5) when the Agency requires all registrants of similar products to submit such data.
2. Submit two (2) copies of your final printed labeling before you release the product for shipment. Final printed labeling means the label or labeling of the product when distributed or sold. Clearly legible reproductions or photo reductions will be accepted for unusual labels, such as those silk-screened directly onto glass or metal containers or large bags or drum labels.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(b). Your release for shipment of the product bearing the amended labeling constitutes acceptance of these conditions.

If you have any questions contact Chris Pfeifer at 703-308-0031 or by email at: pfeifer.chris@epa.gov.

Sincerely,

Sheryl K. Reilly, Associate Branch Chief
Biochemical Pesticides Branch
Biopesticides and Pollution
Prevention Division (7511P)

Enclosure

CONCURRENCES

SYMBOL	>	7511P	7511P					
SURNAME	>	Pfeifer	Reilly					
DATE	>	6/5/14	6/5/14					

2/30

ACCEPTED

Alternate Brand Name for ENDORSE Water Dispersible Granules: JUN 05 2014

- VERANDA™ T Water Dispersible Granules
- Ph-D™ Fungicide
- VERANDA™ O

[Language within brackets is optional text.]

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 66330-56

ENDORSE® Water Dispersible Granules

For use on Turf[*], Ornamentals[*], and Various Crops

For control or suppression of listed diseases on cucurbit vegetables[*], fruiting vegetables[*], ginseng*, grapes, pome fruits[*], root and tuber vegetables[*], berry and small fruit[*], strawberries[*], stone fruits[*], pomegranates[*], tree nuts[*], and non-bearing vine fruit and fruit and nut trees grown in interiorscapes[*], ornamentals[*], Christmas trees (nurseries and plantations)[*], field nursery and container planting, forest nurseries, ornamentals grown in greenhouses, lath and shadehouses and other enclosed structures.

[*Not for use in California.]

GROUP 19 FUNGICIDE

ACTIVE INGREDIENT:

Polyoxin D zinc salt	11.3%
OTHER INGREDIENTS:	88.7%
TOTAL	100.0%

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE SIDE OR BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

FIRST AID	
If on skin or clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If swallowed	<ul style="list-style-type: none"> • Call poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If inhaled	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
HOTLINE NUMBERS	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment.	
FOR 24-HOUR EMERGENCY MEDICAL ASSISTANCE	
Call PROSAR at 1-866-303-6952 or 1-651-632-8946 if calling from outside of the U.S.	

EPA Reg. No. 66330-56
ADxxxxx
Produced for:

EPA Est. No. _____
Net Content: 1 lb, 2 lb 6.4 oz (2.4 lb)

0579300A

3/30

Arysta LifeScience North America, LLC
15401 Weston Parkway, Suite 150
Cary, NC 27513

For Product Information Call 1-866-761-9397

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if absorbed through the skin, swallowed or inhaled. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Avoid breathing dust or spray mist. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

All mixers, loaders, applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Socks
- Shoes
- Chemical-Resistant gloves

User Safety Requirements

Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables are available, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

- Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is moderately toxic to aquatic invertebrates and fish. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate. Do not allow runoff into lakes, streams, ponds or public waterways. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas.

Granules exposed on soil surface may be hazardous to fish and aquatic invertebrates. Collect granules spilled during loading.

Observe the most restrictive of the labeling limitations and precautions of all products used in mixtures.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

For any requirements specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to statements on this label about personal protective equipment (PPE), notification to workers, and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 hours.

PPE required for early entry into treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is: coveralls, socks, shoes, chemical resistant gloves.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in the box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses.

Keep children and pets out of treated area until sprays have dried.

PRODUCT INFORMATION

ENDORSE Water Dispersible Granules is not for use on turf being grown for commercial seed production.

ENDORSE Water Dispersible Granules is not for homeowner use to treat food crops.

ENDORSE Water Dispersible Granules is for use in controlling or suppressing certain diseases on crops listed in this label.

Apply ENDORSE Water Dispersible Granules as a preventive or curative treatment in conjunction with good management practices.

RESISTANCE MANAGEMENT RECOMMENDATIONS

ENDORSE Water Dispersible Granules contains a Group 19 fungicide. Fungal isolates with acquired resistance to Group 19 may eventually dominate the fungal population if Group 19 fungicides are used repeatedly in the same field or in successive years as the primary method of control for targeted species. This may result in partial or total loss of control of those species by ENDORSE Water Dispersible Granules or other Group 19 Fungicides.

To delay fungicide resistance, consider:

- Avoiding the consecutive use of ENDORSE Water Dispersible Granules or other target site of action Group 19 fungicides that have a similar target site of action, on the same pathogens.
- Using tank mixtures or premixes with fungicides from different target site of action Groups as long as the involved products are all registered for the same use and are both effective at the tank mix or prepack rate on the pathogen(s) of concern.
- Basing fungicide use on a comprehensive IPM program.
- Monitoring treated fungal populations for loss of field efficacy.
- Contacting your local extension specialist, certified crop advisors, and/or manufacturer for fungicide resistance management and/or IPM recommendations for specific crops and resistant pathogens.

MIXING AND APPLICATION

When diseases not specified on this label are present or expected, it is appropriate to mix ENDORSE Water Dispersible Granules with appropriately labeled fungicides. When tank mixing ENDORSE Water Dispersible Granules with other products, observe all precautions and limitations on each separate product label. It is always advisable to conduct a tank mix compatibility test when you plan to mix this product with other products. To determine the physical compatibility of this product with other products, use a jar test. Using a quart jar, add the proportionate amounts of the products to approximately one quart of water with agitation. Add dry formulations first, then flowables, then emulsifiable concentrates last. After thorough mixing, allow this mixture to stand for 5 minutes. If the combination remains mixed or can be readily remixed, it is physically compatible. Once compatibility has been proven, use the same procedure for adding required ingredients to the spray tank. To assess the potential for phytotoxicity, test tank mixtures on a small number of plants prior to more widespread application.

Do not apply this product through any type of irrigation system.

Mixing instructions for ENDORSE Water Dispersible Granules:

- Fill spray tank with water to 1/2 of the intended final volume.
- Start spray tank agitation.
- Add the appropriate amount of product to the tank according to the rates in the following tables. Follow the directions above for mixing other products in the spray tank.
- Agitate to ensure thorough mixing while adding the remaining required water.
- DO NOT allow the spray mixture to stand without agitation.

Mix only the amount of spray solution needed to treat the desired area.

For optimum control of labeled diseases, apply ENDORSE Water Dispersible Granules with sufficient water to provide a thorough coverage.

STRAWBERRIES[*]

BERRY AND SMALL FRUIT[*]

Amur river grape; aronia berry; bayberry; bearberry; bilberry; blackberry (including Andean blackberry, arctic blackberry, bingleberry, black satin berry, boysenberry, brombeere, California blackberry, Chesterberry, Cherokee blackberry, Cheyenne blackberry, common blackberry, coryberry, darrowberry, dewberry, Dirksen thornless berry, evergreen blackberry, Himalayaberry, hullberry, lavacaberry, loganberry, lowberry, Lucretiaberry, mammoth blackberry, marionberry, mora, mures deronce, nectarberry, Northern dewberry, olallieberry, Orgeon evergreen berry, phenomenalberry, rangeberry, ravenberry, rossberry, Shawnee blackberry, Southern dewberry, tayberry, youngberry, zarzamora, and cultivars, varieties and/or hybrids of these); blueberry, highbush; blueberry, lowbush; buffalo currant; buffaloberry; che; Chilean guava; chokecherry; cloudberry; cranberry; cranberry, highbush; currant, black; currant, red; elderberry; European barberry; gooseberry; honeysuckle, edible; huckleberry; jostaberry; Juneberry (Saskatoon berry); kiwifruit, fuzzy; kiwifruit, hardy; lingonberry; maypop; mountain pepper berries; mulberry; muntries; native currant; partridgeberry; phalsa; pincherry; raspberry, black and red; riberry; salal; schisandra berry; sea buckthorn; serviceberry; strawberry; wild raspberry; cultivars, varieties, and/or hybrids of these

Disease Control	Rate	Application Notes
<p>Alternaria rot, Black leaf spot[*] (<i>Alternaria alternata</i>)</p> <p>Fruit rots, minor[*] (<i>Alternaria spp.</i>)</p> <p>Botrytis fruit rot and blossom blight, gray mold, Cane Botrytis, Botrytis Fruit Rot and blight, Gray mold leaf blight and dry crown rot, Postharvest rots, Botrytis crown rot[*] (<i>Botrytis cinerea</i>)</p> <p>Fruit rots, minor, Berry rots, Fruit rots (in addition to those appearing elsewhere in this listing) [*] (<i>Cladosporium spp.</i>)</p> <p>Anthracoese Leaf and Fruit Rot[*] (<i>Colletotrichum sp.</i>)</p> <p>Root rot[*] (<i>Cylindrocarpon destructans</i>)</p> <p>Powdery Mildew[*] (<i>Sphaerotheca macularis</i>)</p> <p>Hard brown rot, Rhizoctonia bud and crown rot, leaf blight, web blight, fruit rot[*] (<i>Rhizoctonia solani</i>)</p>	<p>6.2 oz/A (0.7 oz ai/A)</p>	<ul style="list-style-type: none"> • Apply as a foliar spray every 7 – 10 days for foliar diseases. • Apply prior to disease development and when conditions are conducive for disease. • Mix spray volume appropriate for sufficient coverage. • 0 day Preharvest Interval. • Do not apply by air. • Use of adjuvant may enhance coverage. • Apply as a drench every 14 - 28 days for crown and root rot diseases.
<p>Disease Suppression</p>		<ul style="list-style-type: none"> • For Powdery Mildew suppression, begin sprays preventively and continue as needed on a 14 day interval.
<p>Powdery Mildew[*] (<i>Uncinula necator</i>)</p>		

Notes:

- Use in alternation with fungicides that have different modes of action.
- Do not apply more than 3 applications of ENDORSE Water Dispersible Granules (2.1 oz ai/A) per season.

[* Not for use in California.]

CUCURBIT VEGETABLES[*]		
<p>Chayote (fruit); Chinese waxgourd (Chinese preserving melon); Citron melon; Cucumber; Gherkin; Gourd, edible (includes hyotan, cucuzza, hechima, Chinese okra); <i>Momordica</i> spp. (includes balsam apple, balsam pear, bitter melon, Chinese cucumber); Muskmelon (includes true cantaloupe, cantaloupe, casaba, crenshaw melon, golden pershaw melon, honeydew melon, honey balls, mango melon, Persian melon, pineapple melon, Santa Claus melon, and snake melon); Pumpkin; Summer Squash (includes crookneck squash, scallop squash, straightneck squash, vegetable marrow, zucchini); Winter Squash (includes butternut squash, calabaza, hubbard squash, acorn squash, spaghetti squash); Watermelon</p>		
Disease Control	Rate	Application Notes
<p>Powdery Mildew[*] (<i>Sphaerotheca</i> sp.)</p> <p>Gummy Stem Blight (Vine decline)[*] (<i>Didymella bryoniae</i>)</p> <p>Gray Mold[*] (<i>Botrytis</i> sp.)</p> <p>Corynespora leaf spot[*] (<i>Corynespora cassicola</i>)</p> <p>Scab[*] (<i>Gladosporium</i> sp.)</p> <p>Alternaria leaf spot[*] (<i>Alternaria alternata</i>)</p> <p>Early Blight[*] (<i>Alternaria</i> sp.)</p> <p>Belly rot, Damping-off and Web blight[*] (<i>Rhizoctonia solani</i>)</p>	<p>6.2 oz/A (0.7 oz ai/A)</p>	<ul style="list-style-type: none"> • Apply as a foliar spray every 7-10 days for foliar diseases. • Mix spray volume appropriate for sufficient coverage. • Apply prior to disease development and when conditions are conducive for disease. • 0 day Preharvest Interval.
<p>Notes:</p> <ul style="list-style-type: none"> • Use in alternation with fungicides that have different modes of action. • Do not apply more than 5 applications of ENDORSE Water Dispensible Granules (3.5 oz ai/A) per season. <p>[* Not for use in California.]</p>		

FRUITING VEGETABLES (Except Cucurbits)[*]		
Eggplant; Groundcherry; Pepino; Pepper (includes bell pepper, chili pepper, cooking pepper, pimento, sweet pepper); Tomatillo; Tomato		
Disease Control	Rate	Application Notes
Powdery Mildew[*] <i>(Leveillula taurica, Oidiopsis sipula)</i> Botrytis Rot[*] <i>(Botrytis sp.)</i> Early Blight[*] <i>(Alternaria solani)</i> Alternaria stem canker, Black mold rot, Black shoulder[*] <i>(Alternaria alternata)</i> Rhizoctonia damping-off and fruit rot[*] <i>(Rhizoctonia solani)</i>	6.2 oz/A (0.7 oz ai/A)	<ul style="list-style-type: none"> • Apply as a foliar spray every 7 - 10 days for foliar diseases. • Apply prior to disease development and when conditions are conducive for disease. • Mix spray volume appropriate for sufficient coverage. • 0 day Preharvest Interval.
Disease Suppression		
Anthracnose[*] <i>(Colletotrichum coccodes)</i>		
Notes: <ul style="list-style-type: none"> • Use in alternation with fungicides that have different modes of action. • Do not apply more than 5 applications of ENDORSE Water Dispersible Granules (3.5 oz ai/A) per season. [* Not for use in California.]		

POME FRUITS[*]		
Apple; crabapple; mayhaw; pear, Asian; quince; quince, Chinese; quince, Japanese; European pear		
Disease Control	Rate	Application Notes
Powdery Mildew[*] <i>(Podosphaera leucotricha</i> <i>Phyllactinia mali</i> <i>Oidium spp.)</i> Alternaria Leaf Spot, Alternaria Blotch[*] <i>(Alternaria mali)</i> Moldy core and core rot, Alternaria fruit rot[*] <i>(Alternaria spp)</i> Alternaria rot, Alternaria spot and fruit rot, Black spot (of Japanese pear)[*] <i>(Alternaria alternata)</i> Dry eye rot, blossom-end rot, Green fruit rot, Botrytis spur and blossom blight, Gray mold rot[*] <i>(Botrytis cinerea)</i> Moldy core and core rot, Ripe fruit rot[*] <i>(Cladosporium spp.)</i> Blotch Marssonina blotch[*] <i>(Diplocarpon mali)</i>	6.2 oz/A (0.7 oz ai/A)	<ul style="list-style-type: none"> • Mix spray volume appropriate for sufficient coverage. • Apply as a foliar spray every 7 - 10 days for foliar diseases. • Apply prior to disease development and when conditions are conducive for disease. Start sprays for scab control at green tip and continue as needed on a 7 - 10 day interval. • For Powdery Mildew control, begin preventively and repeat as necessary on a 10 - 14 day interval. The use of an alternation program with a sterol inhibiting fungicide is recommended. • 0 day Preharvest Interval.
Disease Suppression		
Scab[*] <i>(Venturia sp.)</i> Apple scab suppression[*] <i>(Venturia inaequalis)</i>		
Notes: <ul style="list-style-type: none"> • Use in alternation with fungicides that have different modes of action. • Do not apply more than 6 applications of ENDORSE Water Dispersible Granules (4.2 oz ai/A) per season. [* Not for use in California.]		

ROOT AND TUBER VEGETABLES[*]		
Carrot; ginseng*; potato		
Disease Control	Rate	Application Notes
<p>Alternaria blight[*] (<i>Alternaria panax</i>)</p> <p>Botrytis blight, Gray mold rot, Watery soft rot, Gray mold[*] (<i>Botrytis cinerea</i>)</p> <p>Early Blight (Potato)[*] (<i>Alternaria solani</i>)</p> <p>Alternaria leaf spot, Brown spot, Black pit[*] (<i>Alternaria alternata</i>)</p> <p>Damping-off, black leg, black root and seedling blight, Canker, Cavity spot, Crown rot, Rhizoctonia canker, and Black scurf[*] (<i>Rhizoctonia solani</i>)</p>	<p>6.2 oz/A (0.7 oz ai/A)</p>	<ul style="list-style-type: none"> • Apply as a foliar spray every 7 - 10 days for foliar diseases. • Apply prior to disease development and when conditions are conducive for disease. • Mix spray volume appropriate for sufficient coverage. • 0 day Preharvest Interval.
Root and Crown Disease Control (Ginseng)		Additional Application Notes for Ginseng
<p>Cylindrocarpon root rot. (<i>Cylindrocarpon destructans</i>)</p> <p>Rhizoctonia root and crown rot (<i>Rhizoctonia solani</i>)</p>		<ul style="list-style-type: none"> • Apply as a drench every 14 days for Cylindrocarpon rot control. • Apply as a drench every 14 – 28 days for root and crown rot diseases.
<p>Notes:</p> <ul style="list-style-type: none"> • For use on Ginseng: <ul style="list-style-type: none"> • Apply within 2 weeks following plant emergence and continue throughout the season. • Consult your local extension agent or local recommendations for information relating to proper timing for control of these diseases. • Use in alternation with fungicides that have different modes of action. • Do not apply more than 3 applications of ENDORSE Water Dispersible Granules (2.1 oz ai/A) per season. <p>* Not for use in California.</p>		

LEAVES OF ROOT AND TUBER VEGETABLES (HUMAN FOOD OR ANIMAL FEED)[*] Carrot		
Disease Control	Rate	Application Notes
Alternaria leaf spot and stem blight, Alternaria storage rot[*] <i>(Alternaria spp.)</i> Gray mold rot[*] <i>(Botrytis cinerea)</i> Rhizoctonia stem canker (sprout rot), Rootlet rot[*] <i>(Rhizoctonia solani)</i>	6.2 oz/A (0.7 oz ai/A)	<ul style="list-style-type: none"> • Apply as a foliar spray every 7 - 10 days for foliar diseases. • Apply prior to disease development and when conditions are conducive for disease. • Mix spray volume appropriate for sufficient coverage. • 0 day Preharvest Interval. • Apply as a drench every 14 - 28 days for crown and root rot diseases.
Notes: <ul style="list-style-type: none"> • Use in alternation with fungicides that have different modes of action. • Do not apply more than 5 applications of ENDORSE Water Dispersible Granules (3.5 oz ai/A) per season. [*Not for use in California.]		

GRAPES		
Disease Control	Rate	Application Notes
<p>Alternaria rot, Black leaf spot[*] (<i>Alternaria alternata</i>)</p> <p>Fruit rots, minor[*] (<i>Alternaria spp.</i>)</p> <p>Botrytis Fruit Rot, Botrytis Bunch Rot (<i>Botrytis cinerea</i>)</p> <p>Botrytis fruit rot and blossom blight, gray mold, Cane Botrytis, and blight, Gray mold leaf blight and dry crown rot, Postharvest rots, Botrytis crown rot[*] (<i>Botrytis cinerea</i>)</p> <p>Fruit rots, minor, Berry rots and raisin molds, Fruit rots (in addition to those appearing elsewhere in this listing)[*] (<i>Cladosporium spp.</i>)</p> <p>Anthracnose Leaf and Fruit Rot[*] (<i>Colletotrichum sp.</i>)</p> <p>Root rot[*] (<i>Cylindrocarpon destructans</i>)</p> <p>Hard brown rot, Rhizoctonia bud and crown rot, leaf blight, web blight, fruit rot[*] (<i>Rhizoctonia solani</i>)</p>	<p>6.2 oz/A (0.7 oz ai/A)</p>	<ul style="list-style-type: none"> • Apply as a foliar spray every 7 – 10 days for foliar diseases. • Apply prior to disease development and when conditions are conducive for disease. • Mix spray volume appropriate for sufficient coverage. • For Botrytis Bunch Rot control, spray product at veraison and 7 days prior to harvest. • 0 day Preharvest Interval. • Do not apply by air. • Use of adjuvant may enhance coverage. • Apply as a drench every 14 - 28 days for crown and root rot diseases.
Disease Suppression		<ul style="list-style-type: none"> • For Powdery Mildew suppression. Begin sprays preventively and continue as needed on a 14 day interval.
<p>Powdery Mildew[*] (<i>Erysiphe (Uncinula) necator</i>), (<i>Oidium tuckeri</i>)</p>		
<p>Notes</p> <ul style="list-style-type: none"> • Use in alternation with fungicides that have different modes of action. • Do not apply more than 3 applications of ENDORSE Water Dispersible Granules (2.1 oz ai/A) per season. <p>[*Not for use in California.]</p>		

STONE FRUITS[*]		
Sweet cherry or Tart cherry; Peach, and Plum or Prune plum; apricot; apricot, Japanese; capulin; cherry, black; cherry, Nanking; Jujube, Chinese; nectarine; peach; plum; plum, American; plum, beach; plum, Canada; plum, cherry; plum, Chickasaw; plum, Damson; plum, Japanese; plum, Klamath; plum, prune; plumcot; sloe; cultivars, varieties, and/or hybrids of these		
Disease Control	Rate	Application Notes
Alternaria leaf spot, Alternaria rot[*] <i>(Alternaria alternata)</i> Alternaria rot[*] <i>(Alternaria spp.)</i> Gray mold rot, Green fruit rot, Botrytis rot[*] <i>(Botrytis cinerea)</i> Scab[*] <i>(Cladosporium spp.)</i> Powdery Mildew[*] <i>(Oidium spp.)</i> Powdery Mildew[*] <i>(Podosphaera leucotricha)</i> Rhizoctonia root rot[*] <i>(Rhizoctonia solani)</i>	6.2 oz/A (0.7 oz ai/A)	<ul style="list-style-type: none"> • Apply as a foliar spray every 7 – 10 days for foliar diseases. • Apply prior to disease development and when conditions are conducive for disease. • Mix spray volume appropriate for sufficient coverage. • 0 day Preharvest Interval. • Apply as a drench every 14 – 28 days for crown and root rot diseases.
[*Not for use in California.]		

POMEGRANATES[*]		
Disease Control	Rate	Application Notes
Alternaria fruit rot[*] <i>(Alternaria spp.)</i> Botrytis fruit rot (Gray mold)[*]	6.2 oz/A (0.7 oz ai/A)	<ul style="list-style-type: none"> • Apply as a foliar spray every 7 - 10 days for foliar diseases. • Apply when conditions are conducive for disease. Beginning during bloom through preharvest. • Mix spray volume appropriate for sufficient coverage. • 0 day Preharvest Interval.
Notes: <ul style="list-style-type: none"> • Use in alternation with fungicides that have different modes of action. • Do not apply more than 3 applications of ENDORSE Water Dispersible Granules (2.1 oz ai/A) per season. [*Not for use in California.]		

TREE NUTS[*]		
almond; pecan; African nut-tree; beechnut; Brazil nut; Brazilian pine; bunya; bur oak; butternut; Cajou nut; candlenut; cashew; chestnut; chinquapin; coconut; coquito nut; dika nut; ginkgo; Guiana chestnut; hazelnut (filbert); heartnut; hickory nut; Japanese horse-chestnut; macadamia nut; mongongo nut; monkey-pot; monkey puzzle nut; Okari nut; Pachira nut; peach palm nut; pequi; Pili nut; pine nut; pistachio; Sapucaia nut; tropical almond; walnut, black; walnut, English; yellowhorn; cultivars, varieties, and/or hybrids of these		
Disease Control	Rate	Application Notes
Alternaria, Alternaria late blight[*] <i>(Alternaria alternata)</i> Leaf spots[*] <i>(Alternaria spp.)</i> Blossom and shoot blight[*] <i>(Botrytis cinerea)</i> Cylindrocarpon dieback[*] <i>Cylindrocarpon destructans</i> Powdery mildew[*] <i>(Oidium spp.)</i> Damping-off[*] Root rot[*] Seedling blight[*] <i>(Rhizoctonia solani)</i>	6.2 oz/A (0.7 oz ai/A)	<ul style="list-style-type: none"> • Mix spray volume appropriate for sufficient coverage (Spray every row to ensure coverage [avoid every other row applications]). • Apply as a foliar spray every 7-10 days for foliar diseases. • Apply prior to disease development and when conditions are conducive for disease. • 0 day Preharvest Interval. • May be applied by air or ground. EXCEPT Pistachios. Do not apply to Pistachios by air. • Use of an adjuvant may enhance coverage.
Hull rot[*] <i>(Rhizopus stolonifer)</i>	6.2 oz/A (0.7 oz ai/A)	For control of Hull rot caused by <i>Rhizopus stolonifer</i> apply Endorse Water Dispersible Granules at the rate of 6.2 oz/A in a dilute spray beginning at 10% hull split. Follow up with a second application at 20 – 40% hull split, if necessary. Mix in appropriate spray volume appropriate for sufficient coverage (Spray every row to ensure coverage [avoid every other row applications]). <ul style="list-style-type: none"> • Begin sprays preventively and repeat on a 10–14 day interval. • 0 day Preharvest Interval. • May be applied by air or ground. • Use of an adjuvant may enhance coverage.

TREE NUTS[*]		
almond; pecan; African nut-tree; beechnut; Brazil nut; Brazilian pine; bunya; bur oak; butternut; Cajou nut; candlenut; cashew; chestnut; chinquapin; coconut; coquito nut; dika nut; ginkgo; Guiana chestnut; hazelnut (filbert); heartnut; hickory nut; Japanese horse-chestnut; macadamia nut; mongongo nut; monkey-pot; monkey puzzle nut; Okari nut; Pachira nut; peach palm nut; pequi; Pili nut; pine nut; pistachio; Sapucaia nut; tropical almond; walnut, black; walnut, English; yellowhorn; cultivars, varieties, and/or hybrids of these		
Disease Control	Rate	Application Notes
Scab[*] (<i>Cladosporium carpophilum</i>)	6.2 oz/A (0.7 oz ai/A)	For control of Almond scab apply Endorse Water Dispersible Granules at the rate of 6.2 oz/A in a dilute spray when infection periods begin. Retreatment with Endorse Water Dispersible Granules or other effective fungicides should be made be on 10 – 14 day intervals throughout the season as disease infection periods occur. Mix in appropriate spray volume appropriate for sufficient coverage (Spray every row to ensure coverage [avoid every other row applications]). <ul style="list-style-type: none"> • Begin sprays preventively and repeat on a 10–14 day interval. • 0 day Preharvest Interval. • May be applied by air or ground. • Use of an adjuvant may enhance coverage.
Rust[*] (<i>Tranzchelia discolor</i>)	6.2 oz/A (0.7 oz ai/A)	For control of Almond rust apply Endorse Water Dispersible Granules at the rate of 6.2 oz/A in a dilute spray when infection periods begin. Retreatment with Endorse Water Dispersible Granules or other effective fungicides should be made on 14 - 21 day intervals throughout the season as disease infection periods occur. Mix in appropriate spray volume appropriate for sufficient coverage (Spray every row to ensure coverage [avoid every other row applications]). <ul style="list-style-type: none"> • Begin sprays preventively and repeat on a 10–14 day interval. • 0 day Preharvest Interval. • May be applied by air or ground. • Use of an adjuvant may enhance coverage.
Disease Suppression		
Botryosphaeria blight[*] (<i>Botryosphaeria</i> sp.)	6.2 oz/A (0.7 oz ai/A)	
Notes: <ul style="list-style-type: none"> • Use in alternation with fungicides that have different modes of action. • Do not apply more than 3 applications of ENDORSE Water Dispersible Granules (2.1 oz ai/A) per season. [*Not for use in California.]		

ORNAMENTALS*, ¹		
Foliar Diseases	Rate	Application Notes
Anthracnose (<i>Colletotrichum</i>) †Botrytis blight (<i>Botrytis cinerea</i>) Curvularia Downy Mildew (<i>Pernospora</i> sp. <i>Plasmopara</i> sp.) Powdery Mildew (<i>Oidium</i> sp. <i>Erysiphe</i> sp. <i>Sphaerotheca</i> sp.) Rhizoctonia aerial blight (<i>Rhizoctonia solani</i>)	0.25 – 0.5 lb/100 gal/A (0.03 – 0.06 lb ai/100 gal/A)	<ul style="list-style-type: none"> • Apply as a foliar spray every 7 – 10 days for foliar diseases. • Apply prior to disease development and when conditions are conducive to disease development. • Mix spray volume appropriate for sufficient coverage.
†Alternaria blight (<i>Alternaria panax</i>) Apple scab (<i>Venturia inaequalis</i>)	0.5 lb/100 gal/A (0.06 lb ai/100 gal/A)	
Root and Crown Diseases	Rate	Application Notes
*Rhizoctonia root and crown rot (<i>Rhizoctonia solani</i>) Black Root Rot (<i>Thielaviopsis basicola</i>)	0.25 – 0.5 lb/100 gal/A (0.03 – 0.06 lb ai/100 gal/A)	<ul style="list-style-type: none"> • Apply as a drench every 14 – 28 days for crown and root rot diseases.
Notes: † Control Consult your local extension agent or local recommendations for information relating to proper timing for control of these diseases. <ul style="list-style-type: none"> • Use in alternation with fungicides that have different modes of action. ¹ Not for use in California. *Includes Christmas trees (nursery and plantations) and non-bearing vine fruit and fruit and nut trees.		

TURF GRASSES ²				
Disease	Rate	Application Notes		
Control	2.4 lb/A (0.27 lb ai/A in a minimum of 88 gal of water)	<ul style="list-style-type: none"> Repeat treatment on a 7-14 day schedule where environmental conditions are conducive to development of diseases.¹ When symptoms are present, best control will be achieved by using a shorter interval. For best results apply after mowing. 		
Brown Patch Large Patch				
Aids in Control (Except California)				
Cool Weather Brown Patch (Yellow Patch) (<i>Rhizoctonia cerealis</i>)				
Foliar and Basal Anthracnose (<i>Colletotrichum graminicola</i>)				
Gray Snow Mold (<i>Typhula ishikariensis</i> and <i>Typhula incarnate</i>)	2.4 lb/A (0.27 lb ai/A in a minimum of 88 gal of water)	<ul style="list-style-type: none"> Repeat treatment on a 7-14 day schedule where environmental conditions are conducive to development of diseases.¹ When symptoms are present, best control will be achieved by using a shorter interval. For best results apply after mowing. 		
Leaf Spot/Melting Out (<i>Dreschlera poae</i>)				
Pink Snow Mold (<i>Microdochium solani</i>)				
Red Thread (<i>Laetisaria fuciformis</i>)				
Rhizoctonia Damping Off (<i>Rhizoctonia solani</i>)				
Zoysia Patch (<i>Rhizoctonia solani</i>)				
Aids in Suppression (Except California)				
Gray Leaf Spot (<i>Pyricularia grisea</i>)				
Suppression and Short Term Control:			1 oz/1,000 sq ft in minimum of 2 gal of water (0.27 lb ai/A in 88 gal of water)	<ul style="list-style-type: none"> Make two-three applications on a 7-day interval schedule. Use a penetrating wetting agent. Immediately following application, water in the treatment with sufficient irrigation (0.5-1") to wet the active root zone.
Fairy Ring (including <i>Marasmius</i> spp., <i>Lepiota</i> spp. and <i>Agaricus</i> spp.)				
<p>Notes:</p> <ul style="list-style-type: none"> With the exception for treatment of Fairy Ring, irrigation or rainfall soon after treatment with ENDORSE Water Dispersible Granules will decrease disease control. For best control, do not irrigate for 12 hours after treatment. <p>¹Consult your local extension agent or local recommendations for information relating to proper timing for control of these diseases.</p> <ul style="list-style-type: none"> Use in alternation with fungicides that have different modes of action. <p>² Not for use in California.</p>				

18/30

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in dry place away from food or feed.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed, by State and local authorities, by burning. If burned, stay out of smoke. Offer for recycling, if available.

**FOR 24-HOUR CHEMICAL EMERGENCY (Spill, Leak, Fire or Accident) ASSISTANCE:
CALL CHEMTREC 1-800-424-9300**

Warranty and Disclaimer Statement

The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Such risks may arise from weather conditions, soil factors, off-target movement, unconventional farming techniques, the presence of other materials, the manner of use or application, or other unknown factors, all of which are beyond the control of Arysta LifeScience North America, LLC ("Arysta"), and can cause crop injury, injury to non-target crops or plants, ineffectiveness of the product, or other unintended consequences. To the extent consistent with applicable law, all such risks shall be assumed by the user or buyer.

Arysta warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks described above, when used in accordance with the Directions for Use under normal conditions.

This warranty does not extend to the use of this product contrary to label instructions or under conditions not reasonably foreseeable to Arysta, and is subject to the inherent risks described above.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, ARYSTA DISCLAIMS ALL OTHER WARRANTIES, EXPRESSED OR IMPLIED, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, ARYSTA, MANUFACTURER, AND SELLER DISCLAIM AND SHALL NOT BE LIABLE FOR ANY SPECIAL, INCIDENTAL, INDIRECT, OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE, HANDLING, APPLICATION, STORAGE, OR DISPOSAL OF THIS PRODUCT OR FOR DAMAGES IN THE NATURE OF PENALTIES, AND THE USER AND BUYER WAIVE ANY RIGHT THAT THEY MAY HAVE TO SUCH DAMAGES. NO AGENT, REPRESENTATIVE OR EMPLOYEE OF ARYSTA IS AUTHORIZED TO MAKE ANY WARRANTY, GUARANTEE OR REPRESENTATION BEYOND THOSE CONTAINED HEREIN OR TO MODIFY THE WARRANTIES CONTAINED HEREIN.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE TOTAL LIABILITY OF ARYSTA, MANUFACTURER, AND SELLER, SHALL BE LIMITED TO THE PURCHASE PRICE PAID, OR AT ARYSTA'S ELECTION, THE REPLACEMENT OF THE PRODUCT.

ENDORSE® is a registered trademark of Arysta LifeScience North America, LLC.

ENDORSE WDG (PENDING) 03/10/14

20/30

[TEXT IN BRACKETS IS OPTIONAL]



Arysta LifeScience

SUPPLEMENTAL LABELING

Ph-D[®]

Fungicide

For use on listed crops

GROUP 19 FUNGICIDE

This supplemental label expires on Month day, yyyy and must not be used or distributed after this date

INGREDIENTS

By wt.

Active Ingredient:	
Polyoxin D zinc salt.....	11.3%
Other Ingredients:	88.7%
Total:	100.0%

KEEP OUT OF REACH OF CHILDREN

CAUTION

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Follow all applicable directions, restrictions, worker protection standard requirements, and precautions on the EPA registered main label for Ph-D Fungicide (EPA Reg. No. 66330-56).

This supplemental label must be in the possession of the user at the time of application.

USE INFORMATION

Ph-D Fungicide is not for homeowner use to treat food crops.

Ph-D Fungicide is for use in controlling or suppressing certain diseases on crops listed on this label. See Container label for more approved uses.

Apply Ph-D Fungicide as a preventive or curative treatment in conjunction with good management practices.

ACCEPTED

JUN 05 2014

Under the Federal Insecticide, Fungicide,
and Rodenticide Act, as amended, for
the pesticide registered under
EPA Reg. No. 66330-56

STRAWBERRIES[*]

BERRY AND SMALL FRUIT[*]

Amur river grape; aronia berry; bayberry; bearberry; bilberry; blackberry (including Andean blackberry, arctic blackberry, bingleberry, black satin berry, boysenberry, brombeere, California blackberry, Chesterberry, Cherokee blackberry, Cheyenne blackberry, common blackberry, coryberry, darrowberry, dewberry, Dirksen thornless berry, evergreen blackberry, Himalayaberry, hullberry, lavacaberry, loganberry, lowberry, Lucretiaberry, mammoth blackberry, marionberry, mora, mures deronce, nectarberry, Northern dewberry, olallieberry, Orgeon evergreen berry, phenomenalberry, rangeberry, ravenberry, rossberry, Shawnee blackberry, Southern dewberry, tayberry, youngberry, zarzamora, and cultivars, varieties and/or hybrids of these); blueberry, highbush; blueberry, lowbush; buffalo currant; buffaloberry; che; Chilean guava; chokecherry; cloudberry; cranberry; cranberry, highbush; currant, black; currant, red; elderberry; European barberry; gooseberry; honeysuckle, edible; huckleberry; jostaberry; Juneberry (Saskatoon berry); kiwifruit, fuzzy; kiwifruit, hardy; lingonberry; maypop; mountain pepper berries; mulberry; muntries; native currant; partridgeberry; phalsa; pincherry; raspberry, black and red; riberry; salal; schisandra berry; sea buckthorn; serviceberry; strawberry; wild raspberry; cultivars, varieties, and/or hybrids of these

Disease Control	Rate	Application Notes
<p>Alternaria rot, Black leaf spot[*] (<i>Alternaria alternata</i>)</p> <p>Fruit rots, minor[*] (<i>Alternaria spp.</i>)</p> <p>Botrytis fruit rot and blossom blight, gray mold, Cane Botrytis, Botrytis Fruit Rot and blight, Gray mold leaf blight and dry crown rot, Postharvest rots, Botrytis crown rot[*] (<i>Botrytis cinerea</i>)</p> <p>Fruit rots, minor, Berry rots, Fruit rots (in addition to those appearing elsewhere in this listing) [*] (<i>Cladosporium spp.</i>)</p> <p>Anthraxnose Leaf and Fruit Rot[*] (<i>Colletotrichum sp.</i>)</p> <p>Root rot[*] (<i>Cylindrocarpon destructans</i>)</p> <p>Powdery Mildew[*] (<i>Sphaerotheca macularis</i>)</p> <p>Hard brown rot, Rhizoctonia bud and crown rot, leaf blight, web blight, fruit rot[*] (<i>Rhizoctonia solani</i>)</p>	<p>6.2 oz/A (0.7 oz ai/A)</p>	<ul style="list-style-type: none"> • Apply as a foliar spray every 7 – 10 days for foliar diseases. • Apply prior to disease development and when conditions are conducive for disease. • Mix spray volume appropriate for sufficient coverage. • 0 day Preharvest Interval. • Do not apply by air. • Use of adjuvant may enhance coverage. • Apply as a drench every 14 - 28 days for crown and root rot diseases.
<p>Disease Suppression</p>		
<p>Powdery Mildew[*] (<i>Uncinula necator</i>)</p>		<ul style="list-style-type: none"> • For Powdery Mildew suppression, begin sprays preventively and continue as needed on a 14 day interval.

Notes:

- Use in alternation with fungicides that have different modes of action.
 - Do not apply more than 3 applications of Ph-D Fungicide (2.1 oz ai/A) per season.
- [* Not for use in California.]

CUCURBIT VEGETABLES[*]

Chayote (fruit); Chinese waxgourd (Chinese preserving melon); Citron melon; Cucumber; Gherkin; Gourd, edible (includes hyotan, cucuzza, hechima, Chinese okra); *Momordica* spp. (includes balsam apple, balsam pear, bitter melon, Chinese cucumber); Muskmelon (includes true cantaloupe, cantaloupe, casaba, crenshaw melon, golden pershaw melon, honeydew melon, honey balls, mango melon, Persian melon, pineapple melon, Santa Claus melon, and snake melon); Pumpkin; Summer Squash (includes crookneck squash, scallop squash, straightneck squash, vegetable marrow, zucchini); Winter Squash (includes butternut squash, calabaza, hubbard squash, acorn squash, spaghetti squash); Watermelon

Disease Control	Rate	Application Notes
Powdery Mildew[*] (<i>Sphaerotheca</i> sp.) Gummy Stem Blight (Vine decline)[*] (<i>Didymella bryoniae</i>) Gray Mold[*] (<i>Botrytis</i> sp.) Corynespora leaf spot[*] (<i>Corynespora cassicola</i>) Scab[*] (<i>Cladosporium</i> sp.) Alternaria leaf spot[*] (<i>Alternaria alternata</i>) Early Blight[*] (<i>Alternaria</i> sp.) Belly rot, Damping-off and Web blight[*] (<i>Rhizoctonia solani</i>)	6.2 oz/A (0.7 oz ai/A)	<ul style="list-style-type: none"> • Apply as a foliar spray every 7-10 days for foliar diseases. • Mix spray volume appropriate for sufficient coverage. • Apply prior to disease development and when conditions are conducive for disease. • 0 day Preharvest Interval.

Notes:

- Use in alternation with fungicides that have different modes of action.
- Do not apply more than 5 applications of Ph-D Fungicide (3.5 oz ai/A) per season.

[* Not for use in California.]

FRUITING VEGETABLES (Except Cucurbits)[*]		
Eggplant; Groundcherry; Pepino; Pepper (includes bell pepper, chili pepper, cooking pepper, pimento, sweet pepper); Tomatillo; Tomato		
Disease Control	Rate	Application Notes
Powdery Mildew[*] <i>(Leveillula taurica, Oidiopsis sipula)</i> Botrytis Rot[*] <i>(Botrytis sp.)</i> Early Blight[*] <i>(Alternaria solani)</i> Alternaria stem canker, Black mold rot, Black shoulder[*] <i>(Alternaria alternata)</i> Rhizoctonia damping-off and fruit rot[*] <i>(Rhizoctonia solani)</i>	6.2 oz/A (0.7 oz ai/A)	<ul style="list-style-type: none"> • Apply as a foliar spray every 7 - 10 days for foliar diseases. • Apply prior to disease development and when conditions are conducive for disease. • Mix spray volume appropriate for sufficient coverage. • 0 day Preharvest Interval.
Disease Suppression		
Anthracnose[*] <i>(Colletotrichum coccodes)</i>		
Notes: <ul style="list-style-type: none"> • Use in alternation with fungicides that have different modes of action. • Do not apply more than 5 applications of Ph-D Fungicide (3.5 oz ai/A) per season. [* Not for use in California.]		

POME FRUITS [*]		
Apple; crabapple; mayhaw; pear, Asian; quince; quince, Chinese; quince, Japanese; European pear		
Disease Control	Rate	Application Notes
Powdery Mildew[*] <i>(Podosphaera leucotricha</i> <i>Phyllactinia mali</i> <i>Oidium spp.)</i> Alternaria Leaf Spot, Alternaria Blotch[*] <i>(Alternaria mali)</i> Moldy core and core rot, Alternaria fruit rot[*] <i>(Alternaria spp)</i> Alternaria rot, Alternaria spot and fruit rot, Black spot (of Japanese pear)[*] <i>(Alternaria alternata)</i> Dry eye rot, blossom-end rot, Green fruit rot, Botrytis spur and blossom blight, Gray mold rot[*] <i>(Botrytis cinerea)</i> Moldy core and core rot, Ripe fruit rot[*] <i>(Cladosporium spp.)</i> Blotch Marssonina blotch[*] <i>(Diplocarpon mali)</i>	6.2 oz/A (0.7 oz ai/A)	<ul style="list-style-type: none"> • Mix spray volume appropriate for sufficient coverage. • Apply as a foliar spray every 7 - 10 days for foliar diseases. • Apply prior to disease development and when conditions are conducive for disease. Start sprays for scab control at green tip and continue as needed on a 7 - 10 day interval. • For Powdery Mildew control, begin preventively and repeat as necessary on a 10 - 14 day interval. The use of an alternation program with a sterol inhibiting fungicide is recommended. • 0 day Preharvest Interval.
Disease Suppression		
Scab[*] <i>(Venturia sp.)</i> Apple scab suppression[*] <i>(Venturia inaequalis)</i>		
Notes: <ul style="list-style-type: none"> • Use in alternation with fungicides that have different modes of action. • Do not apply more than 6 applications of Ph-D Fungicide (4.2 oz ai/A) per season. [* Not for use in California.]		

ROOT AND TUBER VEGETABLES[*]		
Carrot; ginseng*; potato		
Disease Control	Rate	Application Notes
Alternaria blight[*] <i>(Alternaria panax)</i> Botrytis blight, Gray mold rot, Watery soft rot, Gray mold[*] <i>(Botrytis cinerea)</i> Early Blight (Potato)[*] <i>(Alternaria solani)</i> Alternaria leaf spot, Brown spot, Black pit[*] <i>(Alternaria alternata)</i> Damping-off, black leg, black root and seedling blight, Canker, Cavity spot, Crown rot, Rhizoctonia canker, and Black scurf[*] <i>(Rhizoctonia solani)</i>	6.2 oz/A (0.7 oz ai/A)	<ul style="list-style-type: none"> Apply as a foliar spray every 7 - 10 days for foliar diseases. Apply prior to disease development and when conditions are conducive for disease. Mix spray volume appropriate for sufficient coverage. 0 day Preharvest Interval.
Root and Crown Disease Control (Ginseng) Cylindrocarpon root rot <i>(Cylindrocarpon destructans)</i> Rhizoctonia root and crown rot <i>(Rhizoctonia solani)</i>		Additional Application Notes for Ginseng <ul style="list-style-type: none"> Apply as a drench every 14 days for Cylindrocarpon rot control. Apply as a drench every 14 – 28 days for root and crown rot diseases.
Notes: <ul style="list-style-type: none"> For use on Ginseng: <ul style="list-style-type: none"> Apply within 2 weeks following plant emergence and continue throughout the season. Consult your local extension agent or local recommendations for information relating to proper timing for control of these diseases. Use in alternation with fungicides that have different modes of action. Do not apply more than 3 applications of Ph-D Fungicide (2.1 oz ai/A) per season. * Not for use in California.		

LEAVES OF ROOT AND TUBER VEGETABLES (HUMAN FOOD OR ANIMAL FEED)[*]		
Carrot		
Disease Control	Rate	Application Notes
Alternaria leaf spot and stem blight, Alternaria storage rot[*] <i>(Alternaria spp.)</i> Gray mold rot[*] <i>(Botrytis cinerea)</i> Rhizoctonia stem canker (sprout rot), Rootlet rot[*] <i>(Rhizoctonia solani)</i>	6.2 oz/A (0.7 oz ai/A)	<ul style="list-style-type: none"> • Apply as a foliar spray every 7 - 10 days for foliar diseases. • Apply prior to disease development and when conditions are conducive for disease. • Mix spray volume appropriate for sufficient coverage. • 0 day Preharvest Interval. • Apply as a drench every 14 - 28 days for crown and root rot diseases.
Notes: <ul style="list-style-type: none"> • Use in alternation with fungicides that have different modes of action. • Do not apply more than 5 applications of Ph-D Fungicide (3.5 oz ai/A) per season. [*Not for use in California.]		

Grapes		
Disease Control	Rate	Application Notes
<p>Alternaria rot, Black leaf spot[*] (<i>Alternaria alternata</i>)</p> <p>Fruit rots, minor[*] (<i>Alternaria spp.</i>)</p> <p>Botrytis Fruit Rot, Botrytis Bunch Rot and blight, (<i>Botrytis cinerea</i>)</p> <p>Botrytis fruit rot and blossom blight, gray mold, Cane Botrytis, Gray mold leaf blight and dry crown rot, Postharvest rots, Botrytis crown rot[*] (<i>Botrytis cinerea</i>)</p> <p>Fruit rots, minor, Berry rots and raisin molds, Fruit rots (in addition to those appearing elsewhere in this listing)[*] (<i>Cladosporium spp.</i>)</p> <p>Anthracnose Leaf and Fruit Rot[*] (<i>Colletotrichum sp.</i>)</p> <p>Root rot[*] (<i>Cylindrocarpon destructans</i>)</p> <p>Hard brown rot, Rhizoctonia bud and crown rot, leaf blight, web blight, fruit rot[*] (<i>Rhizoctonia solani</i>)</p>	<p>6.2 oz/A (0.7 oz ai/A)</p>	<ul style="list-style-type: none"> • Apply as a foliar spray every 7 – 10 days for foliar diseases. • Apply prior to disease development and when conditions are conducive for disease. • Mix spray volume appropriate for sufficient coverage. • For Botrytis Bunch Rot control, spray product at veraison and 7 days prior to harvest. • 0 day Preharvest Interval. • Do not apply by air. • Use of adjuvant may enhance coverage. • Apply as a drench every 14 - 28 days for crown and root rot diseases.
Disease Suppression		<ul style="list-style-type: none"> • For Powdery Mildew suppression. Begin sprays preventively and continue as needed on a 14 day interval.
<p>Powdery Mildew[*] (<i>Erysiphe (Uncinula) necator</i>) (<i>Oidium tuckeri</i>)</p>		
<p>Notes</p> <ul style="list-style-type: none"> • Use in alternation with fungicides that have different modes of action. • Do not apply more than 3 applications of Ph-D Fungicide (2.1 oz ai/A) per season. <p>[*Not for use in California.]</p>		

STONE FRUITS[*]		
Sweet cherry or Tart cherry; Peach, and Plum or Prune plum; apricot; apricot, Japanese; capulin; cherry, black; cherry, Nanking; Jujube, Chinese; nectarine; peach; plum; plum, American; plum, beach; plum, Canada; plum, cherry; plum, Chickasaw; plum, Damson; plum, Japanese; plum, Klamath; plum, prune; plumcot; sloe; cultivars, varieties, and/or hybrids of these		
Disease Control	Rate	Application Notes
Alternaria leaf spot, Alternaria rot[*] <i>(Alternaria alternata)</i> Alternaria rot[*] <i>(Alternaria spp.)</i> Gray mold rot, Green fruit rot, Botrytis rot[*] <i>(Botrytis cinerea)</i> Scab[*] <i>(Cladosporium spp.)</i> Powdery Mildew[*] <i>(Oidium spp.)</i> Powdery Mildew[*] <i>(Podosphaera leucotricha)</i> Rhizoctonia root rot[*] <i>(Rhizoctonia solani)</i>	6.2 oz/A (0.7 oz ai/A)	<ul style="list-style-type: none"> • Apply as a foliar spray every 7 – 10 days for foliar diseases. • Apply prior to disease development and when conditions are conducive for disease. • Mix spray volume appropriate for sufficient coverage. • 0 day Preharvest Interval. • Apply as a drench every 14 – 28 days for crown and root rot diseases.
[*Not for use in California.]		

POMEGRANATES[*]		
Disease Control	Rate	Application Notes
Alternaria fruit rot[*] <i>(Alternaria spp.)</i> Botrytis fruit rot (Gray mold)[*]	6.2 oz/A (0.7 oz ai/A)	<ul style="list-style-type: none"> • Apply as a foliar spray every 7 - 10 days for foliar diseases. • Apply when conditions are conducive for disease. Beginning during bloom through preharvest. • Mix spray volume appropriate for sufficient coverage of foliage, flower parts, and fruit. • 0 day Preharvest Interval.
Notes: <ul style="list-style-type: none"> • Use in alternation with fungicides that have different modes of action. • Do not apply more than 3 applications of Ph-D Fungicide (2.1 oz ai/A) per season. [*Not for use in California.]		

TREE NUTS[*]

almond; pecan; African nut-tree; beechnut; Brazil nut; Brazilian pine; bunya; bur oak; butternut; Cajou nut; candlenut; cashew; chestnut; chinquapin; coconut; coquito nut; dika nut; ginkgo; Guiana chestnut; hazelnut (filbert); heartnut; hickory nut; Japanese horse-chestnut; macadamia nut; mongongo nut; monkey-pot; monkey puzzle nut; Okari nut; Pachira nut; peach palm nut; pequi; Pili nut; pine nut; pistachio; Sapucaia nut; tropical almond; walnut, black; walnut, English; yellowhorn; cultivars, varieties, and/or hybrids of these

Disease Control	Rate	Application Notes
<p>Alternaria, Alternaria late blight[*] (<i>Alternaria alternata</i>)</p> <p>Leaf spots[*] (<i>Alternaria spp.</i>)</p> <p>Blossom and shoot blight[*] (<i>Botrytis cinerea</i>)</p> <p>Cylindrocarpon dieback[*] (<i>Cylindrocarpon destructans</i>)</p> <p>Powdery mildew[*] (<i>Oidium spp.</i>)</p> <p>Damping-off[*] Root rot[*]</p> <p>Seedling blight[*] (<i>Rhizoctonia solani</i>)</p>	<p>6.2 oz/A (0.7 oz ai/A)</p>	<ul style="list-style-type: none"> • Mix spray volume appropriate for sufficient coverage (Spray every row to ensure coverage [avoid every other row applications]). • Apply as a foliar spray every 7-10 days for foliar diseases. • Apply prior to disease development and when conditions are conducive for disease. • 0 day Preharvest Interval. • May be applied by air or ground. EXCEPT Pistachios. Do not apply to Pistachios by air. • Use of an adjuvant may enhance coverage.
<p>Hull rot[*] (<i>Rhizopus stolonifer</i>)</p>	<p>6.2 oz/A (0.7 oz ai/A)</p>	<p>For control of Hull rot caused by <i>Rhizopus stolonifer</i> apply Ph-D Fungicide at the rate of 6.2 oz/A in a dilute spray beginning at 10% hull split. Follow up with a second application at 20 – 40% hull split, if necessary. Mix in appropriate spray volume appropriate for sufficient coverage (Spray every row to ensure coverage [avoid every other row applications]).</p> <ul style="list-style-type: none"> • Begin sprays preventively and repeat on a 10–14 day interval. • 0 day Preharvest Interval. • May be applied by air or ground. • Use of an adjuvant may enhance coverage.

TREE NUTS[*]		
almond; pecan; African nut-tree; beechnut; Brazil nut; Brazilian pine; bunya; bur oak; butternut; Cajou nut; candlenut; cashew; chestnut; chinquapin; coconut; coquito nut; dika nut; ginkgo; Guiana chestnut; hazelnut (filbert); heartnut; hickory nut; Japanese horse-chestnut; macadamia nut; mongongo nut; monkey-pot; monkey puzzle nut; Okari nut; Pachira nut; peach palm nut; pequi; Pili nut; pine nut; pistachio; Sapucaia nut; tropical almond; walnut, black; walnut, English; yellowhorn; cultivars, varieties, and/or hybrids of these		
Disease Control	Rate	Application Notes
Scab[*] (<i>Cladosporium carpophilum</i>)	6.2 oz/A (0.7 oz ai/A)	For control of Almond scab apply Ph-D Fungicide at the rate of 6.2 oz/A in a dilute spray when infection periods begin. Retreatment with Ph-D Fungicide or other effective fungicides should be made be on 10 – 14 day intervals throughout the season as disease infection periods occur. Mix in appropriate spray volume appropriate for sufficient coverage (Spray every row to ensure coverage [avoid every other row applications]). <ul style="list-style-type: none"> • Begin sprays preventively and repeat on a 10–14 day interval. • 0 day Preharvest Interval. • May be applied by air or ground. • Use of an adjuvant may enhance coverage.
Rust[*] (<i>Tranzchelia discolor</i>)	6.2 oz/A (0.7 oz ai/A)	For control of Almond rust apply Ph-D Fungicide at the rate of 6.2 oz/A in a dilute spray when infection periods begin. Retreatment with Ph-D Fungicide or other effective fungicides should be made on 14 - 21 day intervals throughout the season as disease infection periods occur. Mix in appropriate spray volume appropriate for sufficient coverage (Spray every row to ensure coverage [avoid every other row applications]). <ul style="list-style-type: none"> • Begin sprays preventively and repeat on a 10–14 day interval. • 0 day Preharvest Interval. • May be applied by air or ground. • Use of an adjuvant may enhance coverage.
Disease Suppression		
Botryosphaeria blight[*] (<i>Botryosphaeria</i> sp.)	6.2 oz/A (0.7 oz ai/A)	
Notes: <ul style="list-style-type: none"> • Use in alternation with fungicides that have different modes of action. • Do not apply more than 3 applications of Ph-D Fungicide (2.1 oz ai/A) per season. [*Not for use in California.]		

Arysta LifeScience North America, LLC
 15401 Weston Parkway, Suite 150
 Cary, NC 27513

Ph-D® is a registered trademark of Arysta LifeScience North America, LLC.

Arysta LifeScience and the Arysta LifeScience logo are registered trademarks of Arysta LifeScience Corporation
 Supplemental Label Ph-D Fungicide