

66330-305

2/17/2009

10/19



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

FEB 17 2009

Mr. Bill Washburn
Regulatory Manager
Arysta LifeScience North America LLC
15401 Weston Parkway, Suite 150
Cary, NC 27513

Subject: Label Notifications for Pesticide Registration Notices 2007-4; Update Company Name per Pesticide Registration Notice 98-10

Dear Mr. Washburn,

The Agency is in receipt of your Applications for Pesticide Notification under Pesticide Registration Notices (PRN) 98-10 and 2007-4 dated February 3, 2009 for the following products:

- | | |
|--|--|
| Flomet 4L | EPA Registration Number 66330-260 |
| Ethephon 3.9% S&O | EPA Registration Number 66330-283 |
| Iprodione 2SE Turf | EPA Registration Number 66330-305 |
| Pix[®] Plant Regulator | EPA Registration Number 66330-345 |
| Pix[®] Ultra Plant Regulator | EPA Registration Number 66330-350 |
| Ethephon 3# Plant Regulator | EPA Registration Number 66330-363 |

The Registration Division (RD) has conducted its review of these requests for their applicability under PRNs 98-10 and 2007-4 and finds that the label changes requested fall within the scope of PRNs 98-10 and 2007-4. The labels submitted with the applications have been stamped "Notification" and will be placed in our records.

Please be reminded that 40 CFR Part 156.140(a)(4) requires that a batch code, lot number, or other code identifying the batch of the pesticide distributed and sold be placed on nonrefillable containers. The code may appear either on the label (and can be added by non-notification/PR Notice 98-10) or durably marked on the container itself.

MADE 1/5/10

2019

If you have any questions, please call me directly at 703-305-6249 or Steve Schaible of my staff at 703-308-9362.

Sincerely,



Linda Arrington
Notifications & Minor Formulations Team Leader
Registration Division (7505P)
Office of Pesticide Programs

30419

Please read instructions on reverse before completing form.

Form Approved, OMB No. 2070-0060, Approval expires 05-31-98

 EPA	United States Environmental Protection Agency Washington, DC 20460	<input type="checkbox"/> Registration <input type="checkbox"/> Amendment <input checked="" type="checkbox"/> Other:	OPP Identifier Number
--	---	---	-----------------------

Application for Pesticide - Section I

1. Company/Product Number 66330-305	2. EPA Product Manager MARY WALLER	3. Proposed Classification <input type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) IPRODIONE 2SE TURF	PM# 21	
5. Name and Address of Applicant (Include ZIP Code) Arysta LifeScience North America, LLC 15401 Weston Parkway, Suite 150 Cary, NC 27513 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(I), my product is similar or identical in composition and labeling to: <div style="text-align: center;">NOTIFICATION</div> EPA Reg. No. _____ <div style="text-align: center; font-size: 1.2em;">FEB 17 2009</div> Product Name _____	

Section - II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below

Explanation: Use additional page(s) if necessary. (For Section I and Section II.)

Notification of label change per PR Notice 2007-4. This notification is consistent with guidance in PR Notice 2007-4 and the requirements of EPA's regulations at 40 CFR §§ 156.10, 156.140, 156.144, 156.146, and 156.156. No other changes have been made to the labeling or the Confidential Statement of Formula for this product. I understand that it is a violation of 18 USC Sec 1001 to willfully make any false statement to EPA. I further understand that if the amendment is not consistent with the requirements of 40 CFR §§ 156.10, 156.140, 156.144, 156.146, and 156.156, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

Notification of label change per PR Notice 98-10. This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 USC Sec 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

Section - III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes* <input checked="" type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No			<input type="checkbox"/> Metal <input checked="" type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify)
*Certification must be submitted		If "Yes" Unit Packaging wgt.	No. per container	If "Yes" Package wgt.	No. per container
3. Location of Net Contents Information <input checked="" type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container 1.0 gal., 2.5 gal, 5.0 gal		5. Location of Label Directions <input checked="" type="checkbox"/> On Label <input type="checkbox"/> On labeling accompanying product	
6. Manner in Which Label is Affixed to Product			<input type="checkbox"/> Lithograph <input type="checkbox"/> Other <input checked="" type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled		

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application)		
Name Bill Washburn	Title Regulatory Manager	Telephone No. (Include Area Code) 901-432-5118
I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.		6. Date Application Received <div style="border: 1px solid black; padding: 5px; display: inline-block;">(Stamped)</div>
2. Signature 		
3. Title Regulatory Manager		
4. Typed Name Bill Washburn		5. Date 02/03/09



February 3, 2009

Ms. Mary Waller, PM 21
Document Processing Desk (NOTIF)
Office of Pesticide Programs – 7504P
U.S. Environmental Protection Agency
One Potomac Yard, Room S-4900
2777 South Crystal Drive
Arlington, VA 22202

**Subject: IPRADIONE 2SE TURF
EPA Reg. No. 66330-305**

***Notification of Label Change per PR Notice 2007-4
and PR Notice 98-10***

Dear Ms. Waller:

Please find the following enclosed:

- Application for Pesticide Registration (Other) dated 02/03/09.
- One highlighted copy of subject label, showing all changes.
- One clean copy of the subject label.


Notification of label change per PR Notice 2007-4. This notification is consistent with guidance in PR Notice 2007-4 and the requirements of EPA's regulations at 40 CFR §§ 156.10, 156.140, 156.144, 156.146, and 156.156. No other changes have been made to the labeling or the Confidential Statement of Formula for this product. I understand that it is a violation of 18 USC Sec 1001 to willfully make any false statement to EPA. I further understand that if the amendment is not consistent with the requirements of 40 CFR §§ 156.10, 156.140, 156.144, 156.146, and 156.156, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

Also in this submission, we are using PR Notice 98-10 to update the name of the company, emergency telephone numbers and warranty. Arysta LifeScience North America Corporation was changed to Arysta LifeScience North America, LLC.

Should you have any questions or comments, please do not hesitate to contact me at 901-432-5118 or by e-mail at bill.washburn@arystalifescience.com

Sincerely,

Bill Washburn
Regulatory Manager



5 of 19

IPRODIONE 2SE TURF

A Fungicide For The Prevention And Control Of Certain Diseases Of Turfgrass and Ornamentals

ACTIVE INGREDIENT:

Iprodione: 3-(3,5-dichlorophenyl)-N-(1-methylethyl)-2,4-dioxo-1-imidazolidinecarboxamide)*..... 23.3%

OTHER INGREDIENTS:..... 76.7%

TOTAL:..... 100.0%

This product contains petroleum distillate.
*Equivalent to 2 pounds Iprodione per gallon.

NOTIFICATION

FEB 17 2009

KEEP OUT OF REACH OF CHILDREN

CAUTION

FIRST AID

IF ON SKIN OR CLOTHING:

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes.
- Call a poison control center or doctor for treatment advice.

IF IN EYES:

- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for treatment advice.

IF SWALLOWED:

- Immediately call a poison control center or doctor.
- Do not induce vomiting unless told to do so by a poison control center or doctor.
- Do not give any liquid to the person. Do not give anything by mouth to an unconscious person.

IF INHALED:

- Move person to fresh air.
- If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.
- Call a poison control center or doctor for further treatment advice.

NOTE TO PHYSICIAN: Contains petroleum distillate.

EMERGENCY TELEPHONE NUMBERS: Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

FOR 24-HOUR EMERGENCY MEDICAL ASSISTANCE CALL: 1-866-303-6952

FOR CHEMICAL EMERGENCY: Spill, leak, fire, exposure, or accident, call CHEMTREC 1-800-424-9300

EPA Reg. No. 66330-305

EPA Est. No. 51036-GA-001

AD 110701/Not.071504

NET CONTENTS:

Manufactured For:
ARYSTA LIFESCIENCE NORTH AMERICA, LLC
15401 Weston Parkway, Suite 150
Cary, NC 27513

1

PRECAUTIONARY STATEMENTS

CAUTION

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Avoid contact with skin, eyes, or clothing. Harmful if swallowed or inhaled. In case of contact, immediately flush eyes or skin with plenty of water. Get medical attention if irritation persists. Avoid breathing spray mist.

PERSONAL PROTECTIVE EQUIPMENT

Mixers, loaders, others exposed to the concentrate, cleaners/repairers of equipment, and applicators applying as a dip treatment must wear:

1. Long-sleeved shirt and long pants
2. Chemical-resistant gloves such as barrier laminate, nitrile rubber (>14 mils), neoprene rubber (>14 mils), or viton (>14 mils)
3. Chemical-resistant apron, and
4. Chemical-resistant footwear plus socks.

Applicators using hand held equipment must wear

1. Coveralls over long-sleeve shirt and long pants,
2. Chemical-resistant gloves such as barrier laminate, nitrile rubber (>14 mils), neoprene rubber (>14 mils), or viton (>14 mils)
3. Chemical-resistant footwear plus socks,
4. Chemical-resistant headgear for overhead exposures, and
5. A dust/mist filtering respirator (MSHA/NIOSH approval number prefix TC21C), or a NIOSH approved respirator with any R, P or HE filter.

Applicators using aircraft or mechanical ground equipment (groundboom, airblast, etc.), and flaggers for aerial applications must wear

1. Long-sleeve shirt and long pants, and
2. Shoes plus socks

Applicators using truck-mounted equipment with a handgun at the end of the hose (i.e., for commercial turfgrass and ornamental

7019

applications) and all other handlers not specified above must wear

1. Long-sleeve shirt and long pants,
2. Chemical-resistant gloves such as barrier laminate, nitrile rubber (>14 mils), neoprene rubber (>14 mils), or viton (>14 mils), and
3. Shoes plus socks.

Follow manufacturer's instructions for cleaning/maintaining personal protective equipment (PPE). If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Discard clothing or other materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

ENGINEERING CONTROLS

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

1. Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
2. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
3. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This chemical can contaminate surface water through aerial and ground spray applications. Under some conditions, it may also have a high potential for runoff into surface water after application. These include poorly draining or wet soils with readily visible slopes toward adjacent surface waters, frequently flooded areas, areas overlaying extremely shallow ground water, areas with infield canals or ditches that drain to surface water, areas not separated from adjacent surface waters with vegetated filter strips, and areas overlaying tile drainage systems that drain to surface water.

This pesticide is toxic to invertebrates. Do not apply directly to water, to areas where surface water is present or to

intertidal areas below the mean high water mark. Drift or runoff may be hazardous to invertebrates in neighboring areas. Do not contaminate water when disposing of equipment washwater or rinsate.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Read entire label before using this product.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval of 12 hours for ornamental applications. The restricted entry interval for all other WPS uses is 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

1. Coveralls
2. Chemical-resistant gloves such as barrier laminate, nitrile rubber (>14 mils), neoprene rubber (>14 mils), or viton (>14 mils)
3. Shoes plus socks

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to turf uses (golf courses,

landscape and institutional areas) of this product that are NOT within the scope of the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses.

Do not enter or allow others to enter the treated area until sprays have dried.

STORAGE AND DISPOSAL

STORAGE: Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container $\frac{1}{4}$ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Offer for recycling, if available, or puncture and dispose of in a sanitary landfill or by other procedures approved by State and local authorities.

General Use: In order to assure maximum crop tolerance and disease control, follow recommendations on this label and all the precautions and limitations of the package label.

GENERAL PRECAUTIONS AND RESTRICTIONS

Use of this product at residential sites is prohibited.

Except for use on golf courses, if applying this product adjacent to a water body such as a lake, reservoir, river, permanent stream, marsh or natural pond, estuary, or commercial fish pond, there must be at least a 25-foot vegetative buffer strip between the water body and the point of application.

For golf courses only, do not apply to turf cut higher than 1" on golf courses where water bodies are present.

Do not apply this product when wind direction is toward aquatic areas.

TURF

(Golf Courses, Sod Farms And Institutional Areas)

Do not prepare more spray solution than can be used within 12 hours to minimize potential active ingredient degradation.

Initiate application when presence of disease is detected or if weather conditions are favorable for disease development.

DISEASE	APPLICATION SITE	APPLICATION RATE (fl. ozs./1000 FT ²)	SPRAY INTERVAL (days)
Dollar Spot (<i>Lanzia spp.</i> and <i>Moellerodiscus spp.</i>)	Golf course greens and tee boxes	3 - 4	14 - 21
Brown Patch (<i>Rhizoctonia solanii</i>)	Golf course fairways and other turf Areas	2 - 4	14 - 28
Leaf Spot such as Helminthosporium Leaf Spot caused by (<i>Drechslera spp.</i>)			
COMMENTS: Continue applications until disease pressure has reached acceptable level. Rates may be reduced to 2 fluid ounces to control Dollar Spot on fairways.			
Large Patch * (<i>Rhizoctonia solanii</i>)	All turf areas (except residential lawns)	4	14 - 21
COMMENTS: Initial application should be made in the Fall when conditions become favorable for disease development. Repeat applications in the Spring if disease pressure remains.			
Fusarium Blight (<i>Fusarium spp.</i>)	All turf areas (except residential lawns)	8	28
Necrotic Ring Spot* (<i>Leptosphaeria korrae</i>)			
COMMENTS: Apply prior to disease development as a preventative measure when weather conditions favor disease development.			
Fusarium Patch (<i>Microdochium nivale</i>)	All turf areas (except residential lawns)	4 - 8	14 - 21
(Pacific Northwest Only - West of the Cascade Mountains]			
COMMENTS: Continue applications until disease pressure has reached acceptable level.			
Gray Snow Mold (<i>Typhula spp.</i>)	All turf areas (except residential lawns)	4 - 8	Make initial application prior to first snow cover.
Pink Snow Mold (<i>Fusarium nivale</i>)			

DISEASE	APPLICATION SITE	APPLICATION RATE (fl. ozs./1000 FT ²)	SPRAY INTERVAL (days)
COMMENTS: A follow up application may be applied during a mid-winter thaw.			
Corticium Red Thread (<i>Laetisaria fuciformis</i>)	All turf areas (except residential lawns)	4	14
COMMENTS: Make preventative applications until conditions no longer favor disease development.			

Apply the recommended rates in the tables in 0.5 to 10 gallons of water per 1000 FT².

Applications of IPRDIONE 2SE must not exceed a maximum of 35 fluid oz. product/1000 ft.² a year.

Applications of IPRDIONE 2SE must be limited to no more than six a year.

Application of higher rates and shorter intervals are appropriate for all diseases when conditions are severe. Applications of lower rates and longer intervals are suggested for times when disease pressure is light to moderate.

Areas treated with IPRDIONE 2SE should not be mowed or irrigated until foliage has fully dried. A 24-hour period following application before mowing or irrigating is ideal.

IPRDIONE 2SE should not be combined with any type of sticker, extender or wetting agents. Do not allow livestock or poultry to graze on treated areas or feed on clippings from these areas.

*Not registered for use in California.

ORNAMENTALS

FIELD, LANDSCAPE AND GREENHOUSE ORNAMENTALS AND CONIFER NURSERIES*

* Conifer Nurseries not registered for use in California.

NOT FOR RESIDENTIAL USE. RECOMMENDED FOR USE BY COMMERCIAL NURSERY AND LANDSCAPE PERSONNEL.

Due to the wide variety of ornamental plants, it is not possible to determine the potential phytotoxicity for IPRDIONE 2SE on every species. Users should apply a label rate on a small number of plants prior to large scale use to evaluate tolerance.

ORNAMENTALS

ORNAMENTAL VARIETY		DISEASE
Ageratum	Holly	Aerial Web Blight (<i>Rhizoctonia sp.</i>)
Ajuga	Hoya	Alternaria Leaf Blight (<i>Alternaria euphorbiae</i>)
Almond (ornamental)	Hydrangea	Alternaria Leaf Spot (<i>Alternaria panax, Alternaria tenuissima</i>)
Alyssum	Impatiens*	Botrytis Blight (<i>Botrytis Sp.</i>)
Andromeda	Iris	Fusarium Leaf Spot (<i>Fusarium moniliforme</i>)
Aphelandra	Juniper	Helminthosporium Leaf Spot (<i>Helminthosporium sp.</i>)
Artemisia	Kalanchoe	Rhizoctonia stem and root rot (<i>Rhizoctonia sp.</i>)
Aster	Lillies	
Azalea	Lipstick vine	
Boxwood	(<i>Aeschynanthus</i>)	
Cactus	Marigold	
Calendula	Monarda (Bee Balm)	
Carnation	Pachysandra	
Cherry (ornamental)	Palm	
Chrysanthemum	Pansy	
Cineraria	Peach (ornamental)	
Cistena Plum	Peperomia	
Coleus	Periwinkle	
Columbine	Philodendron	
Coral Bells (Heuchera)	Phlox	
Crape Myrtle	Pilea	
Crassula	Pine	
Croton	Pitosporum	
Cyclamen	Plum (ornamental)	
Daffodils	Poinsettia	
Dahlia	Poppy	
Delphinium	Pothos*	
Deutzia	Primrose	
Dianthus	Privet	
Dieffenbachia	Protea	
Dizygotheca	Pyracantha	
Dogwood	Rhododendron	
Dracena	Rose	
English Ivy	Rose Tree of China	
Episcia	Salvia	
Euonymous	Schefflera	
Ficus	Snapdragon	
Forsythia	Statice	
Gazania	Tree Ivy	
Geranium	Tulip	
Gladiolus	Viburnum	
Gloxinia	Violet	
Gypsophila	Zinnia	
Hawthorn		
Iris		Ink Spot (<i>Drechslera iridis</i>)
Tulip		Tulip Fire (<i>Botrytis tulipae</i>)
Zinnia		Alternaria Leaf Blight (<i>Alternaria zinniae</i>)
Chrysanthemum		Ray Blight (<i>Ascochyta chrysanthami</i>)
Gladiolus		Fusarium Corm rot (<i>Fusarium oxysporum</i>)

Daffodils	Daffodil Leaf Scorch (<i>Stagnospora curtissi</i>)
Cistena Plum Plum (ornamental)	Blossom Blight (<i>Monilinia fructicola</i>)
Rose	Botrytis Storage Rot (<i>BotrytisSp.</i>)
Azalea Rhododendron	Cylindrocladium Blight and Wilt* (<i>Cylindrocladium scoparium</i>)

*NOTE: Do not use IPRODIONE 2SE Fungicide as a soil drench on Impatiens, and Pathos. Do not use IPRODIONE 2SE Fungicide on Spathiphyllum.

FOLIAR APPLICATION INSTRUCTIONS

Apply 1 to 2 ½ quarts of IPRODIONE 2SE in 100 gallons of water every 7 to 14 days until disease pressure is within acceptable levels. Spray plants to the point of run-off to insure thorough coverage.

Limit individual applications of IPRODIONE 2SE to a maximum of 2.5 quarts product/acre.

Limit total applications of Iprodione 2SE to a maximum of 4 per year.

Limit total applications to no more than 10 quarts of product/acre/year.

Application of IPRODIONE 2SE at higher rates and shorter intervals are appropriate for all diseases when conditions are severe. Applications of lower rates and longer intervals are suggested for times when disease pressure is light to moderate.

DRENCH APPLICATION INSTRUCTIONS

To control Rhizoctonia Stem and Root Rot (*Rhizoctonia spp.*), mix 13 fluid ounces in 100 gallons of water. Apply at seeding or transplant time 1 to 2 pints of this dilution per square foot of soil. Repeat application every 14 days as disease pressure warrants.

Limit applications of IPRODIONE 2SE to a maximum of 35 fluid ounces product/1000 ft² a year.

Limit individual applications of Iprodione 2SE to a maximum of 6 per year.

Application of IPRODIONE 2SE at higher rates and shorter intervals are appropriate for all diseases when conditions are severe. Applications of lower rates and longer intervals are suggested for times when disease pressure is light to moderate.

NOTE: Do not apply IPRDIONE 2SE *Spathiphyllum*. Do not apply as a drench on *impatiens* and *pathos*.

DIP APPLICATION INSTRUCTIONS

Roses - To control Botrytis Storage Rot (*Botrytis sp.*), mix 1 quart of IPRDIONE 2SE in 100 gallons of water and dip bare root for 5 minutes prior to cold storage.

AZALEA AND RHODODENDRON - To control Cylindrocladium Blight and Wilt* (*Cylindrocladium scoparium*), mix 1 quart of IPRDIONE 2SE in 100 gallons of water and dip cuttings for 5 minutes before planting.

GLADIOLUS - To control Fusarium Corm Rot (*Fusarium oxysporum*), mix 2 quarts of IPRDIONE 2SE in 100 gallons of water and dip corms for 5 minutes prior storage.

* Not registered for use in California.

TANK MIXTURES FOR TURF

To control additional diseases in turf such as Summer Stress Complex/Decline and Pythium Blight, IPRDIONE 2SE may be tank mixed with other fungicides.

Gray Snow Mold control will be enhanced by a tank mixture with a properly labeled chlorothalonil product such as Daconil 2787 Flowable or Daconil WG. Initial application of tank mix should be made prior to first snow cover and followed by another application if a mid-winter thaw occurs.

User should adhere to any applicable restrictions on the tank mix product. If compatibility of tank mix partners is unknown, mixing a small amount of the products in proper ratios in a clear jar is advised prior to mixing a large tank.

Do not exceed a total of 35 fluid oz. IPRDIONE 2SE TURF per 1000 ft² per year with a maximum of 6 applications.

DIRECTIONS THROUGH SPRINKLER IRRIGATION SYSTEMS

Do not use through sprinkler irrigation systems in California.

Apply this product only through sprinkler irrigation systems including center pivot. Do not apply this product through any other type of irrigation system.

SPRAY PREPARATION: Remove scale, pesticide residues, and other foreign matter from the chemical tank and entire injector system. Flush with clean water.

APPLICATION INSTRUCTIONS: First prepare a suspension of IPRODIONE 2SE TURF in a mix tank. Fill tank with 1/2 to 3/4 the desired amount of water. Start mechanical or hydraulic agitation. Add the required amount of IPRODIONE 2SE TURF, and then the remaining volume of water. (Suspension concentrations using the appropriate dosage per acre recommended on this label of IPRODIONE 2SE TURF per 1 to 4 gallons of water are recommended). Then set sprinkler to deliver 0.1 to 0.3 inch of water per acre. Start sprinkler and uniformly inject the suspension of IPRODIONE 2SE TURF into the irrigation water line so as to deliver the desired rate per acre. The suspension of IPRODIONE 2SE TURF should be injected with a positive displacement pump into the main line ahead of a right angle turn to insure adequate mixing. If you should have any other questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.

NOTE: When treatment with IPRODIONE 2SE TURF has been completed, further field irrigation over the treated area should be avoided for 24 to 48 hours to prevent washing the chemical off the crop.

GENERAL PRECAUTIONS FOR APPLICATIONS THROUGH SPRINKLER IRRIGATION SYSTEMS

Maintain continuous agitation in mix tank during mixing and application to assure a uniform suspension.

Greater accuracy in calibration and distribution will be achieved by injecting a larger volume of a more dilute solution per unit time.

The system must contain a functional check valve, vacuum relief valve, and low-pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump. The pesticide injection pipeline must always contain a functional, normally closed solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shutdown. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected. Systems must use a metering pump, such as a positive displacement injection pump (e. g., diaphragm pump) effectively designed and constructed of

materials that are compatible with pesticides and capable of being flitted with a system interlock.

Do not apply when wind speed favors drift, when system connection or fittings leak, when nozzles do not provide uniform distribution or when lines containing the product must be dismantled and drained.

Crop injury, lack of effectiveness, or illegal pesticide residues in the crop may result from nonuniform distribution of treated water.

Allow sufficient time for pesticide to be flushed through all lines and all nozzles before turning off irrigation water. A person knowledgeable of the chemigation system and responsible for its operation must shut the system down and make necessary adjustments should the need arise.

Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the label described safety devices for public water supplies are in place.

SPRAY DRIFT MANAGEMENT

Sensitive Areas:

This pesticide should only be applied when the potential for drift to adjacent sensitive areas (e.g., residential areas, bodies of water, known habitats for threatened or endangered species, non-target crops) is minimal (e.g., when wind is blowing away from the sensitive areas).

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment-and-weather-related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions.

The following drift management requirements must be followed to avoid off-target drift movement from aerial applications to agricultural field crops. These requirements do not apply to forestry applications, public health uses or to applications using dry formulations.

1. The distance of the outer most nozzles on the boom must not exceed $3/4$ the length of the wingspan or rotor.
2. Nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees.

Where states have more stringent regulations, they should be observed. The applicator should be familiar with and take into account the information covered in the Aerial Drift Reduction Advisory Information.

The following section is advisory in nature and does not supersede the mandatory label requirements.

INFORMATION ON DROPLET SIZE (This section is advisory in nature and does not supersede the mandatory label requirements)

The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions (See Wind, Temperature and Humidity, and Temperature Inversions).

CONTROLLING DROPLET SIZE (This section is advisory in nature and does not supersede the mandatory label requirements)

Volume - Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.

Pressure - Do not exceed the nozzle manufacturer's recommended pressures. For many nozzle types lower pressure produces larger droplets. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.

Number of Nozzles - Use the minimum number of nozzles that provide uniform coverage.

Nozzle Orientation - Orienting nozzles so that the spray is released parallel to the airstream produces larger droplets than other orientations and is the recommended practice. Significant deflection from horizontal will reduce droplet size and increase drift potential.

Nozzle Type - Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce the largest droplets and the lowest drift.

BOOM LENGTH (This section is advisory in nature and does not supersede the mandatory label requirements)

For some use patterns, reducing the effective boom length to less than $3/4$ of the wingspan or rotor length may further reduce drift without reducing swath width.

APPLICATION HEIGHT (This section is advisory in nature and does not supersede the mandatory label requirements)

Applications should not be made at a height greater than 10 feet above the top of the target plants unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind.

SWATH ADJUSTMENT (This section is advisory in nature and does not supersede the mandatory label requirements)

When applications are made with a crosswind, the swath will be displaced downwind. Therefore, on the up and downwind edges of the field, the applicator should compensate for the displacement by adjusting the path of the aircraft upwind. Swath adjustment distance should increase, with the increasing drift potential (higher wind, smaller drops, etc.).

WIND (This section is advisory in nature and does not supersede the mandatory label requirements)

Drift potential is lowest between wind speeds of 2 - 10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given speed. Application should be avoided below 2 mph due to variable wind direction and high inversion potential. NOTE: Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

TEMPERATURE AND HUMIDITY (This section is advisory in nature and does not supersede the mandatory label requirements)

When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

TEMPERATURE INVERSIONS (This section is advisory in nature and does not supersede the mandatory label requirements)

Applications should not occur during a temperature inversion because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to light variable winds common during

inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

Warranty and Disclaimer Statement

The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Such risks may arise from weather conditions, soil factors, off-target movement, unconventional farming techniques, the presence of other materials, the manner of use or application, or other unknown factors, all of which are beyond the control of Arysta LifeScience North America, LLC ("Arysta"), and can cause crop injury, injury to non-target crops or plants, ineffectiveness of the product, or other unintended consequences. All such risks shall be assumed by the user or buyer.

Arysta warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks described above, when used in accordance with the Directions for Use under normal conditions. This warranty does not extend to the use of this product contrary to label instructions or under conditions not reasonably foreseeable to Arysta, and is subject to the inherent risks described above.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, ARYSTA DISCLAIMS ALL OTHER WARRANTIES, EXPRESSED OR IMPLIED, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, ARYSTA, MANUFACTURER, AND SELLER DISCLAIM AND SHALL NOT BE LIABLE FOR ANY SPECIAL, INCIDENTAL, INDIRECT, OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE, HANDLING, APPLICATION, STORAGE, OR DISPOSAL OF THIS PRODUCT OR FOR DAMAGES IN THE NATURE OF PENALTIES, AND THE USER AND BUYER WAIVE ANY RIGHT THAT THEY MAY HAVE TO SUCH DAMAGES. NO AGENT, REPRESENTATIVE OR EMPLOYEE OF ARYSTA IS AUTHORIZED TO MAKE ANY WARRANTY, GUARANTEE OR REPRESENTATION BEYOND THOSE CONTAINED HEREIN OR TO MODIFY THE WARRANTIES CONTAINED HEREIN. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE TOTAL LIABILITY OF ARYSTA, MANUFACTURER, AND SELLER, SHALL BE LIMITED TO THE PURCHASE PRICE PAID, OR AT ARYSTA'S ELECTION, THE REPLACEMENT OF THE PRODUCT.

NOTIFICATION

FEB 17 2009