

66330-362

02-08-2011

1/43



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

FEB 8 2011

Gary R. Orr, Ph.D.
Product Registration
Arysta Lifescience NA Corporation
15401 Weston Parkway, Suite 150
Cary, NC 27513

SUBJECT: Application for Pesticide Notification (PRN 98-10)
Request Alternate Brand Name "Rolling 72"
EPA Reg. No. 66330-362
Application Dated November 5, 2010

Dear: Ms. Thomas,

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 dated 11/05/10 for the above product. The Registration Division (RD) has conducted a preliminary screen of this request for its applicability under PRN 98-10 and finds that the action(s) requested fall within the scope of PRN 98-10. The label submitted with the application has been stamped "Notification" and will be placed in our records.

If you have any questions, please call me directly at 703-305-5335 or Owen F. Beeder of my staff at 703-308-8899.

Sincerely,

A handwritten signature in black ink that reads "Paul J. Mastradone".

Paul J. Mastradone, Ph.D., Acting
Notifications & Minor Formulations Team Leader
Registration Division (7505P)
Office of Pesticide Programs

2
43

 EPA United States Environmental Protection Agency Washington, DC 20460	<input type="checkbox"/> Registration <input type="checkbox"/> Amendment <input checked="" type="checkbox"/> Other:	OPP Identifier Number
---	--	-----------------------

Application for Pesticide - Section I

1. Company/Product Number 66330-362	2. EPA Product Manager Tony Kish	3. Proposed Classification <input type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Chlorothalonil 720 SC	PM# 22	
5. Name and Address of Applicant (Include ZIP Code) Arysta LifeScience North America Corporation 15401 Weston Parkway, Suite 150 Cary, NC 27513 <input type="checkbox"/> Check if this is a new address		6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(1), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____

Section - II

NOTIFICATION
FEB - 8 2011

<input type="checkbox"/> Amendment - Explain below. <input type="checkbox"/> Resubmission in response to Agency letter dated _____ <input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____ <input type="checkbox"/> "Me Too" Application <input type="checkbox"/> Other - Explain below
--	---

Explanation: Use additional page(s) if necessary. (For Section I and Section II.)


NOTIF Addition of an **alternate brand name** Rolling 72 SC for the product bearing the **primary brand name** Chlorothalonil 720 SC (EPA Reg. No. 66330-362).

Contact: gary.orr@arystalifescience.com

Section - III

1. Material This Product Will Be Packaged In:			
Child-Resistant Packaging <input type="checkbox"/> Yes* <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No If "Yes" Unit Packaging wgt. No. per container	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No If "Yes" Package wgt. No. per container	2. Type of Container <input type="checkbox"/> Metal <input type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify)
* Certification must be submitted			
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container	4. Size(s) Retail Container 40 LB BAG	5. Location of Label Directions <input type="checkbox"/> On Label <input type="checkbox"/> On labeling accompanying product	
6. Manner in Which Label is Affixed to Product <input type="checkbox"/> Lithograph <input type="checkbox"/> Other _____ <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled			

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application)		
Name Gary R. Orr, PhD	Title Regulatory Manager	Telephone No. (Include Area Code): 919-678-4879
I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.		6. Date Application Received (Stamped)
2. Signature 	3. Title Regulatory Manager	
4. Typed Name Gary R. Orr, PhD	5. Date November 5, 2010	

4
43

NOTIFICATION

FEB - 8 2011

ROLLING 72 SC
Flowable Fungicide

Agricultural Fungicide

ACTIVE INGREDIENT:

Chlorothalonil (tetrachloroisophthalonitrile) 54.0%

INERT INGREDIENTS:..... 46.0%

TOTAL:..... 100.0%

This product contains 6.0 pounds of Chlorothalonil per gallon (720 g per liter)

KEEP OUT OF REACH OF CHILDREN

CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label find someone to explain it to you in detail.)

See inside booklet for additional precautionary statements

For Product Use Information Call 1-866-761-9397

Arysta LifeScience North America, LLC
15401 Weston Parkway, Suite 150
Cary, NC 27513

EPA Reg. No. 66330-362
EPA Est. No.

NET WEIGHT: _____

FIRST AID	
IF SWALLOWED	<ul style="list-style-type: none"> • Call a poison control center or doctor for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
IF ON SKIN OR CLOTHING	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15–20 minutes. • Call a poison control center or doctor for treatment advice.
IF INHALED	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth to mouth, if possible. • Call a poison control center or doctor for further treatment advice.
IF IN EYES	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
NOTE TO PHYSICIAN	
Persons suffering with temporary allergic skin reactions may respond to treatment with oral antihistamines and topical or oral steroids.	
Have the product container or label with you when calling a poison control center or doctor or going for treatment.	
HOTLINE NUMBERS:	
FOR 24-HOUR MEDICAL EMERGENCY ASSISTANCE:	
Call PROSAR at 1-866-303-6952 or 1-651-632-8946 if calling from outside of the U.S.	
FOR 24-HOUR CHEMICAL EMERGENCY ASSISTANCE:	
Spill, leak, fire, exposure, or accident	
Call CHEMTREC at 1-800-424-9300 or 1-703-527-3887 if calling from outside of the U.S.	

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION:

Harmful if swallowed, absorbed through skin, or inhaled. Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Avoid breathing spray mist. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical resistant to this product are made of any waterproof material. If you want more options, follow the instructions for category A on an EPA chemical resistance category selection chart.

Mixers, loaders, applicators and all others who handle this pesticide must wear:

- Long-sleeved shirt and long pants
- Chemical resistant gloves made of any waterproof material – Category A (e.g. barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, natural rubber, polyethylene, polyvinyl chloride (PVS) or viton)
- Shoes plus socks

USER SAFETY REQUIREMENTS

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing or other materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

ENGINEERING CONTROLS STATEMENT

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS**Users should:**

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to aquatic invertebrates and wildlife. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment wash water or rinsate.

This chemical is known to leach through soil into groundwater under certain conditions as a result of label use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

This chemical can contaminate surface water through spray drift. Under some conditions, it may also have a high potential for runoff into surface water for several days to weeks after application. These include poorly draining or wet soils with readily visible slopes toward adjacent surface waters, frequently flooded areas, areas overlaying extremely shallow ground water, areas with infield canals or ditches that drain to surface water, areas not separated from adjacent surface waters with vegetated filter strips, and areas over-laying tile drainage systems that drain to surface water.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, or pets either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of **12 hours**.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is: long-sleeved shirt and long pants or coveralls, shoes plus socks, chemical resistant gloves made of any waterproof material, such as nitrile, butyl, neoprene, and / or barrier laminate, and protective eyewear.

Special Eye Irritation Provisions: This product is a severe eye irritant. Although the restricted entry interval expires after 12 hours, for the next 6.5 days entry is permitted only when the following safety measures are provided:

1. At least one container designed specifically for flushing eyes must be available in operating condition at the WPS required decontamination site intended for workers entering the treated area.
2. Workers must be informed, in a manner they can understand:
 - that residues in the treated area may be highly irritating to their eyes
 - that they should take precautions, such as refraining from rubbing their eyes, to keep the residues out of their eyes
 - that if they do get residues in their eyes, they should immediately flush their eyes using the eyeflush container that is located at the decontamination site or using other readily available clean water, and
 - how to operate the eyeflush container

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE

Store in original container and keep tightly closed. Store in a cool, dry place. Protect from excessive heat.

PESTICIDE DISPOSAL

Pesticide wastes are toxic. Improper disposal of excess pesticide, pesticide spray or rinsate is a violation of federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL

Plastic Containers equal to or less than 5 gallons: Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip.

Plastic Containers greater than 5 gallons: Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

Offer for recycling, if available, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Returnable Refillable Containers/ Bulk and Minibulk Containers:

Return container intact to point of purchase. Refillable container. Refill this container with pesticide only. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

GENERAL INFORMATION

Integrated Pest Management

ROLLING 72 SC is an effective fungicide when used according to label directions for control of a broad spectrum of plant diseases. **ROLLING 72 SC** is recommended for use in programs that are compatible with the principles of Integrated Pest Management (IPM), which include the use of disease resistant crop varieties, cultural practices, pest scouting and disease forecasting systems, which reduce unnecessary applications of pesticides.

Fungicide Resistance Management

ROLLING 72 SC is an effective product for strategic use in programs that attempt to minimize disease resistance to fungicides. Some other fungicides that are at risk from disease resistance exhibit a single-site mode of fungicidal action. **ROLLING 72 SC**, with a multi-site mode of action, may be used to delay or prevent the development of resistance to single-site fungicides. Consult your federal or state Cooperative Extension Service representatives for guidance on the proper use of **ROLLING 72 SC** in programs which seek to minimize the occurrence of disease resistance to other fungicides.

ROLLING 72 SC can be used effectively in dilute or concentrate sprays. Thorough, uniform coverage is essential for disease control.

GENERAL PRECAUTIONS AND RESTRICTIONS

DO NOT use on greenhouse-grown crops.

Tank Mixing

DO NOT combine **ROLLING 72 SC** in spray tank with pesticides, surfactants or fertilizers, unless prior use has shown the combination physically compatible, effective and noninjurious under your conditions of use. DO NOT combine **ROLLING 72 SC** with DiPel® or Latron B-1956® or Latron AG-98® as phytotoxicity may result from the combination when applied to crops listed on this label.

This product must not be applied within 150 feet (for aerial applications) or 25 feet (for ground applications) of marine/estuarine water bodies unless there is an untreated buffer area of that width between the area to be treated and the water body.

SPRAY DRIFT PRECAUTIONS

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment and weather related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions.

The following drift management requirements must be followed to avoid off-target drift movement from aerial applications to agricultural field crops. These requirements do not apply to forestry applications, public health uses or to applications using dry formulations.

1. The distance of the outer most nozzles on the boom must not exceed ¾ the length of the wingspan or rotor.
2. Nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees.

Where states have more stringent regulations, they should be observed.

The applicator should be familiar with and take into account the information covered in the *Aerial Drift Reduction Advisory* section.

Aerial Drift Reduction Advisory

This section is advisory in nature and does not supersede the mandatory label requirements.

Information on Droplet Size

The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions (see Wind, Temperature and Humidity, and Temperature Inversions below).

Controlling Droplet Size

- **Volume** - Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.
- **Pressure** - Do not exceed the nozzle manufacturer's recommended pressures. For many nozzle types lower pressure produces larger droplets. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.
- **Number of nozzles** - Use the minimum number of nozzles that provide uniform coverage.
- **Nozzle Orientation** - Orienting the nozzles so that the spray is released parallel to the airstream produces larger droplets than other orientations and is the recommended practice. Significant deflection from horizontal will reduce droplet size and increase drift potential.
- **Nozzle Type** - Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce the largest droplets and the lowest drift potential.

Boom Length

For some use patterns, reducing the effective boom length to less than 3/4 of the wingspan or rotor length may further reduce drift without reducing swath width.

Application Height

Applications should not be made at a height greater than 10 feet above the top of the largest plants unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind.

Swath Adjustment

When applications are made with a crosswind, the swath will be displaced downwind. Therefore, on the upwind and downwind edges of the field, the applicator should compensate for this displacement by adjusting the path of the aircraft upwind. Swath adjustment distance should increase with increasing drift potential (higher wind, smaller drops, etc.).

Wind

Drift potential is lowest between wind speeds of 2 - 10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given speed. Application should be avoided below 2 mph due to variable wind direction and high inversion potential. **NOTE:** Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

Temperature and Humidity

When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

Temperature Inversions

Applications should not occur during a temperature inversion because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

Sensitive Areas

The pesticide should only be applied when the potential for drift to adjacent sensitive areas (e.g., residential areas, bodies of water, known habitat for threatened or endangered species, nontarget crops) is minimal (e.g., when wind is blowing away from the sensitive areas).

APPLICATION INSTRUCTIONS

Dosage rates on this label indicate pints of **ROLLING 72 SC** per acre, unless otherwise stated. Under conditions favoring disease development, the higher rate specified and shortest application interval should be used.

Note: Slowly invert container several times to assure uniform mixture.

The required amount of **ROLLING 72 SC** should be added slowly into the spray tank during filling. With concentrate sprays, pre-mix the required amount of **ROLLING 72 SC** in a clean container and add to the spray tank as it is being filled. Keep agitator running when filling spray tank and during spray operations.

Apply **ROLLING 72 SC** in sufficient water to obtain adequate coverage of foliage. Gallonage to be used will vary with crop and amount of plant growth.

For field and row crops, spray volume usually will range from 20 to 150 gallons per acre for dilute sprays and 5 to 10 gallons per acre for concentrate ground sprays and aircraft applications.

For tree and orchard crops, apply **ROLLING 72 SC** in sufficient water and with proper calibration to obtain uniform coverage of tree canopy. For fruit and nut bearing crops, the maximum volume is 300 gallons per acre unless indicated otherwise in the specific use directions. For conifers, the maximum volume is 100 gallons per acre.

APPLICATION AND CALIBRATION TECHNIQUES FOR CHEMIGATION

Apply this product only through sprinkler irrigation systems including center pivot, motorized lateral move, traveling gun, solid set or portable (wheel move, side roll, end tow, or hand move) irrigation system(s). Do not apply this product through any other type of irrigation system.

Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.

If you have questions about calibration, you should contact State Extension Service Specialists, equipment manufacturers or other irrigation experts.

Do not apply this product through irrigation systems connected to a public water system. "Public water system" means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days per year.

Controls for both irrigation water and pesticide injection systems must be functionally interlocked, so as to automatically terminate pesticide injection when the irrigation water pump motor stops. A person knowledgeable of the irrigation system and responsible for its operation shall be present so as to discontinue pesticide injection and make necessary adjustments, should the need arise.

The irrigation water pipeline must be fitted with a functional, automatic, quick-closing check valve to prevent the flow of treated irrigation water back toward the water source. The pipeline must also be fitted with a vacuum relief valve and low pressure drain, located between the irrigation water pump and the check valve, to prevent back-siphoning of treated irrigation water into the water source.

Always inject **ROLLING 72 SC into irrigation water after it discharges from the irrigation pump and after it passes through the check valve. Never inject pesticides into the intake line on the suction side of the pump.**

Pesticide injection equipment must be fitted with a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump. Interlock this valve to the power system, so as to prevent fluid from

being withdrawn from the chemical supply tank when the irrigation system is either automatically or manually turned off.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Spray mixture in the chemical supply tank must be agitated at all times, otherwise settling and uneven application may occur. Do not apply when wind speed favors drift beyond the area intended for treatment.

Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump), effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

ROLLING 72 SC may be used through two basic types of sprinkler irrigation systems as outlined in Sections A and B below. Determine which type of system is in place, then refer to the appropriate directions provided for each type.

A. Center Pivot, Motorized Lateral Move and Traveling Gun Irrigation Equipment

For injection of pesticides, these continuously moving systems must use a positive displacement injection pump, of either diaphragm or piston type, constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock and capable of injection at pressures approximately 2 – 3 times those encountered within the irrigation water line. Venturi applicator units cannot be used on these systems.

Thoroughly mix recommended amount of **ROLLING 72 SC** for acreage to be covered into same amount of water used during calibration and inject into system continuously for one revolution or run. Mixture in the chemical supply tank must be continuously agitated during the injection run. Shut off injection equipment after one revolution or run, but continue to operate irrigation system until **ROLLING 72 SC** has been cleared from last sprinkler head.

B. Solid Set and Portable (Wheel Move, Side Roll, End Tow, or Hand Move) Irrigation Equipment

With stationary systems, an effectively designed in-line Venturi applicator unit is preferred which is constructed of materials that are compatible with pesticides; however, a positive-displacement pump can also be used.

Determine acreage covered by sprinkler. Fill tank of injection equipment with water and adjust flow to use contents over a 30 to 45 minute period. Mix desired amount of **ROLLING 72 SC** for acreage to be covered with water so that the total mixture of **ROLLING 72 SC** plus water in the injection tank is equal to the quantity of water used during calibration, and operate entire system at normal pressures recommended by the manufacturer of injection equipment used, for amount of time established during calibration. Agitation is recommended. **ROLLING 72 SC** can be injected at the beginning or end of the irrigation cycle or as a separate application. Stop injection equipment after treatment is completed and continue to operate irrigation system until **ROLLING 72 SC** has been cleared from last sprinkler head.

DIRECTIONS FOR APPLICATION - AGRICULTURAL CROPS

Crop	Diseases (Pathogen)	Pints/Acre (lbs a.i./Acre)	Application Directions

Crop	Diseases (Pathogen)	Pints/Acre (lbs a.i./Acre)	Application Directions
Asparagus	Rust (<i>Puccinia asparagi</i>) Purple Spot (<i>Pleospora herbarum</i>) Cercospora blight (<i>C. asparagi</i>)	2 – 4 (1.5 – 3.0)	Apply in water volumes of 25-50 gallons per acre. Begin applications after final harvest of spears. Repeat applications at 14-28 day intervals (the minimum re-treatment interval is 14 days), depending on disease pressure. Use the higher rate and shorter interval if disease severity begins to increase or weather conditions favor disease development. Ground applications only.
Restrictions and Limitations: <ul style="list-style-type: none"> DO NOT apply more than 12 pints ROLLING 72:SC (9.0 lbs a.i.) per acre during each growing season. DO NOT apply within 190 days (120 days in CA and AZ) of the harvest of spears in the following season. 			
Bean (Snap)	Rust (<i>Uromyces appendiculatus</i>)	1 3/8 – 3 (1.0 – 2.25)	Apply with sufficient water to obtain adequate coverage. Begin applications during early bloom stage or when disease first threatens and repeat as necessary (the minimum re-treatment interval is 7 days) to maintain control. Apply by ground, air or chemigation.
	Botrytis blight (gray mold) (<i>Botrytis cinerea</i>)	3 (2.25)	
Restrictions and Limitations: <ul style="list-style-type: none"> DO NOT apply more than 12 pints ROLLING 72:SC (9.0 lbs a.i.) per acre during each growing season. DO NOT apply within 7 days of harvest. 			
Beans (Dry) (except soybeans) bean, adzuki bean, broad bean, dry bean, lablab bean, navy bean, kidney bean, lima bean, moth bean, mung bean, pink bean, pinto bean, tepary bean, urd bean, yardlong catjang chickpea (garbanzo) cowpea lupin, grain lupin bean, rice	Rust (<i>Uromyces appendiculatus</i>) Anthracnose (<i>Colletotrichum lindemuthianum</i>) Downy mildew (<i>Phytophthora nicotianae</i>) Cercospora leaf blotch (<i>C. cruenta</i>) Ascochyta blight (<i>A. phaseolorum</i>)	1 3/8 – 2 (1.0 – 1.5)	Apply with sufficient water to obtain adequate coverage. Begin applications at first onset of disease, which may occur as early as 2 to 4 weeks before flowering, and repeat at 7 to 10 day intervals (the minimum re-treatment interval is 7 days). For use only on beans to be harvested dry with pods removed. Apply by ground, air or chemigation.

Crop	Diseases (Pathogen)	Pints/Acre (lbs a.i./Acre)	Application Directions
bean, runner bean, jackbean pea, blackeyed pea, southern			
Restrictions and Limitations: <ul style="list-style-type: none"> DO NOT apply more than 8 pints ROLLING 72 SC (6.0 lbs a.i.) per acre during each growing season. DO NOT apply within 14 days of harvest. 			
Blueberries	Suppression: Anthracnose (ripe rot) <i>(C. gloeosporoides)</i> Mummy Berry <i>(M. vaccinicorymbosi)</i>	3 – 4 (2.25 – 3.0)	ROLLING 72 SC should be used as part of an overall disease management strategy which includes alternation with a fungicide with a different mode of action. Diseases may only be suppressed and russetting may occur under heavy disease pressure or unfavorable environmental conditions. Apply with sufficient water to obtain adequate coverage, normally 20-100 gallons per acre. Begin applications at budbreak (green tip) and repeat at 10-day intervals through early bloom (the minimum re-treatment interval is 10 days). Under heavy disease pressure, use the higher rate. Ground or air applications only.
	Septoria leaf spot <i>(Septoria albopunctata)</i> Rust <i>(Pucciniastrum vaccinii)</i>	3 – 4 (2.25 – 3.0)	Post-Harvest Foliar Use (after all berries have been harvested): To maintain healthy leaves for the following season, apply with sufficient water to obtain adequate coverage (normally 20-100 gallons per acre). Repeat at 10-14 day intervals (the minimum re-treatment interval is 10 days). Ground or air applications only.
Restrictions and Limitations: <ul style="list-style-type: none"> DO NOT apply more than 12 pints ROLLING 72 SC (9.0 lbs a.i.) per acre during each growing season. DO NOT apply after full bloom (except for foliar use after harvest) or within 42 days of harvest. 			
Cabbage Chinese Cabbage (tight-headed varieties only) Cauliflower Broccoli Chinese Broccoli Brussels Sprouts	Alternaria leaf spot <i>(Alternaria spp.)</i> Downy mildew <i>(Peronospora parasitica)</i>	1 ½ (1.125)	Apply with sufficient water to obtain adequate coverage. Begin applications after transplants are set in field, or shortly after emergence of field-seeded crop, or when conditions favor disease development. Repeat at 7 to 10 day intervals (the minimum re-treatment interval is 7 days) or as necessary to maintain control.

Crop	Diseases (Pathogen)	Pints/Acre (lbs a.i./Acre)	Application Directions
			Apply by ground, air or chemigation.
	Ring spot (California only)	2 (1.5)	For field-seeded Brussels sprouts, begin applications at time of early sprout development or when conditions favor disease development. Repeat at 7 to 10 day intervals (the minimum re-treatment interval is 7 days) to maintain control.
Restrictions and Limitations: <ul style="list-style-type: none"> DO NOT apply more than 16 pints ROLLING 72 SC (12.0 lbs a.i.) per acre during each growing season. DO NOT apply within 7 days of harvest. 			
Carrot	Cercospora leaf spot (<i>C. carotae</i>) Alternaria leaf blight (<i>A. dauci</i>)	1 ½ - 2 (1.125 - 1.5)	Apply with sufficient water to obtain adequate coverage. Start applications when disease threatens and repeat at 7 to 10 day intervals (the minimum re-treatment interval is 7 days) to maintain control. Apply by ground, air or chemigation.
Restrictions and Limitations: <ul style="list-style-type: none"> DO NOT apply more than 20 pints ROLLING 72 SC (15.0 lbs a.i.) per acre during each growing season. ROLLING 72 SC may be applied the day of harvest. 			
Celery	Early blight (<i>Cercospora apii</i>) Late blight (<i>Septoria apicola</i>) Basal stalk rot (<i>Rhizoctonia solani</i>) Suppression: (7 day schedule) Pink rot (<i>Sclerotinia sclerotiorum</i>) Early blight (<i>Cercospora apii</i>) Late blight (<i>Septoria apicola</i>)	2 - 3 (1.5 - 2.25) 3 (2.25) 1 ½ - 2 (1.125 to 1.5) per 100 gal.	Apply with sufficient water to obtain adequate coverage. Start applications when transplants are set in the field and repeat at 7 day interval as needed to maintain control (the minimum re-treatment interval is 7 days). Apply by ground, air or chemigation. For celery seedbeds, apply in a spray volume of 125 gallons per acre twice weekly or as needed to maintain control. Start applications shortly after crop emergence. Use the higher rate under severe disease conditions.
Restrictions and Limitations: <ul style="list-style-type: none"> DO NOT apply more than 24 pints ROLLING 72 SC (18.0 lbs a.i.) per acre during each growing season. DO NOT apply within 7 days of harvest. 			

Crop	Diseases (Pathogen)	Pints/Acre (lbs a.i./Acre)	Application Directions
Corn (Sweet)	Helminthosporium leaf blights	¾ - 2 (0.6 - 1.5)	Apply with sufficient water to obtain adequate coverage. Begin applications when conditions favor disease development and repeat at 7 day intervals (the minimum re-treatment interval is 7 days). Under severe disease conditions, use 1 ½ to 2 pints ROLLING 72 SC per acre. Apply by ground, air or chemigation.
Corn (grown for seed)	Rust (<i>Puccinia</i> spp.)		

Restrictions and Limitations:

- DO NOT apply more than 12 pints ROLLING 72 SC (9.0 lbs a.i.) per acre during each growing season.
- DO NOT apply within 14 days of harvest.
- DO NOT apply to sweet corn to be processed.
- DO NOT allow livestock to graze in treated fields.
- DO NOT ensile treated corn or use as livestock forage.

Cranberry	Fruit Rots	4 - 6 ½ (3.0 - 4.9)	Apply at early bloom and repeat at 10 to 14 day intervals (the minimum re-treatment interval is 10 days). Under severe disease conditions, use the 6 ½ pint per acre rate on a 10 day schedule. Apply by ground, air or chemigation. When applying by chemigation, apply in 300 gallons of water per acre through solid set systems only.
	Lophodermium leaf/twig blight (<i>L. hypophyllum</i>)		
	Upright Dieback (<i>Phomopsis vaccinii</i>)	4 - 6 ½ (3.0 - 4.9)	Apply with sufficient water to obtain coverage of uprights and runners. Make the first application prior to bloom, when shoots begin growth in the spring. Make additional applications at 10-14 day intervals. Apply by ground, air or chemigation. When applying by chemigation, use 300 gallons of water per acre through solid set systems only.

Restrictions and Limitations:

- DO NOT apply more than 20 pints ROLLING 72 SC (15.0 lbs a.i.) per acre during each growing season.
- DO NOT apply within 50 days of harvest.
- DO NOT apply to beds when flooded or allow release of irrigation water from beds for at least 3 days following application.

Cucurbits Cucumber Cantaloupe Muskmelon Honeydew melon Watermelon Squash	Anthracnose (<i>Colletotrichum</i> spp.) Downy mildew (<i>Pseudoperonospora cubensis</i>)	1 ½ - 2 (1.125 - 1.5)	Apply with sufficient water to obtain adequate coverage. Begin applications when plants are in first true leaf stage or when conditions are favorable for disease development. Repeat applications at 7 day intervals (the minimum re-treatment interval is 7
--	---	-----------------------	---

Crop	Diseases (Pathogen)	Pints/Acre (lbs a.i./Acre)	Application Directions
Pumpkin	Target spot (<i>Corynespora cassiicola</i>)		<p>days).</p> <p>Note: Spraying mature watermelons may result in sunburn of the upper surface of the fruit. DO NOT apply ROLLING 72 SC to watermelons when any of the following conditions are present:</p> <ul style="list-style-type: none"> • intense heat and sunlight • drought conditions • poor vine canopy • other crop and environmental conditions which may be conducive to increased natural sunburn. <p>DO NOT combine ROLLING 72 SC with anything except water for application to watermelons unless your prior use has shown the combination to be non-injurious to watermelons under your conditions of use.</p> <p>Apply by ground, air or chemigation.</p>
	Cercospora leaf spot (<i>C. citrullina</i>)	2 – 3 (1.5 – 2.25)	
	Gummy stem blight/vine decline (<i>Didymella bryoniae</i>)		
	Alternaria leaf blight (<i>A. cucumerina</i>)		
	Alternaria leaf spot (<i>A. alternata</i>)		
	Scab (<i>Cladosporium cucumerinum</i>) Powdery mildew (<i>Sphaerotheca</i> only)		
<p>Restrictions and Limitations:</p> <ul style="list-style-type: none"> • DO NOT apply more than 21 pints ROLLING 72 SC (15.75 lbs a.i.) per acre during each growing season. • ROLLING 72 SC may be applied the day of harvest. 			
Grasses Grown for Seed	Stem rust	1 – 1 ½ (0.75 – 1.125)	<p>Apply with sufficient water to obtain adequate coverage. Begin applications during stem elongation when conditions favor disease development. Re-apply at flag (top) leaf emergence and repeat at 14 day intervals (the minimum re-treatment interval is 14 days).</p> <p>Apply by ground, air or chemigation.</p>
	Leaf rust		
	Stripe rust		
	Septoria leaf spot		
	Glume blotch		
	Bipolaris and Drechslera leaf spots		
	Selenophoma (eyespot)	1 – 2 (0.75 – 1.5)	
<p>Restrictions and Limitations:</p> <ul style="list-style-type: none"> • DO NOT apply more than 6 pints ROLLING 72 SC (4.5 lbs a.i.) per acre during each growing season. • DO NOT apply within 14 days of harvest. • DO NOT allow livestock to graze in treated areas or feed hay produced before harvest. Feeding of treated plant parts after harvest of seed is allowed. 			
Mango	Anthracnose (<i>Colletotrichum</i> spp.)	2 – 3 ½ (1.5 – 2.6)	<p>Apply in a water volume of 20 to 300 gallons per acre. Begin applications at early bloom and repeat on a 7 - 14 day interval until early fruit development. Begin the season with the 2 pint rate</p>

Crop	Diseases (Pathogen)	Pints/Acre (lbs a.i./Acre)	Application Directions												
			<p>on a 14 day interval (the minimum re-treatment interval is 7 days). If disease pressure is severe, use the higher rate and shorter interval.</p> <p>Ground or air applications only.</p>												
<p>Restrictions and Limitations:</p> <ul style="list-style-type: none"> DO NOT apply more than 32 pints ROLLING 72 SC (24 lbs a.i.) per acre during each growing season. DO NOT apply within 21 days of harvest. 															
<p>Mint (Indiana, Michigan and Wisconsin only)</p>	<p>Rust (<i>Puccinia menthae</i>)</p> <p>Septoria leaf spot (<i>S. menthae</i>)</p>	<p>1 3/8 (1.0)</p>	<p>Apply with sufficient water to obtain adequate coverage, normally 20 to 150 gallons per acre for dilute sprays and 5 to 10 gallons per acre for concentrate ground and aircraft applications. Begin applications when emerging plants are 4 to 8 inches high and repeat at 7 to 10 day intervals to maintain control (the minimum re-treatment interval is 7 days).</p>												
<p>Restrictions and Limitations:</p> <ul style="list-style-type: none"> DO NOT apply more than 4 pints ROLLING 72 SC (3 lbs a.i.) per acre during each growing season. DO NOT apply within 80 days of harvest. DO NOT feed fresh or extracted mint hay from treated fields to livestock. 															
<p>Onion (Dry bulb) Garlic</p>	<p>Botrytis leaf blight (<i>Botrytis</i> spp.)</p> <p>Purple blotch (<i>Alternaria porri</i>)</p> <p>Suppression: Botrytis neck rot</p> <p>Downy mildew (<i>Peronospora destructor</i>)</p>	<p>1 – 3 (0.75 – 2.25)</p>	<p>Apply with sufficient water to obtain thorough coverage of tops. ROLLING 72 SC is recommended for use with disease monitoring systems which adjust fungicide rates and frequency of application according to disease hazard. Apply as follows:</p> <table border="1" data-bbox="971 1308 1409 1549"> <thead> <tr> <th></th> <th>Low Disease Hazard & Prior to Infection</th> <th>Low Disease Hazard & Some Disease Present</th> <th>High Disease Hazard</th> </tr> </thead> <tbody> <tr> <td>Rate per Acre</td> <td>1 pt.</td> <td>1 3/8 pts.</td> <td>3 pts.</td> </tr> <tr> <td>Frequency</td> <td>10 days</td> <td>7 to 10 days</td> <td>7 days</td> </tr> </tbody> </table> <p>For suppression of neck rot (<i>Botrytis</i> spp.) during storage, a minimum of three weekly applications prior to lifting, using 1 3/8 to 3 pints of ROLLING 72 SC per acre is recommended.</p> <p>The minimum re-treatment interval is 7 days.</p> <p>Apply by ground, air or chemigation.</p>		Low Disease Hazard & Prior to Infection	Low Disease Hazard & Some Disease Present	High Disease Hazard	Rate per Acre	1 pt.	1 3/8 pts.	3 pts.	Frequency	10 days	7 to 10 days	7 days
	Low Disease Hazard & Prior to Infection	Low Disease Hazard & Some Disease Present	High Disease Hazard												
Rate per Acre	1 pt.	1 3/8 pts.	3 pts.												
Frequency	10 days	7 to 10 days	7 days												

Crop	Diseases (Pathogen)	Pints/Acre (lbs a.i./Acre)	Application Directions
Restrictions and Limitations: <ul style="list-style-type: none"> DO NOT apply more than 20 pints ROLLING 72 SC (15 lbs a.i.) per acre during each growing season. DO NOT apply within 7 days of harvest. 			
Onion (green bunching) Leek Shallots Onion and Garlic (grown for seed)	Botrytis leaf blight (<i>Botrytis</i> spp.) Purple blotch (<i>Alternaria porri</i>) Suppression: Downy mildew (<i>Peronospora destructor</i>)	1 ½ - 3 (1.125 – 2.25)	Apply with sufficient water to obtain thorough coverage of tops. Begin applications prior to favorable infection periods, and repeat at 7 - 10 day intervals for as long as conditions favor disease (the minimum re-treatment interval is 7 days). Use the high rate and a 7 day schedule of applications when heavy dew or rains persist. Apply by ground, air or chemigation.
Restrictions and Limitations: <ul style="list-style-type: none"> DO NOT apply more than 9 pints ROLLING 72 SC (6.75 lbs a.i.) per acre during each growing season. DO NOT apply within 7 days of harvest on garlic. DO NOT apply within 14 days of harvest on green bunching onions, leeks or shallots. 			
Papaya	Alternaria fruit spot (<i>A. alternata</i>) Anthracnose (<i>Colletotrichum</i> spp.) Stem end rot (<i>A. alternata</i> , <i>Colletotrichum</i> spp.)	1 ½ - 3 (1.125 – 2.25)	Apply by ground only, in sufficient water to obtain adequate coverage of fruit and leaves. Begin treatment when conditions favor development of disease and continue treatments at 14 day intervals until weather conditions no longer favor disease development (the minimum re-treatment interval is 14 days.)
Restrictions and Limitations: <ul style="list-style-type: none"> DO NOT apply more than 9 pints ROLLING 72 SC (6.75 lbs a.i.) per acre during each growing season. ROLLING 72 SC may be applied the day of harvest. 			
Parsnip	Alternaria leaf spot (<i>Alternaria</i> spp.) Downy mildew (<i>Plasmopara crustosa</i>) Anthracnose (<i>Colletotrichum</i> spp.) Botrytis blight (gray mold) (<i>B. cinerea</i>) Bottom rot (<i>Rhizoctonia</i>)	1 ½ - 2 (1.125 – 1.5)	Apply with sufficient water to obtain adequate coverage. Make the first application at the first sign of disease or when conditions are favorable for infection. Continue applications on a 7 - 10 day schedule (the minimum re-treatment interval is 7 days). Apply by ground, air or chemigation.
Restrictions and Limitations: <ul style="list-style-type: none"> DO NOT apply more than 8 pints ROLLING 72 SC (6 lbs a.i.) per acre during each growing season. DO NOT apply within 10 days of harvest. 			

Crop	Diseases (Pathogen)	Pints/Acre (lbs a.i./Acre)	Application Directions
Passion Fruit (Hawaii only)	Alternaria fruit and leaf spot (<i>Alternaria</i> spp.) Anthracnose (<i>Colletotrichum</i> spp.) Cercospora fruit spot	2 (1.5)	Apply by ground only in sufficient water to obtain adequate coverage of fruit and leaves. Begin applications during late bloom and repeat at 14 day intervals until weather conditions no longer favor disease development (the minimum re-treatment interval is 14' days.)
Restrictions and Limitations: <ul style="list-style-type: none"> DO NOT apply more than 10 pints ROLLING 72 SC (7.5 lbs a.i.) per acre during each growing season. DO NOT apply within 7 days of harvest. 			
Peanut	Early leaf spot (<i>Cercospora arachidicola</i>) Late leaf spot (<i>Cercosporidium personatum</i>) Pepper spot (<i>Leptosphaerulina crassiasca</i>)	1 – 1 ½ (0.75 – 1.125)	Apply with sufficient water for coverage when leaf wetness first occurs or 30 to 40 days after planting; repeat at 14 day intervals (the minimum re-treatment interval is 14 days). When conditions favor late leaf spot or when rust or web blotch occur, apply 1 ½ pints ROLLING 72 SC per acre at 14 day intervals for the remainder of the season.
	Rust (<i>Puccinia arachidis</i>) Web blotch (<i>Phoma arachidicola</i>)	1 ½ (1.125)	Apply by ground, air, or chemigation. If applying by chemigation, use 1 ½ pints ROLLING 72 SC per acre. It is recommended to alternate chemigation applications with ground or aerial applications.
Restrictions and Limitations: <ul style="list-style-type: none"> DO NOT apply more than 12 pints ROLLING 72 SC (9 lbs a.i.) per acre during each growing season. DO NOT apply within 14 days of harvest. DO NOT allow livestock to graze in treated areas. DO NOT feed hay or threshings from treated fields to livestock. 			
Potato	Late blight (<i>Phytophthora infestans</i>) Early blight (<i>Alternaria solani</i>) Botrytis vine rot (<i>B. cinerea</i>) Black dot (<i>Colletotrichum coccodes</i>)	¾ (0.6) then 1 – 1 ½ (0.75 – 1.125)	Begin applications at the low rate when vines are first exposed and leaf wetness occurs. Repeat applications at 5 - 10 day intervals (the minimum re-treatment interval is 5 days). Begin applying the higher label rates at 5 - 10 days intervals when any one of the following events occur: <ul style="list-style-type: none"> Vines close within the rows Late blight forecasting measures 18 disease severity values (DSV) The crop reaches 300 P-days Increase water spray volume as canopy density increases. Use the highest rate and shortest interval when plants are rapidly growing and disease

Crop	Diseases (Pathogen)	Pints/Acre (lbs a.i./Acre)	Application Directions
			<p>conditions are severe.</p> <p>Apply by ground, air, or chemigation. DO NOT exceed a 10 day interval between applications when using chemigation.</p>

Restrictions and Limitations:

- DO NOT apply more than 15 pints ~~ROLLING 72 SC~~ (11.25 lbs a.i.) per acre during each growing season.
- DO NOT apply within 7 days of harvest.

Soybean	<p>Anthracnose (<i>Colletotrichum truncatum</i>)</p> <p>Diaprothe pod and stem rot (<i>D. phaseolorum</i>)</p> <p>Frogeye leaf spot (<i>Cercospora sojina</i>)</p> <p>Purple seed stain (<i>C. kikuchii</i>)</p>		<p>Apply with sufficient water to obtain complete coverage, using at least 5 gallons of water per acre for aerial application. Use the 3 application program in areas having a history of moderate to severe disease intensity. The minimum re-treatment interval is 14 days.</p> <p>Apply by ground, air, or chemigation.</p>
	<p>Cercospora leaf blight (<i>C. kikuchii</i>)</p> <p>Septoria brown spot (<i>S. glycines</i>)</p> <p>Suppression: Rust (<i>Phakipsora pachyrhizi</i>)</p>	<p>1 ½ - 2 ¼ (1.125 - 1.7)</p>	<p>Two application program: For determinate varieties, make the first application at early pod set stage (R3) and the second application at seed formation (R5). For indeterminate varieties, make the first application when largest pods are 1 - 1 ¼ inches in length. Make the second application 14 days later.</p>
		<p>1 - 2 (0.75 - 1.5)</p>	<p>Three application program: For determinate varieties, make the first application at the beginning of flowering (R1), the second at early pod set (R3), and the third at beginning of seed formation (R5). For indeterminate varieties, make the first application one week after first flowering and continue applications at 14 day intervals.</p>
	<p>Stem canker (<i>Diaporthe phaseolorum</i>)</p>	<p>1 (0.75)</p>	<p>Apply with 10 to 20 gallons of water per acre, as a band treatment directing spray to provide coverage of entire plant. Make the first application at time of emergence of the second trifoliate leaves (V2). If conditions favor stem canker disease make a second and third application. Make all applications at 14 day intervals.</p>

Restrictions and Limitations:

- DO NOT apply more than 6 pints ~~ROLLING 72 SC~~ (4.5 lbs a.i.) per acre during each growing season.
- DO NOT apply within 6 weeks of harvest.

Crop	Diseases (Pathogen)	Pints/Acre (lbs a.i./Acre)	Application Directions
<ul style="list-style-type: none"> DO NOT feed hay or threshings from treated fields to livestock. 			
Tomato	<u>FOLIAGE</u> Early blight <i>(Alternaria solani)</i> Late blight <i>(Phytophthora infestans)</i> Gray leaf spot <i>(Stemphyllium botryosum)</i> Gray leaf mold <i>(Fluvia fluva; Cladosporium)</i> Septoria leaf spot <i>(S. lycopersici)</i> Target spot <i>(Corynespora cassiicola)</i>	1 3/8 – 2 (1.0 – 1.5)	Apply with sufficient water to obtain adequate coverage. Begin applications when dew or rain occurs and disease threatens. Apply on a 7 - 10 day interval for foliage diseases. For fruit diseases, begin at fruit set and apply on a 7 - 14 day interval. Use the highest rate and shortest interval specified when disease conditions are severe. The minimum re-treatment interval is 7 days. Apply by ground, air, or chemigation.
	<u>FRUIT</u> Anthracnose <i>(Colletotrichum spp.)</i> Alternaria fruit rot (black mold) <i>(A. alternata)</i> Botrytis gray mold <i>(B. cinerea)</i> Late blight fruit rot <i>(P. infestans)</i> Rhizoctonia fruit rot <i>(R. solani)</i>	2 – 2 3/4 (1.5 – 2.1)	
Restrictions and Limitations: <ul style="list-style-type: none"> DO NOT apply more than 20 pints ROLLING 72 SC (15 lbs a.i.) per acre during each growing season. ROLLING 72 SC may be applied the day of harvest. 			

DIRECTIONS FOR APPLICATION - TREE AND ORCHARD CROPS

Apply **ROLLING 72 SC** in sufficient water and with proper calibration to obtain uniform coverage of tree canopy. For fruit and nut bearing crops, the maximum volume is 300 gallons per acre unless indicated otherwise in the specific use directions. For conifers, the maximum volume is 100 gallons per acre.

Application with ground equipment is preferable to aerial application because ground applications generally give better coverage of the tree canopy. If application with ground equipment is not feasible, **ROLLING 72 SC** may be applied with aircraft using at least 20 gallons of spray per acre. The minimum volume for application by aircraft to forest stands and Christmas trees is 10 gallons per acre. When concentrate sprays are used or when treating non-bearing or immature trees, the lower rate of **ROLLING 72 SC** listed may be used.

DO NOT allow livestock to graze in treated areas.

Crop	Diseases (Pathogens)	ROLLING 72 SC Use Rate		Application Instructions
		Pints/Acre (lbs a.i./acre)	100 Gallons*	
Almonds	Blossom blight/ brown rot (<i>Monilinia</i> spp.)	4 (3.0)	1.33 (1.0)	Apply with water volumes of 20 – 300 gallons per acre. For blossom blight, begin application at popcorn (pink bud) and follow with an application at full bloom. If weather is still conducive for disease development, another application may be made at petal fall. For control of shothole, make an application in the autumn at leaf fall. In the spring, make the first application at budbreak, followed by an application at shuck split to control nut infections and to control scab. Ground or air applications only.
	Shothole (<i>Wilsonomyces carpophilus</i>)			
	Scab (<i>Venturia carpophila</i>)			
Restrictions and Limitations:				
<ul style="list-style-type: none"> DO NOT apply more than 25 pints ROLLING 72 SC (18.75 lbs a.i.) per acre during each growing season (leaf fall through shuck split). DO NOT apply within 150 days of harvest. 				
Filberts (Hazelnuts)	Eastern filbert blight (<i>Anisogramma anomala</i>)	4 (3.0)	1.33 (1.0)	Apply with water volumes of 20 – 300 gallons per acre. Begin applications at the onset of disease or when weather conditions favor disease development. Make applications on a 14 - 28 day schedule, using the shorter interval under heavy disease pressure (the minimum re-treatment interval is 14 days).
Restrictions and Limitations:				
<ul style="list-style-type: none"> DO NOT apply more than 12 pints ROLLING 72 SC (9 lbs a.i.) per acre during each growing season. DO NOT apply within 120 days of harvest. DO NOT apply through irrigation. DO NOT apply with oils, other pesticides, surfactants or fertilizers. DO NOT apply within one week of an oil-based pesticide application. 				
Peach	Leaf curl	3 1/8 - 4 1/8	1 – 1 3/8	For best control apply at leaf fall in late

23
43

Crop	Diseases (Pathogens)	ROLLING 72 SC Use Rate		Application Instructions
		Pints/Acre (lbs a.i./acre)	100 Gallons*	
Nectarine Apricot Cherry Plum Prune	(<i>Taphrina deformans</i>) Shot hole (<i>Wilsonomyces carpophilus</i>)	(2.3 – 3.1)	(0.75 – 1.0)	autumn, using sufficient water and proper sprayer calibration to obtain uniform coverage. When conditions favor high disease levels, use the high rate of application and apply once or twice more in mid to late winter before budswell. If the leaf fall application is not practical, application of ROLLING 72 SC for control of leaf curl may be made at any time prior to budswell the following spring. Where shothole occurs, also apply at budbreak to protect newly emerging leaves and at shuck split to prevent fruit infections. Ground or air applications only.
	Lacy (russet) scab (Plum / Prune) Brown rot blossom blight (<i>Monilinia</i> spp.)	3 1/8 - 4 1/8 (2.3 – 3.1)	1 – 1 3/8 (0.75 – 1.0)	Make one application at popcorn (pink, red or early white bud) and a second application at full bloom. If weather conditions favor disease development, make an additional application at petal fall.
	Cherry leaf spot (<i>Blumeriella jaapii</i>) Scab (<i>Cladosporium carpophilum</i>) Black knot (Cherry, Plum) (<i>Apiosporina morbosa</i>)	3 1/8 - 4 1/8 (2.3 – 3.1)	1 – 1 3/8 (0.75 – 1.0)	In addition to the bloom application listed above, make one application at shuck split. Do not apply ROLLING 72 SC after shuck split and before harvest. If additional disease control is needed before harvest, use another registered fungicide. For control of cherry leaf spot after harvest, make one application to foliage within 7 days after fruit is removed. In orchards with a history of high leaf spot incidence, make a second application 10 – 14 days later. Ground or air applications only.
Restrictions and Limitations: <ul style="list-style-type: none"> DO NOT apply more than 20.5 pints ROLLING 72 SC (15.4 lbs a.i.) per acre during each growing season. ROLLING 72 SC may be applied the day of harvest. The minimum re-treatment interval is 10 days. 				
Pistachio	Botryosphaeria blight (<i>B. dothidea</i>)	6 (4.5)	3 (2.25)	Apply with water volumes of 20 – 200 gallons per acre. Make the first application at the beginning of the blossom period followed by an application at full bloom. Make additional applications as required on a 28 day interval. (The minimum re-treatment interval is 28 days). For Septoria and Botrytis, use the higher rate if disease pressure is severe.
	Suppression: Alternaria late blight (<i>A. alternata</i>) Septoria leaf spot	4 – 6	2 – 3	

Crop	Diseases (Pathogens)	ROLLING 72 SC Use Rate		Application Instructions
		Pints/Acre (lbs a.i./acre)	100 Gallons*	
	(<i>S. pistacina</i>) Botrytis blight (<i>B. cinerea</i>)	(3.0 - 4.5)	(1.50 - 2.25)	<p>Note: Use of this product may result in speckling or reddening of the fruit hull (epicarp). This effect is superficial and has not resulted in any change in nut quality.</p> <p>Ground or air applications only.</p>

Restrictions and Limitations:

- DO NOT apply more than 30 pints ROLLING 72 SC (22.5 lbs a.i.) per acre during each growing season.
- DO NOT apply within 14 days of harvest.

Conifers (Pines, Spruces)	Swiss needlecast (<i>Phaeocryptopus gaeumannii</i>)	2 ¾ - 5 ½ (2.1 - 4.125)	2 ¾ - 5 ½ (2.1 - 4.125)	<p>Single application technique: In Christmas tree plantations or forest stands, make one application in the spring when new shoot growth is ½ to 2 inches in length.</p>
	Scleroderris canker (Pines) (<i>Gremmeniella abietina</i>) Swiss needlecast (<i>Phaeocryptopus gaeumannii</i>)	1 ½ - 2 ¾ (1.125 - 2.1)	1 ½ - 2 ¾ (1.125 - 2.1)	
	Sirococcus tip blight (<i>S. conigenus</i>)	2 - 3 ½ (1.5 - 2.6)	2 - 3 ½ (1.5 - 2.6)	
	Rhizosphaera needlecast (Spruces) (<i>Rhizosphaera</i> spp.)	5 ½ (4.125)	5 ½ (4.125)	
	Scirrhia brown spot (Pines) (<i>Mycosphaerella dearnessii</i>)			
	Cyclaneusma and Lophodermium needlecasts (Pines)	2 ¾ - 5 ½ (2.1 - 4.125)	2 ¾ - 5 ½ (2.1 - 4.125)	<p>Apply in early spring prior to budbreak. Repeat applications at approximately 6 - 8 week intervals, until spore release ceases in late fall. Apply monthly during periods of frequent rainfall, and where Lophodermium infections occur during dormancy (Pacific Northwest). During drought periods, applications may be suspended, then resumed upon next occurrence of needle wetness.</p>
	Rhabdocline needlecast	1 ½ - 2 ¾ (1.125 - 2.1)	1 ½ - 2 ¾ (1.125 - 2.1)	

25
43

Crop	Diseases (Pathogens)	ROLLING 72 SC Use Rate		Application Instructions
		Pints/Acre (lbs a.i./acre)	100 Gallons*	
	(Douglas-fir)			elongated and conditions no longer favor disease development. In plantations of mixed provenance, or when irregular budbreak occurs, apply weekly until all trees have broken bud, then every 3 to 4 weeks as specified above. In nursery beds, use the high rate on a 3-week schedule.
	Botrytis seedling blight Phoma twig blight	1 ½ - 2 ¾ (1.125 - 2.1)	1 ½ - 2 ¾ (1.125 - 2.1)	Begin applications in nursery beds when seedlings are 4 inches tall and when cool, moist conditions favor disease development. Make additional applications at 7 - 14 day intervals as long as disease favorable conditions persist.
	Autoecious needle rust (Weir's cushion) (Spruce)	5 ½ (4.125)	5 ½ (4.125)	Begin applications when 10% of buds have broken and twice thereafter at 7 - 10 day intervals.

Restrictions and Limitations:

- DO NOT apply more than 22 pints ROLLING 72 SC (16.5 lbs a.i.) per acre during each growing season.
- The minimum re-treatment interval is 21 days. The minimum re-treatment interval in nursery beds is 7 days.

*Volumetric rates to be used only with full dilute spray volume specified on this label for tree and orchard crops.

Crop	Disease	ROLLING 72 SC Use Rate	Application Instructions
		fl oz / 1000 sq ft	
Mushroom	Verticillium brown spot Dry bubble	2.75 - 5.5	Apply as a drench to the mushroom bed surface in at least 12.5 gallons of water per 1,000 sq. ft. of mushroom bed. Make two applications. Apply the high rate (5.5 fl oz) of ROLLING 72 SC in the first application and the low rate (2.75 fl oz) of ROLLING 72 SC in the second application. The first application should be made within two days of top-dressing the spawn-colonized mushroom compost with a casing layer. The second application should be made at pinning.

Restrictions and Limitations:

- DO NOT apply within 5 days of first harvest.
- DO NOT make more than two applications per cropping cycle.
- DO NOT apply more than 8.25 fl oz of ROLLING 72 SC per 100 sq ft per cropping cycle.

Warranty and Disclaimer Statement

The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Such risks may arise from weather conditions, soil factors, off-target movement, unconventional farming techniques, the presence of other materials, the manner of use or application, or other unknown factors, all of which are beyond the control of Arysta LifeScience North America, LLC ("Arysta"), and can cause crop injury, injury to non-target crops or

plants, ineffectiveness of the product, or other unintended consequences. All such risks shall be assumed by the user or buyer. Arysta warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks described above, when used in accordance with the Directions for Use under normal conditions. This warranty does not extend to the use of this product contrary to label instructions or under conditions not reasonably foreseeable to Arysta, and is subject to the inherent risks described above.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, ARYSTA DISCLAIMS ALL OTHER WARRANTIES, EXPRESSED OR IMPLIED, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, ARYSTA, MANUFACTURER, AND SELLER DISCLAIM AND SHALL NOT BE LIABLE FOR ANY SPECIAL, INCIDENTAL, INDIRECT, OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE, HANDLING, APPLICATION, STORAGE, OR DISPOSAL OF THIS PRODUCT OR FOR DAMAGES IN THE NATURE OF PENALTIES, AND THE USER AND BUYER WAIVE ANY RIGHT THAT THEY MAY HAVE TO SUCH DAMAGES. NO AGENT, REPRESENTATIVE OR EMPLOYEE OF ARYSTA IS AUTHORIZED TO MAKE ANY WARRANTY, GUARANTEE OR REPRESENTATION BEYOND THOSE CONTAINED HEREIN OR TO MODIFY THE WARRANTIES CONTAINED HEREIN.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE TOTAL LIABILITY OF ARYSTA, MANUFACTURER, AND SELLER, SHALL BE LIMITED TO THE PURCHASE PRICE PAID, OR AT ARYSTA'S ELECTION, THE REPLACEMENT OF THE PRODUCT.

Dipel is a registered trademark of Valent BioSciences Corporation
Latron B-1956 and latron AG-98 are registered trademarks of Dow AgroSciences LLC
Arysta LifeScience North America, LLC

27
43

ROLLING 72 SC

Flowable Fungicide

Turf and Ornamental Fungicide

ACTIVE INGREDIENT:

ROLLING 72 SC (tetrachloroisophthalonitrile) 54.0%

INERT INGREDIENTS: 46.0%

TOTAL: 100.0%

This product contains 6.0 pounds of ROLLING 72 SC per gallon (720 g per liter)

KEEP OUT OF REACH OF CHILDREN

CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label find someone to explain it to you in detail.)

See inside booklet for additional precautionary statements

For Product Use Information Call 1-866-761-9397

Arysta LifeScience North America, LLC
15401 Weston Parkway, Suite 150
Cary, NC 27513

EPA Reg. No. 66330-362
EPA Est. No.

NET WEIGHT: _____

28
43

FIRST AID	
IF SWALLOWED	<ul style="list-style-type: none"> • Call a poison control center or doctor for treatment advice. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Have person sip a glass of water if able to swallow.
IF ON SKIN OR CLOTHING	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin with plenty of water for 15–20 minutes. • Get medical attention if irritation persists.
IF INHALED	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth to mouth, if possible. • Call a poison control center or doctor for further treatment advice.
IF IN EYES	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. • Call a physician if irritation persists.
NOTE TO PHYSICIAN	
Persons suffering with temporary allergic skin reactions may respond to treatment with oral antihistamines and topical or oral steroids.	
Have the product container or label with you when calling a poison control center or doctor or going for treatment.	
HOTLINE NUMBERS:	
FOR 24-HOUR MEDICAL EMERGENCY ASSISTANCE:	
Call PROSAR at 1-866-303-6952 or 1-651-8946 if calling from outside of the U.S.	
FOR 24-HOUR CHEMICAL EMERGENCY ASSISTANCE:	
Spill, leak, fire, exposure, or accident	
Call CHEMTREC at 1-800-424-9300 or 1-703-527-3887 if calling from outside of the U.S.	

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION. Harmful if swallowed, absorbed through skin, or inhaled. Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Avoid breathing spray mist. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical resistant to this product are made of any waterproof material. If you want more options, follow the instructions for category A on an EPA chemical resistance category selection chart.

Mixers, loaders, applicators and all other handlers who handle this pesticide must wear:

- Long-sleeved shirt and long pants
- Chemical resistant gloves made of any waterproof material – Category A (e.g. barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, natural rubber, polyethylene, polyvinyl chloride (PVS) or viton)
- Shoes plus socks
- A NIOSH approved dust/mist filtering respirator (MSHA/NIOSH approval number prefix TC-21C) or a NIOSH approved respirator with any N, R, P or HE filter for applicators and handlers in enclosed areas such as a greenhouse.

29
43

USER SAFETY REQUIREMENTS

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing or other materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

ENGINEERING CONTROLS STATEMENT

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to aquatic invertebrates and wildlife. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwater or rinsate.

This chemical is known to leach through soil into groundwater under certain conditions as a result of label use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

This chemical can contaminate surface water through spray drift. Under some conditions, it may also have a high potential for runoff into surface water for several days to weeks after application. These include poorly draining or wet soils with readily visible slopes toward adjacent surface waters, frequently flooded areas, areas overlaying extremely shallow ground water, areas with infield canals or ditches that drain to surface water, areas not separated from adjacent surface waters with vegetated filter strips, and areas over-laying tile drainage systems that drain to surface water.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, or pets either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of **12 hours**.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is: long-sleeved shirt and long pants or coveralls, shoes plus socks, chemical resistant gloves made of any waterproof material, such as nitrile, butyl, neoprene, and / or barrier laminate, and protective eyewear.

Special Eye Irritation Provisions: This product is a severe eye irritant. Although the restricted entry interval expires after 12 hours, for the next 6.5 days entry is permitted only when the following safety measures are provided:

1. At least one container designed specifically for flushing eyes must be available in operating condition at the WPS required decontamination site intended for workers entering the treated area.
2. Workers must be informed, in a manner they can understand:
 - That residues in the treated area may be highly irritating to their eyes
 - That they should take precautions, such as refraining from rubbing their eyes, to keep the residues out of their eyes
 - That if they do get residues in their eyes, they should immediately flush their eyes using the eyeflush container that is located at the decontamination site or using other readily available clean water, and
 - How to operate the eyeflush container

NON-AGRICULTURAL USE REQUIREMENTS

THE REQUIREMENTS IN THIS BOX APPLY TO USES OF THIS PRODUCT THAT ARE not WITHIN THE SCOPE OF THE Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not enter or allow others to enter the treated area until sprays have dried.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE

Store in original container and keep tightly closed. Store in a cool dry place. Protect from excessive heat.

PESTICIDE DISPOSAL

Pesticide wastes are toxic. Improper disposal of excess pesticide, pesticide spray or rinsate is a violation of federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL

Plastic Containers equal to or less than 5 gallons: Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip.

Plastic Containers greater than 5 gallons: Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth ensuring at least one complete revolution, for 30 seconds. Stand

31
43

the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

Offer for recycling, if available, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Returnable Refillable Containers/ Bulk and Minibulk Containers:

Return container intact to point of purchase. Refillable container. Refill this container with pesticide only. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

GENERAL INFORMATION

ROLLING 72 SC is an excellent disease control agent when used according to label directions for control of a broad spectrum of plant diseases. ROLLING 72 SC is recommended for use in programs that are compatible with the principles of Integrated Pest Management (IPM), which include the use of disease resistant crop varieties, cultural practices, post scouting and disease forecasting systems, which reduce unnecessary applications of pesticides.

ROLLING 72 SC is effective in programs to minimize disease resistance to fungicides. ROLLING 72 SC, with a multi-site mode of action, may be used to delay or prevent the development of resistance to single-site fungicides. Consult your federal or state Cooperative Extension Service representatives for guidance on the proper use of ROLLING 72 SC in programs which seek to minimize the occurrence of disease resistance to other fungicides.

ROLLING 72 SC can be used effectively in dilute or concentrate sprays. Thorough, uniform coverage is essential for disease control.

GENERAL PRECAUTIONS AND RESTRICTIONS

Do not use on home lawns and turf sites associated with apartment buildings, day care centers, playgrounds, play fields, recreational park athletic fields, athletic fields located on or next to schools (i.e. elementary, middle and high schools), campgrounds, churches and theme parks.

Agricultural Use Sites Only (sod farms, golf courses, nurseries and greenhouses): This product must not be applied within 150 feet (for aerial applications) or 25 feet (for ground applications) of marine/estuarine water bodies unless there is an untreated buffer area of that width between the area to be treated and the water body.

Do not combine ROLLING 72 SC in the spray tank with pesticides, surfactants or fertilizers, unless prior use has shown the combination physically compatible, effective and noninjurious under your conditions of use. Do not combine ROLLING 72 SC with DiPel® or Latron B-1956® as phytotoxicity may result from the combination when applied to some species on this label.

The required amount of ROLLING 72 SC should be added slowly into the spray tank during filling. With concentrate sprays, pre-mix the required amount of ROLLING 72 SC in a clean container and add to the spray tank as it is being filled. Keep agitator running when filling spray tank and during spray operations.

SPRAY DRIFT PRECAUTIONS

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment and weather related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions.

The following drift management requirements must be followed to avoid off-target drift movement from aerial applications to agricultural use sites. These requirements do not apply to forestry applications, public health uses or to applications using dry formulation.

1. The distance of the outer most nozzles on the boom must not exceed ¼ the length of the wingspan or rotor.

- 2. Nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees.

Where states have more stringent regulations, they should be observed.

The applicator should be familiar with and take into account the information covered in the *Aerial Drift Reduction Advisory Information*.

Aerial Drift Reduction Advisory Information

This section is advisory in nature and does not supersede the mandatory label requirements.

Information on Droplet Size

The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions (see Wind, Temperature and Humidity, and Temperature Inversions below).

Controlling Droplet Size

- **Volume** - Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.
- **Pressure** - DO NOT exceed the nozzle manufacturer's recommended pressures. For many nozzle types lower pressure produces larger droplets. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.
- **Number of nozzles** - Use the minimum number of nozzles that provide uniform coverage.
- **Nozzle Orientation** - Orienting nozzles so that the spray is released parallel to the airstream produces larger droplets than other orientations and is the recommended practice. Significant deflection from horizontal will reduce droplet size and increase drift potential.
- **Nozzle Type** - Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce the largest droplets and the lowest drift.

Boom Length

For some use patterns, reducing the effective boom length to less than ¾ of the wingspan or rotor length may further reduce drift without reducing swath width.

Application Height

Applications should not be made at a height greater than 10 feet above the top of the largest plants unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind.

Swath Adjustment

When applications are made with a crosswind, the swath will be displaced downwind. Therefore, on the upwind and downwind edges of the field, the applicator should compensate for this displacement by adjusting the path of the aircraft upwind. Swath adjustment distance should increase, with increasing drift potential (higher wind, smaller drops, etc.)

Wind

Drift potential is lowest between wind speeds of 2 - 10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given speed. Application should be avoided below 2 mph due to variable wind direction and high inversion potential. **NOTE:** Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

Temperature and Humidity

When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

Temperature Inversions

Applications should not occur during a temperature inversion because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain in a concentrated cloud.

This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

Sensitive Areas

The pesticide should only be applied when the potential for drift to adjacent sensitive areas (e.g., residential areas, bodies of water, known habitat for threatened or endangered species, nontarget crops) is minimal (e.g., when wind is blowing away from the sensitive areas).

APPLICATION INSTRUCTIONS

APPLICATION AND CALIBRATION TECHNIQUES FOR SPRINKLER IRRIGATION – CHEMIGATION:

Apply this product only through sprinkler irrigation systems including center pivot, motorized lateral move, solid set or portable (wheel move, side roll, end tow, or hand move) irrigation system(s). Do not apply this product through any other type of irrigation system. Do not use ROLLING 72 SC through sprinkler irrigation equipment on golf courses.

Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.

If you have questions about calibration, you should contact State Extension Service Specialists, equipment manufacturers or other irrigation experts.

Do not apply this product through irrigation systems connected to a public water system. "Public water system" means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days per year.

Controls for both irrigation water and pesticide injection systems must be functionally interlocked, so as to automatically terminate pesticide injection when the irrigation water pump motor stops. A person knowledgeable of the irrigation system and responsible for its operation shall be present so as to discontinue pesticide injection and make necessary adjustments, should the need arise.

The irrigation water pipeline must be fitted with a functional, automatic, quick-closing check valve to prevent the flow of treated irrigation water back toward the water source. The pipeline must also be fitted with a vacuum relief valve and low pressure drain, located between the irrigation water pump and the check valve, to prevent back-siphoning of treated irrigation water into the water source.

Always inject ROLLING 72 SC into irrigation water after it discharges from the irrigation pump and after it passes through the check valve. Never inject pesticides into the intake line on the suction side of the pump.

Pesticide injection equipment must be fitted with a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump. Interlock this valve to the power system, so as to prevent fluid from being withdrawn from the chemical supply tank when the irrigation system is either automatically or manually turned off.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Spray mixture in the chemical supply tank must be agitated at all times, otherwise settling and uneven application may occur. Do not apply when wind speed favors drift beyond the area intended for treatment.

Posting of areas to be chemigated is required when 1) any part of a treated area is within 300 feet of sensitive areas such as residential areas, labor camps, businesses, day care centers, hospitals, in-patient clinics, nursing homes or any public areas such as schools, parks, playgrounds, or other public facilities not including public roads, or 2) when the chemigated area is open to the public.

Posting must conform to the following requirements. Treated areas shall be posted with signs at all usual points of entry and along likely routes of approach from the listed sensitive areas. When there are no usual points of entry, signs must be posted in the corners of the treated areas and in any other location affording maximum visibility to sensitive areas. The printed side of the sign should face away from the treated area towards the sensitive area. The signs shall be printed in English. Signs must be posted prior to application and must remain posted until foliage has dried and soil surface water has disappeared. Signs may remain in place indefinitely as long as they are composed of materials to prevent deterioration and maintain legibility for the duration of the posting period.

All words shall consist of letters at least 2 1/2 inches tall, and all letters and the symbol shall be a color that sharply contrasts with their immediate background. At the top of the sign shall be the words KEEP OUT, followed by an octagonal stop sign symbol at least 8 inches in diameter containing the word STOP. Below the symbol shall be the words PESTICIDES IN IRRIGATION WATER.

The sign is in addition to any sign posted to comply with the Worker Protection Standard.

ROLLING 72 SC may be used through two basic types of sprinkler irrigation systems as outlined in Sections A and B below. Determine which type of system is in place, then refer to the appropriate directions provided for each type.

A. Center Pivot, Motorized Lateral Move and Traveling Gun Irrigation Equipment

For injection of pesticides, these continuously moving systems must use a positive displacement injection pump, of either diaphragm or piston type, constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock and capable of injection at pressures approximately 2 – 3 times those encountered within the irrigation water line. Venturi applicator units cannot be used on these systems.

Fill chemical supply tank of injection equipment with water. Operate system for one complete revolution or run across the field, measuring time required, amount of water injected, and acreage covered.

Thoroughly mix recommended amount of ROLLING 72 SC for acreage to be covered to same amount of water used during calibration and inject into system continuously for one revolution or run. Mixture in the chemical supply tank must be continuously agitated during the injection run. Shut off injection equipment after one revolution or run, but continue to operate irrigation system until ROLLING 72 SC has been cleared from last sprinkler head.

B. Solid Set and Portable (Wheel Move, Side Roll, End Tow, or Hand Move) Irrigation Equipment

With stationary systems, an effectively designed in-line Venturi applicator unit is preferred which is constructed of materials that are compatible with pesticides; however, a positive-displacement pump can also be used.

Determine acreage covered by sprinkler. Fill tank of injection equipment with water and adjust flow to use contents over a thirty to forty-five minute period. Mix desired amount of ROLLING 72 SC for acreage to be covered with water so that the total mixture of ROLLING 72 SC plus water in the injection tank is equal to the quantity of water used during calibration, and operate entire system at normal pressures recommended by the manufacturer of injection equipment used, for amount of time established during calibration. Agitation is recommended. ROLLING 72 SC can be injected at the beginning or end of the irrigation cycle or as a separate application. Stop injection equipment after treatment is completed and continue to operate irrigation system until ROLLING 72 SC has been cleared from last sprinkler head.

DIRECTIONS FOR APPLICATION - TURF

ROLLING 72 SC can be used to control diseases on turf on golf courses and sod farms.

35
43

Do not use on home lawns and turf sites associated with apartment buildings, day care centers, playgrounds, play fields, recreational park athletic fields, athletic fields located on or next to schools (i.e. elementary, middle and high schools), campgrounds, churches and theme parks.

Turf Group A: For Golf Course Fairways and Sod Farms:

- DO NOT use for sod farms at application rates greater than 13 lbs. a.i. per acre per year.
- For fairways, DO NOT apply more than 34.7 pints/acre (12.7 fl oz/1000 sq ft) of ROLLING 72 SC per growing season (26 lbs. a.i./acre/growing season).
- The minimum re-treatment interval for single application rates up to 9.75 pints/acre (3.6 fl oz/1000 sq ft) of ROLLING 72 SC (7.3 lbs a.i./acre) is 7 days.
- The minimum re-treatment interval after an application of a rate greater than 9.75 pints/acre (3.6 fl oz/1000 sq ft) of ROLLING 72 SC (7.3 lbs a.i./acre) is 14 days.
- DO NOT apply more than one application of a rate greater than 9.75 pints/acre (3.6 fl oz/1000 sq ft) of ROLLING 72 SC (7.3 lbs a.i./acre) per growing season.
- The maximum single application rate is 15.1 pints/acre (5.5 fl oz/1000 sq ft) of ROLLING 72 SC (11.3 lbs a.i./acre).
- Apply ROLLING 72 SC in 43.5 - 87 gallons of water per acre. Begin applications when conditions favor disease development and repeat applications as long as these conditions persist. Under severe disease conditions use the highest rate and shortest interval corresponding with the application schedule selected from the table below.
- DO NOT mow or water after treatment until spray deposited on turfgrass is thoroughly dry; ROLLING 72 SC should always be used in conjunction with good turf management practices.
- Sod farm turf treated with ROLLING 72 SC prior to harvest must be mechanically cut, rolled and harvested.

Turf Group B: For Golf Course Tees and Greens

Golf Course Tees:

- DO NOT apply more than 69.3 pints/acre (25.4 fl oz/1000 sq ft) of ROLLING 72 SC per growing season (52 lbs. a.i./acre/growing season).
- The minimum re-treatment interval for single application rate up to 9.75 pints/acre (3.6 fl oz/1000 sq ft) of ROLLING 72 SC (7.3 lbs a.i./acre) is 7 days.
- The minimum re-treatment interval after an application of a rate greater than 9.75 pints/acre (3.6 fl oz/1000 sq ft) of ROLLING 72 SC (7.3 lbs a.i./acre) is 14 days.
- DO NOT apply more than two applications of a rate greater than 9.75 pints/acre (3.6 fl oz/1000 sq ft) of ROLLING 72 SC (7.3 lbs a.i./acre) per growing season.
- The maximum single application rate is 15.1 pints/acre (5.5 fl oz/1000 sq ft) of ROLLING 72 SC (11.3 lbs a.i./acre).
- Apply ROLLING 72 SC in 43.5 - 87 gallons of water per acre. Begin applications when conditions favor disease development and repeat applications as long as these conditions persist. Under severe disease conditions use the highest rate and shortest interval corresponding with the application schedule selected from the table below.
- DO NOT mow or water after treatment until spray deposited on turfgrass is thoroughly dry; ROLLING 72 SC should always be used in conjunction with good turf management practices.

Golf Course Greens:

- DO NOT apply more than 97.3 pints/acre (35.7 fl oz/1000 sq ft) of ROLLING 72 SC per growing season (73 lbs. a.i./acre/growing season).
- The minimum re-treatment interval for single application rate up to 9.75 pints/acre (3.6 fl oz/1000 sq ft) of ROLLING 72 SC (7.3 lbs a.i./acre) is 7 days.

- The minimum re-treatment interval after an application of a rate greater than 9.75 pints/acre (3.6 fl oz/1000 sq ft) of ROLLING 72 SC (7.3 lbs a.i./acre) is 14 days.
- DO NOT apply more than two applications of a rate greater than 9.75 pints/acre (3.6 fl oz/1000 sq ft) of ROLLING 72 SC (7.3 lbs a.i./acre) per growing season.
- The maximum single application rate is 15.1 pints/acre (5.5 fl oz/1000 sq ft) of ROLLING 72 SC (11.3 lbs a.i./acre).
- Apply ROLLING 72 SC in an adequate amount of water to provide complete coverage. This amount may vary from 87 to 450 gallons per acre. See table below for suggested rates and timing. Under severe disease conditions use the highest rate and shortest interval corresponding with the application schedule selected from the table below.
- DO NOT mow or water after treatment until spray deposited on turfgrass is thoroughly dry; ROLLING 72 SC should always be used in conjunction with good turf management practices.

Diseases Controlled	App. Interval (days)	Pre-Disease Rates ¹		
		fl. oz. / 1000 sq. ft.	pints / acre	lbs. a.i. / acre
Algae²	7 – 14	2.0 – 3.6	5.5 – 9.75	4.1 – 7.3
Anthracnose (<i>Colletotrichum graminicola</i>)	7 – 14 14	3.0 – 3.6 3.6 – 5.5	8.3 – 9.75 9.9 – 15.1	6.2 – 7.3 7.4 – 11.3
Brown Patch (<i>Rhizoctonia solani</i>)	7 – 14	2.0 – 3.6	5.5 – 9.75	4.1 – 7.3
Copper Spot (<i>Gloeocercospora sorghi</i>)	14	4.0 – 5.5	11 – 15.1	8.25 – 11.3
Dichondra Leaf Spot (CA only) (<i>Alternaria</i> spp.)	14	4.0 – 5.5	11 – 15.1	8.25 – 11.3
Dollar Spot (<i>Sclerotinia homeocarpa</i> ; <i>Lanzia</i> or <i>Moellerodiscus</i> spp.)	7 – 10 7 – 21	1.0 ⁵ – 2.0 2.0 – 3.6	2.8 ⁵ – 5.0 5.5 – 9.75	2.1 ⁵ – 4.1 4.1 – 7.3
Fusarium Patch (Gerlachia)³ (<i>Microdochium nivale</i>)	21 – 28	5.5	15.1	11.3
Gray Leaf Spot (<i>Pyricularia grisea</i> , <i>P. oryzae</i>)	7 – 10	2.0 – 3.6	5.5 – 9.75	4.1 – 7.3
Gray Snow Mold⁴ (<i>Typhula</i> spp.)	30	5.5	15.1	11.3
Leaf Spot , Melting Out, Brown Blight (<i>Bipolaris sorokiniana</i> , <i>Drechslera</i> spp. including <i>D. poae</i> , <i>D. siccans</i>) <i>Curvularia</i> spp.)	7 – 10 7 – 21	2.0 2.0 – 3.6	5.5 5.5 – 9.75	4.1 4.1 – 7.3
Red Thread (<i>Laetisaria fuciformis</i>)	7 – 10 14	2.0 – 3.6 3.6 – 5.5	5.5 – 9.75 9.9 – 15.1	4.1 – 7.3 7.4 – 11.3

37
43

Stem Rust (Bluegrass) (<i>Puccinia graminis</i>)	14	4.0 – 5.5	11 – 15.1	8.25 – 11.3
--	----	-----------	-----------	-------------

Diseases Controlled	App. Interval (days)	Post-Disease Rates ¹		
		fl oz / 1000 sq ft	pints / acre	lbs a.i. / acre
Algae²	7 - 14	2.0 – 3.6	5.5 – 9.75	4.1 – 7.3
	14	4.0 – 5.5	11 – 15.1	8.25 – 11.3
Brown Patch (<i>Rhizoctonia solani</i>)	14	4.0 – 5.5	11 – 15.1	8.25 – 11.3
Copper Spot (<i>Gloeocercospora sorghi</i>)	14	5.5	15.1	11.3
Dichondra Leaf Spot (CA only) (<i>Alternaria</i> spp.)	14	5.5	15.1	11.3
Dollar Spot (<i>Sclerotinia homeocarpa</i> ; <i>Lanzia</i> or <i>Moellerodiscus</i> spp.)	14	4.0 – 5.5	11 – 15.1	8.25 – 11.3
Gray Leaf Spot (<i>Pyricularia grisea</i> , <i>P. oryzae</i>)	14	4.0 – 5.5	11 – 15.1	8.25 – 11.3
Leaf Spot , Melting Out, Brown Blight (<i>Bipolaris sorokiniana</i> , <i>Drechslera</i> spp. including <i>D. poae</i> , <i>D. siccans</i>) <i>Curvularia</i> spp.)	14	4.0 – 5.5	11 – 15.1	8.25 – 11.3
Red Thread (<i>Laetisaria fuciformis</i>)	14	5.5	15.1	11.3
Stem Rust (Bluegrass) (<i>Puccinia graminis</i>)	14	5.5	15.1	11.3

NOTES:

- Turf Group A:** Limit of one application per season at rates greater than 7.3 lbs a.i./acre (9.75 pints/acre or 3.6 fl oz/1000 sq ft) of ROLLING 72 SC.
Turf Group B: Limit of two applications per season at rates greater than 7.3 lbs a.i./acre (9.75 pints/acre or 3.6 fl oz/1000 sq ft) of ROLLING 72 SC.
- Turf Groups A & B:** For prevention of algae on turfgrasses, apply ROLLING 72 SC at the rate of 5.5 – 9.75 pints/acre (2.0 – 3.6 fl. oz/1000 sq ft) (4.1 – 7.3 lbs a.i./acre) on a 7 – 14 day schedule. Under severe algae conditions use the 9.75 pints/acre (3.6 fl oz/1000 sq ft) rate and apply on a 7-day interval. When algae is well established, every attempt should be made to dry out the afflicted area. Once dry, spiking or verticutting should be done to enhance turfgrass recover in conjunction with a ROLLING 72 SC application at the rate of 11 to 15.1 pints/acre (4.0 – 5.5 fl oz/1000 sq ft). **Turf Group B:** A second application of ROLLING 72 SC at the 15.1 pints/acre (5.5 fl oz/1000 sq ft) rate may be made 14 days after the first application. **Turf Groups A & B:** Following applications of the 15.1 pints/acre (5.5 fl oz/1000 aw ft) rate, several applications of ROLLING 72 SC at a rate of 5.5 – 9.75 pints/acre (2.0 – 3.6 fl oz/1000 sq ft) (4.1 – 7.3 lbs a.i./acre) on a 7 to 14 day interval may be necessary for turfgrass recovery. Only a preventive spray program with ROLLING 72 SC will prevent a recurrence of the algae when environmental conditions are favorable.
- Turf Groups A & B:** In areas where pink snow mold (*Geriachia* or *Fusarium* patch) is likely to occur, apply ROLLING 72 SC at 15.1 pints/acre (5.5 fl oz/1000 sq ft) (11.3 lbs a.i./acre) in combination with products containing iprodione at 88 ozs a.i./acre (2 oz a.i./1000 sq ft) of turf area. Read and observe all label directions for products containing these active ingredients. For control of *Fusarium* patch only in areas

38
43

where snow cover is intermittent or lacking during the winter, apply 15.1 pints/acre (5.5 fl oz/1000 sq ft) of ROLLING 72 SC (11.3 lbs a.i./acre). Make application in late autumn. **Turf Group B:** Apply a second application of 15.1 pints/acre (5.5 fl oz/1000 sq ft) of ROLLING 72 SC 21 to 28 days after the first application unless conditions favorable for Fusarium patch no longer prevail.

4. **Turf Group A & B:** For Gray snow mold caused by *Typhula* spp., apply in sufficient water to obtain adequate coverage (2 to 10 gallons per 1000 sq ft). Apply one application 15.1 pints/acre (5.5 fl oz/1000 sq ft) of ROLLING 72 SC (11.3 lbs a.i./acre). Application must be made before snow cover in autumn. **Turf Group B:** If snow cover is intermittent or lacking during the winter, a second application of ROLLING 72 SC at 15.1 pints/acre (5.5 fl oz/1000 sq ft) may be applied one month after the first application.

5. Low rate is not effective on intensively mowed turfgrasses such as golf course and greens.

DIRECTIONS FOR APPLICATION - ORNAMENTAL PLANTS

Apply ROLLING 72 SC at a rate of 1 3/8 pints (1.0 lb a.i.) per 100 gallons of water unless other directions are given in the tables below. DO NOT apply more than 48.5 pints ROLLING 72 SC (36.4 lbs a.i./acre) per growing season to field grown ornamentals. Apply in a spray to the point of drip, when conditions are favorable for disease development. Repeat applications at 7 to 14 day intervals until conditions are no longer favorable. During periods when conditions favor severe disease incidence, generally cloudy or wet weather, apply ROLLING 72 SC at 7 day intervals. The minimum re-treatment interval is 7 days. ROLLING 72 SC should be applied to plants when both foliage and flowers are dry, or nearly dry.

DO NOT combine ROLLING 72 SC in the spray tank with pesticides, surfactants or fertilizers, unless prior use has shown the combination to be physically compatible, effective and noninjurious under your conditions of use.

ROLLING 72 SC may be used in greenhouses. DO NOT use mistblowers or high pressure spray equipment when making applications of ROLLING 72 SC in greenhouses.

ROLLING 72 SC is recommended for control of fungal diseases referred to by numbers in parentheses following each ornamental. Ornamentals listed on this label have been tested and found to tolerate applications of ROLLING 72 SC at the recommended rates. The user should test for possible phytotoxic responses, using recommended rates on ornamental plants on a small area prior to commercial use. Applications made during bloom may damage flowers and/or fruits.

Treated plants and fruits from treated plants MUST NOT BE EATEN.

ORNAMENTALS RECOMMENDED FOR TREATMENT WITH ROLLING 72 SC:

Broadleaf Shrubs and Trees		
Andromeda (<i>Pieris</i>) ⁴	Flowering Almond ^{1,2}	Oregon-Grape (<i>Mahonia</i>) ⁶
Ash (<i>Fraxinus</i>) ¹	Flowering Cherry ^{1,2}	Photinia ¹
Aspen ¹	Flowering Peach ^{1,2}	Poplar ¹
Azalea ^{1,2,4}	Flowering Plum ^{1,2}	Privet (<i>Ligustrum</i>) ¹
Buckeye, Horsechestnut ¹	Flowering Quince ^{1,2}	Rhododendron ^{1,2,4}
Cherry-Laurel ¹	Hawthorn ^{1,6}	Sand Cherry ^{1,2}
Crabapple ^{1,6,8}	Holly ¹	Sequoia ¹
Dogwood ¹	Lilac ⁵	Spiraea ¹
Eucalyptus ³	Magnolia ¹	Sycamore, Planetree ¹
Euonymus ¹	Mountain Laurel ¹	Viburnum ⁵
Firethorn (<i>Pyracantha</i>) ¹	Oak (red group only) ^{1,7}	Walnut (<i>Juglans</i>) ¹

Flowering Plants ^a and Bulbs		
Arabian Violet ² Begonia ¹ Camellia ¹ Carnation ^{1,2} Chrysanthemum ^{1,2} Crocus ¹ Daffodil ¹ Daisy ¹ Geranium ^{1,6}	Gladiolus ^{1,2} Hollyhock ⁶ Hydrangea (foliage only) ^{1,6} Iris ^{1,2} Lily ¹ Marigold ¹ Narcissus ¹ Pansy ¹ Petunia ^{1,4}	Phlox ¹ Poinsetta ^{b 1} Rose ^{c 1} Statice ¹ Tulip ¹ Zinnia ^{1,5}

- ^a Avoid applications during bloom period on plants where flower injury is unacceptable.
^b Discontinue applications prior to bract formation; phytotoxicity is possible on the bracts.
^c Use 1 pint of ROLLING 72 SC per 100 gallons of water.

Foliage Plants		
Aglaonema ¹ Areca palm ¹ Artemisia ¹ Boston fern ¹ Dumbcane (<i>Dieffenbachia</i>) ¹ Dracaena ¹ Fatsia (<i>Aralia</i>) ¹	Ficus ¹ Florida Ruffle Fern ¹ Leatherleaf Fern ¹ Lipstick plant ¹ Ming aralia ¹ Oyster plant (<i>Rhoeo</i>) ¹ Pachysandra ^{d 1}	Parlor palm (<i>Chamaedorea</i>) ¹ Peperomia ¹ Philodendron ^{1,4} Prayer plant (<i>Maranta</i>) ¹ Syngonium ¹ Zebra plant (<i>Aphelandra</i>) ¹

^d Use 2 3/4 pints of ROLLING 72 SC per 100 gallons of water.

Note: DO NOT apply ROLLING 72 SC to either green or variegated Pittosporum or to Schefflera, as multiple applications have been demonstrated to cause phytotoxic responses.

DISEASES CONTROLLED WITH ROLLING 72 SC:

1. Leafspots & Foliar Blights:	
Actinopelte leafspot Alternaria leafspot or leaf blight Anthracnose leaf blotch, spot Anthracnose blight (<i>Discula</i>) Ascochyta blight Bipolaris leafspot (<i>Helminthosporium</i>) Black spot on roses Botrytis leafspot, leaf blight Cephalosporium leafspot Cercospora leafspot Cercosporidium leafspot Corynespora leafspot Coryneum blight (shothole) Curvularia leafspot Cylindrosporium leafspot Dactylaria leafspot Didymellina leafspot	Drechslera leafspot Fabraea leafspot (<i>Entomosporium</i>) Gloeosporium black leafspot Ink spot (<i>Drechslera</i>) Marssonina leafspot Monilinia blossom blight, twig blight Mycosphaerella ray blight Myrothecium leafspot, brown rot Nematostoma leaf blight Phyllosticta leafspot Ramularia leafspot Rhizoctonia web blight Septoria leafspot Sphaeropsis leafspot Stagonospora leaf scorch Tan leaf spot (<i>Curvularia</i>) Volutella leaf blight

42
43

2. Flower Spots and Blights	
Botrytis flower spot, flower blight Curvularia flower spot Monilinia blossom blight	Ovulinia flower blight Rhizopus blossom blight Sclerotinia flower blight

3. Cylindrocadium stem canker

4. Phytophthora leaf blight, dieback

5. Powdery mildews	
<i>Erysiphe cichoracearum</i>	<i>Microsphaera</i> spp.

6. Rusts	
<i>Gymnosporangium</i> spp. <i>Pucciniastrum hydrangeae</i>	<i>Puccinia</i> spp.

7. Taphrina blister

8. Scab (<i>Venturia inaequalis</i>)

DIRECTIONS FOR APPLICATION - TREE AND ORCHARD CROPS

Apply ROLLING 72 SC in sufficient water and with proper calibration to obtain uniform coverage of tree canopy. Application with ground equipment is preferable to aerial application because ground applications generally give better coverage of the tree canopy. If application with ground equipment is not feasible, ROLLING 72 SC may be applied with aircraft using at least 20 gallons of spray per acre. When concentrate sprays are used or when treating non-bearing or immature trees, the lower rate of ROLLING 72 SC listed may be used. DO NOT allow livestock to graze in treated areas. The following spray volumes are recommended as gallons of spray per acre:

Crop	Spray Volume (Gallons per Acre)	
	Concentrate	Dilute
Peach Nectarine Apricot Tart Cherry Plum Prune	20	300
Sweet Cherry	20	400
Conifers		
Forest stands	10 – 20 (aircraft)	Not used
Christmas trees	10 – 50 (aircraft or ground equipment)	100
Nursery beds	5 – 10 (ground equipment only)	100

41
43

Crop	Diseases	ROLLING 72 SC Use Rate		Application Instructions
		Pints/Acre (lbs a.i./acre)	100 Gallons* (lbs a.i./100 gallons)	
Peach Nectarine Apricot Cherry Plum Prune	Leaf curl Coryneum blight (shothole)	3 1/8 - 4 1/8 (2.3 - 3.1)	1 - 1 3/8 (0.75 - 1.0)	For best control of both diseases apply at leaf fall in late autumn, using sufficient water and proper sprayer calibration to obtain uniform coverage. When conditions favor high disease levels use the high rate of application and apply once or twice more in mid to late winter before budswell. If the leaf fall application is not practical, application of ROLLING 72 SC for control of leaf curl may be made at any time prior to budswell the following spring. Where Coryneum blight (shothole) occurs, also apply at budbreak to protect newly emerging leaves and at shuck split to prevent fruit infections.
	Lacy (russet) scab (Plum / Prune)	3 1/8 - 4 1/8 (2.3 - 3.1)	1 - 1 3/8 (0.75 - 1.0)	Make one application at popcorn (pink, red or early white bud) and a second application at full bloom. If weather conditions favor disease development, make an additional application at petal fall.
	Cherry leaf spot Peach, Nectarine, Apricot scab Black knot (Cherry, Plum)	3 1/8 - 4 1/8 (2.3 - 3.1)	1 - 1 3/8 (0.75 - 1.0)	In addition to the bloom application listed above, make one application at shuck split. DO NOT apply ROLLING 72 SC after shuck split and before harvest. If additional disease control is needed before harvest, use another registered fungicide. For control of cherry leaf spot after harvest, make one application to foliage within 7 days after fruit is removed. In orchards with a history of high leaf spot incidence, make a second application 10 - 14 days later.
Restrictions and Limitations:				
<ul style="list-style-type: none"> • DO NOT apply more than 20.5 pints ROLLING 72 SC (15.4 lbs a.i.) per acre during each growing season. • The minimum re-treatment interval is 10 days. 				
Conifers (Pine, Spruce)	Swiss needlecast	2 3/4 - 5 1/2 (2.1 - 4.125)	2 3/4 - 5 1/2 (2.1 - 4.125)	Single application technique: In Christmas tree plantations or forest stands make one application in the spring when new shoot growth is 1/2 to 2 inches in length.
	Scleroderris canker (Pines) Swiss needlecast	1 1/2 - 2 3/4 (1.125 - 2.1)	1 1/2 - 2 3/4 (1.125 - 2.1)	Make the first application in spring when new shoot growth is 1/2 to 2 inches in length. Make additional applications at 3 to 4 week intervals until conditions no longer favor disease development. For use in nursery beds, apply the highest rate specified on a 3-week schedule.
	Sirococcus tip blight	2 - 3 1/2 (1.5 - 2.6)	2 - 3 1/2 (1.5 - 2.6)	

42
43

Crop	Diseases	ROLLING 72 SC Use Rate		Application Instructions
		Pints/Acre (lbs a.i./acre)	100 Gallons* (lbs a.i./100 gallons)	
	Rhizosphaera needlecast (Spruces) Scirrhia brown spot (Pines)	5 ½ (4.125)	5 ½ (4.125)	
	Cyclaneusma and Lophodermium needlecasts (Pines)	2 ¾ to 5 ½ (2.1 - 4.125)	2 ¾ - 5 ½ (2.1 - 4.125)	Apply in early spring prior to budbreak. Repeat applications at approximately 6 - 8 week intervals, until spore release ceases in late fall. Apply monthly during periods of frequent rainfall, and where Lophodermium infections occur during dormancy (Pacific Northwest). During drought periods, applications may be suspended, then resumed upon next occurrence of needle wetness.
	Rhabdocline needlecast (Douglas-fir)	1 ½ - 2 ¾ (1.125 - 2.1)	1 ½ - 2 ¾ (1.125 - 2.1)	Apply at budbreak and repeat at 3 - 4 week intervals until needles are fully elongated and conditions no longer favor disease development. In plantations of mixed provenance, or when irregular budbreak occurs, apply weekly until all trees have broken bud, then every 3 - 4 weeks as specified above. In nursery beds, use the high rate on a 3-week schedule.
	Botrytis seedling blight Phoma twig blight	1 ½ - 2 ¾ (1.125 - 2.1)	1 ½ - 2 ¾ (1.125 - 2.1)	Begin applications in nursery beds when seedlings are 4 inches tall and when cool, moist conditions favor disease development. Make additional applications at 7 to 14 day intervals as long as disease favorable conditions persist.
	Autoecious needle rust (Weir's cushion) (Spruce)	5 ½ (4.125)	5 ½ (4.125)	Begin applications when 10% of buds have broken and twice thereafter at 7 - 10 day intervals.

Restrictions and Limitations:

- DO NOT apply more than 22 pints ROLLING 72 SC (16.5 lbs a.i.) per acre during each growing season.
- The minimum re-treatment interval is 21 days.
- The minimum re-treatment interval in nursery beds is 7 days.

*Volumetric rates to be used only with full dilute spray volume specified on this label for tree and orchard crops.

43
43

Warranty and Disclaimer Statement

The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Such risks may arise from weather conditions, soil factors, off-target movement, unconventional farming techniques, the presence of other materials, the manner of use or application, or other unknown factors, all of which are beyond the control of Arysta LifeScience North America, LLC ("Arysta"), and can cause crop injury, injury to non-target crops or plants, ineffectiveness of the product, or other unintended consequences. All such risks shall be assumed by the user or buyer. Arysta warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks described above, when used in accordance with the Directions for Use under normal conditions. This warranty does not extend to the use of this product contrary to label instructions or under conditions not reasonably foreseeable to Arysta, and is subject to the inherent risks described above.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, ARYSTA DISCLAIMS ALL OTHER WARRANTIES, EXPRESSED OR IMPLIED, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, ARYSTA, MANUFACTURER, AND SELLER DISCLAIM AND SHALL NOT BE LIABLE FOR ANY SPECIAL, INCIDENTAL, INDIRECT, OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE, HANDLING, APPLICATION, STORAGE, OR DISPOSAL OF THIS PRODUCT OR FOR DAMAGES IN THE NATURE OF PENALTIES, AND THE USER AND BUYER WAIVE ANY RIGHT THAT THEY MAY HAVE TO SUCH DAMAGES. NO AGENT, REPRESENTATIVE OR EMPLOYEE OF ARYSTA IS AUTHORIZED TO MAKE ANY WARRANTY, GUARANTEE OR REPRESENTATION BEYOND THOSE CONTAINED HEREIN OR TO MODIFY THE WARRANTIES CONTAINED HEREIN.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE TOTAL LIABILITY OF ARYSTA, MANUFACTURER, AND SELLER, SHALL BE LIMITED TO THE PURCHASE PRICE PAID, OR AT ARYSTA'S ELECTION, THE REPLACEMENT OF THE PRODUCT.

Dipel is a registered trademark of Valent BioSciences Corporation
Latron B-1956 and latron AG-98 are registered trademarks of Dow AgroSciences LLC
Arysta LifeScience North America, LLC

ROLLING 72 SC (Master Label) AD061307, Notification 02/27/09, Notification 01/20/11