66330-367

05-19-20101



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, D.C. 20460

> OFFICE OF PREVENTION, PESTICIDES AND TOXIC SUBSTANCES

Bill Washburn Arysta LifeScience North America, LLC 15401 Weston Parkway, Suite 150 Cary, NC 27513

MAY 19 2010

Subject: Label Notification(s) Per Agency Letter Dated June 4, 2009; Environmental Hazard and General Labeling for Pyrethroid Non-Agricultural Outdoor Products Notification

Dear Mr. Washburn:

The Agency is in receipt of your applications for pesticide notifications, dated May 7, 2010, per Agency letter dated June 4, 2009.

EPA Registration 66330-365 EPA Registration 66330-366 EPA Registration 66330-367 EPA Registration 66330-368 Aloft GC SC Insecticide Aloft LC SC Insecticide Aloft GC G Insecticide Aloft LC G Insecticide

The Registration Division (RD) has conducted a review of these requests for applicability under Agency letter dated June 4, 2009, and find that the label changes requested fall within the scope of the letter. The labels have been date-stamped "*Notification*" and will be placed in our records.

If you have any questions, please contact me at (703) 306-0415 or <u>davis.kable@epa.gov</u>.

Sincerely,

Kable Bo Davis Entomologist Insecticide-Rodenticide Branch Registration Division (7505P)

| | | | | | ر ا) 060, Approval expires 05-31-98 |
|------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------|------------------------------------------------------------------|
| EPA | United St Environmental Pro Washington, D | tection Agency | ☐ Registra ☐ Amendn ⊠ Other: | ition | OPP Identifier Number |
| | Appl | lication for Pest | ticide - Sectior | n l | |
| 6330-367 | | | oduct Manager agle | | 3. Proposed Classification |
| 4. Company/Product (Name) Aloft GC G Insecticide | | PM# | | | None Restricted |
| 5. Name and Address of Applic Arysta LifeScience Nortl 15401 Weston Parkway, Cary, NC 27513 | n America, LLC Suite 150 | 6. Expect (b)(I), my to: EPA Reg | lited Review . In / product is simila g. No Name | MAY 1.9.2 | |
| Check if this is a n | ew address | Sectior | | · ···· · · · · · · · · · · · · · · · · | <u></u> |
| violation of 18 USC Sec | onal page(s) if neces ange per June 4, 200 in the June 4, 2009 li e to the labeling or th 1001 to willfully make quirements of 40 CF | 09, letter from En etter and requirem le Confidential Sta any false stateme R part 156, this pr | vironmental Pro lents of EPA's re tement of Formu ent to EPA. I furt oduct may be in v |) p tection Agenc gulations at 40 (la for this produ her understand | CFR part 156. No other ct. I understand that it is a |
| | | Section |) - 111 | | |
| 1. Material This Product Will E Child-Resistant Packaging Yes* No | Yes* | | Water Soluble Packaging Yes No No. per | | 2. Type of Container Metal Plastic Glass |
| *Certification must be submitted | Unit Packaging v | wgt. container | Package wgt. | container | Paper Other (Specify) |
| 3. Location of Net Contents In | formation 4. S Container | Size(s) Retail Containe | r | On Label | Label Directions |
| 6. Manner in Which Label is A | | Lithograph Paper glued Stenciled | Other | | |
| | | Section | - IV | | |
| 1. Contact Point (Complete iter | ms directly below for ident | ification of individual to | be contacted, if nec | essary, to process | this application) |
| Name Bill Washburn | | Title Regulatory | Manager | | elephone No. ((nclude Area |
| I certify that the statements I h | Certi | ification | | I | ode) 901-432-5118 |
| acknowledge that any knowing under applicable law. | ave made on this form and ly false or misleading state | | | | 901-432-5118 6. Date Application Received (Stamped) |
| acknowledge that any knowing | ly false or misleading state | | ble by fine or impriso | onment or both o | 901-432-5118 6. Date Aβplication cos Received (Stamped) |



Arysta LifeScience

May 7, 2010

Ms. Venus Eagle, PM 1 Document Processing Desk (NOTIF -PYRETHROID) Office of Pesticide Programs – 7504P U.S. Environmental Protection Agency Room S-4900, One Potomac Yard 2777 South Crystal Drive Arlington, VA 22202-4501

Subject: Aloft GC G Insecticide EPA Reg. No. 66330-367

Notification of label change per June 4, 2009, letter from Environmental Protection Agency.

 \mathcal{A}

Dear Ms. Eagle:


Notification of label change per June 4, 2009, letter from Environmental Protection Agency. This notification is consistent with guidance in the June 4, 2009 letter and requirements of EPA's regulations at 40 CFR part 156. No other changes have been made to the labeling or the Confidential Statement of Formula for this product. I understand that it is a violation of 18 USC Sec 1001 to willfully make any false statement to EPA. I further understand that if the amendment is not consistent with the requirements of 40 CFR part 156, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.


Enclosed are two copies of the revised label. One clean copy and one copy with changes highlighted. Please acknowledge acceptance of this notification. Should you have any questions or comments, please do not hesitate to contact me at 901-432-5118 or by e-mail at <u>bill.washburn@arystalifescience.com.com</u>

Sincerely,

ie (vooler

Bill Washburn Regulatory Manager





0000 0000

RESTRICTED USE PESTICIDE

Toxic to Fish and Aquatic Organisms

For retail sale to and use only by certified applicators, or persons under their direct supervision and only for those uses covered by certified applicator's certification.



| GROUP | 4A | INSECTICIDE |
|-------|----|-------------|
| GROUP | 3 | INSECTICIDE |

For contact and systemic insect pest control in turf and landscape ornamentals on golf courses, lawns and landscape ornamentals around residential, institutional, public, commercial and industrial buildings, parks, recreational areas, athletic fields, and sod farms.

DO NOT USE THIS PRODUCT ON GOLF COURSES AND SOD FARMS IN NASSAU COUNTY OR SUFFOLK COUNTY, NEW YORK.

| INGREDIENTS | By wt. |
|---------------------|--------|
| Active Ingredients: | |
| Clothianidin* | |
| Bifenthrin** | |
| Other Ingredients | |
| Total | |
| | |

*(E)-1-(2-chloro-1,3-thiazol-5-ylmethyl)-3-methyl-2-nitroguanidine.

** 2-methylbiphenyl-3-ylmethyl (1RS,3RS)-3-[(Z)-2-chloro-3,3,3-trifluoroprop-1-enyl]-2,2-

dimethylcyclopropanecarboxylate (cis isomers 97% minimum, trans isomers 3% maximum)

KEEP OUT OF REACH OF CHILDREN CAUTION

See inside booklet for additional precautionary statements

EPA Registration No. 66330-367 EPA Est. No.

NOTIFICATION

MAY 1 9 2010

Produced for: Arysta LifeScience North America, LLC 15401 Weston Parkway, Suite 150 Cary, NC 27513

For Product Use Information, call 1-866-761-9397

| | FIRST AID |
|------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| If swallowed | Call poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person. |
| If inhaled | Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice. |
| If in eyes | Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. |
| If on skin or clothing | Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice. |
| NOTE TO PHYSICIAN. | This pesticide contains a pyrethroid. If significant quantities have been |

NOTE TO PHYSICIAN: This pesticide contains a pyrethroid. If significant quantities have been swallowed, promptly eliminate the ingested quantities from the stomach and intestines. Treatment is symptomatic and supportive. Alcohol, milk, cream, and other digestible fats and oils should NOT be administered as they may increase absorption.

HOTLINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. **FOR 24-HOUR EMERGENCY MEDICAL ASSISTANCE:** CALL PROSAR at 1-866-303-6952.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wear protective eyewear. Wear: long-sleeved shirt and long pants, socks, shoes, and chemical-resistant gloves (such as Natural Rubber, Selection Category A).

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some of the materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart. Applicators and other handlers must wear:

- long-sleeved shirt and long pants
- shoes plus socks
- chemical-resistant gloves
- protective eyewear

Follow the manufacturer's instructions for cleaning/maintaining PPE. If no instructions for washables are available, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

- Users should:
 Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

T

This product is extremely toxic to fish and aquatic invertebrates. <u>To protect the environment, do not</u> <u>allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters.</u> <u>Applying this product in calm weather when rain is not predicted for the next 24 hours will help to</u> <u>ensure that wind or rain does not blow or wash pesticide off the treatment area</u>. <u>Sweeping any</u> <u>product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden</u> <u>will help to prevent run off to water bodies or drainage systems</u>. Drift and runoff may be hazardous to aquatic organisms in neighboring areas. Do not apply where runoff is likely to occur. Do not apply <u>directly to water, or to areas where surface water is present or to intertidal areas below the mean high</u> <u>water mark</u>. Do not contaminate water when cleaning equipment or disposing of equipment washwaters or rinsate.

This product is toxic to bees exposed to treatment and for more than 5 days following treatment. Do not apply this product to blooming, pollen-shedding or nectar-producing parts of plants if bees may forage on the plants during this time period, unless the application is made in response to a public health emergency declared by appropriate state or federal authorities.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Carefully read and understand the Directions for Use and restrictions before applying this product.

Do not apply this product in any way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

RESTRICTIONS

Apply using ground equipment only; do not apply by air.

Do not apply within a 25 foot radius of the following: Lakes, reservoirs, rivers, permanent streams, marshes or natural ponds, estuaries, and commercial fish farm ponds.

<u>Application is prohibited directly into sewers or drains, or to any area like a gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur. Do not allow the product to enter any drain during or after applications.</u>

Do not apply on the following: Commercial nurseries, commercial greenhouses, and grass grown for seed.

Apply this product directly to the lawn or garden area. Water treated area as directed on this label. Do not water to the point of run-off.

Do not make applications during rain.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions on this label about personal protective equipment (PPE), notification to workers and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry into treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

Coveralls

1

- Shoes plus socks
- Chemical-resistant gloves

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in the box apply to use of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries and commercial greenhouses.

Keep children and pets off treated area until the dust has settled.

INFORMATION

ALOFT[™] GC G is a broad spectrum, long lasting contact and systemic insecticide for control of listed insects infesting turfgrass, sod and ornamentals in landscapes and surrounding areas. ALOFT GC G can be used around residences, business complexes, industrial complexes, airports, playgrounds, school yards, parks and recreation areas, athletic fields, golf courses, cemeteries and arboretums.. ALOFT GC G can be used on commercial sod farms.

RESISTANCE MANAGEMENT

The active ingredients in **ALOFT GC G** are clothianidin, a Group 4A insecticide, and bifenthrin a Group 3 insecticide. Insect biotypes with acquired resistance to Group 4A or Group 3 insecticides may eventually dominate the insect population if Group 4A or Group 3 insecticides are used alone or repeatedly in the same site or in successive years as the primary method of control for targeted species. Aloft GC G contains two active ingredients with different modes of action and provides a recognized built in insecticide resistance management strategy to minimize the development of insecticide resistance in insect pests.

To delay insecticide resistance in insects that are at risk of developing resistance to insecticides consider:

- Avoid the consecutive use of insecticides in the same group on the same insect species.
- Use tank-mixtures or pre-mixes of insecticides from a different target site of action.

Filename: Aloft GC G (367-112408)(adjacent to turf 061809) CLEAN RVSD.doc ----------066330-00367.20090619.adjacent to turf clean-rvsd.pdf

- Base insecticide use on a comprehensive IPM program.
- Monitor treated insect populations for loss of field efficacy.
- Contact your State Agricultural Experiment Station, State Extension Specialists, other turf experts and/or Arysta LifeScience North America Corporation for insecticide resistance management and/or IPM recommendations for the specific site and resistant pest problems.
- For further information or to report suspected resistance, you may contact Arysta LifeScience North America Corporation at 1-866-761-9397.

APPLICATION EQUIPMENT

Apply **ALOFT GC G** uniformly over the area being treated with standard granular application equipment. Use either drop-type or rotary-type spreaders to apply **ALOFT GC G**.

Avoid the use of spreaders that apply the material in narrow, concentrated bands. Calibrate application equipment prior to use according to the manufacturer's directions. Check frequently to be sure equipment is working properly and distributing granules uniformly and accurately. In order to insure continuing delivery of the correct amount of **ALOFT GC G** further adjustments to the settings should be made periodically during application.

APPLICATION INSTRUCTIONS – TURFGRASS AND PERIMETERS AROUND LAWNS AND LANDSCAPED AREAS

ALOFT GC G can be used for broad spectrum insect control in turfgrass and perimeters. **ALOFT GC G** can be used around residences, business complexes, industrial complexes, airports, playgrounds, school yards, parks and recreation areas, athletic fields, golf courses, cemeteries and arboretums.

ALOFT GC G has sufficient contact and residual activity to provide high levels of control. The need for an application can be based on historical monitoring of the presence of or anticipated occurrence of individual or multiple insect pest species, the site, previous records or experiences, current season adult trapping or other methods. Optimum control of white grubs will be achieved when applications are made just prior to egg laying or to early instar larvae of target pests, followed by sufficient irrigation or rainfall to move the active ingredient into the zone where the insect pests are active. Applications made in advance of egg laying or to areas being damaged by existing white grubs or scavenging animals will be controlled with an application of Aloft LC G anytime white grubs are active. Multiple applications can be made each growing season but do not use more than 160 lbs of product (0.4 lb of clothianidin active ingredient, 0.2 lb bifenthrin) per acre per year as a broadcast application. Note: You may also use additional products containing bifenthrin as long as the maximum amount of bifenthrin does not exceed 0.4 lbs active ingredient per acre per year for use in lawns and perimeters.

ALOFT GC G can be applied to turf at 80 - 160 lbs per acre per year (1.8 - 3.6 lbs/1000 sq ft). The rate is dependent on the target pest(s), their stage of development and the desired level of control. Optimum control of white grubs (larvae) will be achieved when applications are made just prior to or just after peak egg laying. Higher rates should be used when long residual control is desired for control of multiple insect pests or late season curative (rescue) applications are made to turf already showing grub damage or damage due to scavenging of animals. Multiple applications can be made but do not exceed the maximum amount per year (160 lbs/acre or 3.6 lbs/1,000 sq ft). Consult your local State Agricultural Experiment Station, State Extension Turf Specialists or other turf experts for specific information concerning the timing of applications.

For early season control of insect pests (such as annual bluegrass weevil) and for full season control of white grubs, make the first application of **ALOFT GC G** at peak adult activity or when other insect pests reach a damaging population in the spring at a rate of 100 - 160 lbs per acre (2.3 - 3.6 lbs/1000 sq ft). For control of white grubs at standard application timing, apply 80 - 160 lbs per acre (1.8 - 3.6 lbs/1000 sq ft) just prior peak adult activity or to early instar larvae.

Filename: Aloft GC G (367-112408)(adjacent to turf 061809) CLEAN RVSD.doe 066330-00367.20090619.adjacent to turf clean rvsd.pdf

For surface insects, best control will be obtained if Aloft GC G is not watered in until at least 24 hours after the application. If the target insects are in the thatch, lightly water in after application with enough water to wet the thatch zone. For control of armyworms, cutworms and other surface feeders, apply 80 - 160 lbs per acre (1.8 - 3.6 lbs/1000 sq ft) at the first sign of insects or insect damage. Repeat the application as needed to control additional flushes of insects. Up to a total of 160 lbs per acre per year (3.6 lbs/1,000 sq ft) can be applied.

Ί

For control of existing grub populations, best results will be obtained if you irrigate after application with enough water to move the insecticide into the zone where the grubs are feeding.

Avoid mowing turfgrass until after irrigation or rainfall has occurred so that uniformity of application will not be affected.

| TURFGRASS AND PERIMETERS AROUND LAWNS | | | | |
|----------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--|--|--|
| APPLICATION RATE | INSECT PEST | | | |
| 1.8 - 3.6 lbs per 1,000 sq. ft. (80-160 lbs per acre) | Annual Bluegrass Weevil (Adults)/Hyperodes ¹ Annual Bluegrass Weevil (Larvae)/Hyperodes ¹ Ants ² , Nuisance (Excluding fire, harvester, pharaoh, and carpenter ants) Armyworms Billbugs ³ Black Turfgrass Ataenius ⁴ Chiggers Chinch Bugs Cutworms European Crane Fly Grasshoppers Leafhoppers Mole Crickets ⁵ Pillbugs Sod webworms Sowbugs Spittle bug Sugarcane grub White grubs ⁶ (Asiatic Garden Beetle, European Chafer, Green June Beetle, Japanese Beetle, Northern Masked Chafer, <i>Phyllophaga</i> spp. (May or June Beetle), Oriental Beetle, Southern Masked Chafer) | | | |

¹ Annual Bluegrass Weevil: Application can be timed for adult weevils as they move from their over-wintering sites into turf. This movement usually begins before or when Forsythia is in full bloom and continues to full bloom of dogwoods. Larvae from adult weevils that escape the application will be controlled with the application directed at adult weevils. Applications made after adult weevil activity will control larvae. Consult your State Cooperative Extension Service for more specific information regarding timing of application. Full season control of white grubs will be obtained with an application of 100 – 160 lb/Acre targeted towards Annual Bluegrass Weevil/Hyperodes.

² Applications of Aloft GC G in the late fall or early spring will reduce populations of nuisance ants, excluding fire, harvester, pharaoh, and carpenter ants. Make an application in the fall before ants stop foraging or after they become active in the spring.

TURFGRASS AND PERIMETERS AROUND LAWNS

- ³ Billbugs: Application can be made when adult Billbugs are first observed, usually in April and May in temperate regions. A spring application will also control Billbug larvae. Consult your State Cooperative Extension Service for more specific information regarding timing of application. White grubs will be controlled with an application targeted towards Billbugs.
- ⁴ Black Turfgrass Ataenius: Application can be timed to coincide with the first generation of Black Turfgrass Ataenius that usually appears during May. White grubs will be controlled with an application targeted towards Black Turfgrass Ataenius.
- ⁵ **Mole Crickets:** Application should be made during peak adult flight and egg lay. If the soil is dry, irrigate before application to stimulate insect activity and water after application to release the insecticide from the granule carrier.
- ⁶ Aloft GC G will control white grubs anytime applications are made when they are active from early spring to late fall. The residual activity of Aloft GC G is long enough to provide season long control of white grubs when applied for control of early season insects such as Annual Bluegrass Weevil, Billbugs, Nuisance Ants, overwintering White Grubs or other early season insects.

RESTRICTIONS

ł

- Do not apply more than a total of 160 lbs of **ALOFT GC G** (0.4 lb of clothianidin active ingredient, 0.2 lb bifenthrin) as a broadcast application per acre per year.
- If the maximum season limit of ALOFT GC G has been applied and pest populations require additional treatments, use another pesticide labeled for control of the targeted insects that does not contain clothianidin. You may use additional products containing bifenthrin as long as the maximum amount of bifenthrin does not exceed 0.4 lbs active ingredient per acre per year for use in lawns and perimeters.

| | Granular Spreader Settings | | | | | |
|----------------------|----------------------------|---------------------------|--------------------|--|--|--|
| Spreader | 2 lb/1000 sq ft | 3 lb/ 1000 sq ft | 4 lb/ 1000 sq ft | | | |
| Cyclone | 3 | 4 | 4 1/4 | | | |
| Lely | 3 | 4 | 5 | | | |
| Lesco | 10 | 15 | 16 | | | |
| Scotts R8A/SR-1 | I | K | L | | | |
| Spyker | 3 | 3 1/2 | 4 1/4 | | | |
| Vicon | 20 | 22 | 23 | | | |
| Note: Spreader setti | ings are approximate | starting points for calib | prating equipment. | | | |

When treating perimeters of structures, make the application to the ground from 5-10 feet adjacent to the structure.

| | Band Width | Lbs. of ALOFT GC G/ Linear Distance (Feet) Around Structure | | | | | | | |
|-------|---------------|-------------------------------------------------------------|-----|-----|-----|------|------|------|-------|
| sq ft | in Feet | 100 | 150 | 200 | 250 | 300 | 400 | 500 | 1,000 |
| 1.8 | 5 | 1.0 | 1.5 | 2.0 | 2.5 | 3.0 | 4.0 | 5.0 | 10.0 |
| 1.8 | 10 | 2.0 | 3.0 | 4.0 | 5.0 | 6.0 | 8.0 | 10.0 | 20.0 |
| 2.8 | 5 | 1.4 | 2.0 | 2.8 | 3.5 | 4.2 | 5.6 | 7.0 | 14.0 |
| 2.8 | 10 | 3.0 | 4.5 | 6.0 | 7.5 | 9.0 | 12.0 | 15.0 | 30.0 |
| 3.6 | 5 | 2.0 | 3.0 | 4.0 | 5.0 | 6.0 | 8.0 | 10.0 | 20.0 |
| 3.6 | 10 | 3.6 | 5.4 | 7.2 | 9.0 | 10.8 | 14.4 | 18.0 | 36.0 |

area of a circle whose radius is the bandwidth (this accounts for the area treated at the four corners of the structure) plus the linear distance treated times the bandwidth.

APPLICATION INSTRUCTIONS – LANDSCAPE ORNAMENTALS

1

ALOFT GC G is a broad spectrum, contact and long lasting systemic insect control product for a wide spectrum of insects infesting landscape ornamentals adjacent to turf areas. ALOFT GC G can be used around residences, business complexes, industrial complexes, airports, playgrounds, school yards, parks and recreation areas, athletic fields, golf courses, cemeteries and arboretums.

For control of listed surface insects and insects that crawl along the surface of the soil or mulch under ornamentals, best results will be obtained if Aloft GC G is watered in lightly after application to move the insecticide into the areas where insects are active. Water in with enough irrigation or rainfall to move the active ingredient into the zone where the target insects are active. Soil surface applications will provide control of foliage feeding insects; however, optimum control will not be achieved until the active ingredient is taken up by the plant roots and translocated to the area where insects are feeding.

| LANDSCAPE ORNAMENTALS | | | | | |
|--------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--|--|
| SOIL SURFACE or MULCH APPLICATION | SOIL INSECT PE | APPLICATION INSTRUCTIONS | | | |
| 1.8 – 3.6 lb per 1,000 sq ft (80 – 160 lb per acre) | Ants, Nuisance (Excluding fire, harvester, pharaoh, and carpenter ants) Beet Armyworm Beetles Black Vine Weevils (Adults) Centipedes Crickets Cutworms Diaprepes (Adults) Earwigs Fall Armyworms Grasshoppers Millipedes | Pillbugs White grubs (Asiatic Garden Beetle, European Chafer, Green June Beetle, Japanese Beetle, Northern Masked Chafer, <i>Phyllophaga</i> spp. (May or June Beetle), Oriental Beetle, Southern Masked Chafer) | Aloft LC G must be watered in or receive rainfall after application to be activated. Control of leaf feeding insects will be delayed until the insecticide is taken up by the roots and moved into the areas where insects are feeding, usually 2-4 weeks. For ornamental trees or non-bearing fruit and nut trees and woody shrubs see below. | | |

| LANDSCAPE ORNAMENTALS | | | | | | |
|---------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--|--|--|
| SOIL SURFACE or MULCH APPLICATION | ABOVE GROU | APPLICATION INSTRUCTIONS | | | | |
| 3.0 – 3.6 lb per 1,000 sq ft (132 - 160 lb per acre) | Aphids Azalea lace bug Caterpillars Flea Beetles Florida wax scale Glassy winged sharpshooter Gypsy Moth Caterpillars Japanese Beetles (Adults) | Leafhoppers Leafminers Leafrollers Mealybugs Plant Bugs (incl. <i>Lygus</i> spp) Root Weevils (incl. Black Vine Weevils) Scales, Soft Whiteflies | Apply in enough water to wet the root zone of the plants. Control of leaf feeding insects will be delayed until the insecticide is taken up by the roots and moved into the areas where insects are feeding, usually 2-4 weeks. For ornamental trees or non-bearing fruit and nut trees and woody shrubs see below. | | | |

- Do not apply more than a total of 160 lbs of **ALOFT GC G** (0.4 lb of clothianidin active ingredient, 0.2 lb bifenthrin) as a broadcast application per acre per year.
- If the maximum season limit of ALOFT GC G has been applied and pest populations require additional treatments, use another pesticide labeled for control of the targeted insects that does not contain clothianidin. You may use additional products containing bifenthrin as long as the maximum amount of bifenthrin does not exceed 0.4 lbs active ingredient per acre per year for use in lawns and perimeters.

APPLICATION INSTRUCTIONS - ORNAMENTAL TREES, NON-BEARING FRUIT AND NUT TREES AND WOODY SHRUBS IN LANDSCAPE AREAS ADJACENT TO TURF AREAS.

Aloft GC G can be used for systemic control of insects with an application on trees in landscapes adjacent to turf areas. Aloft LC G contains clothianidin which is systemic and will be translocated throughout the tree. Treat the area over the root zone of the trees or apply the insecticide around the "root well" of the plant.

Aloft GC G is a systemic product and will be translocated upward into the plant system from root uptake. It must be placed in an area where the roots can absorb the active ingredient. It may take 2-4 weeks for the active ingredient to move from the root zone to the areas of the trees. If insect pests are already present and causing damage, consider making a foliar application and then following up with a granule application. Water in after application with enough water to move the insecticide into the root zone of the trees.

For trees and shrubs adjacent to turf areas, use 0.4-0.8 lb Aloft GC G per inch of cumulative trunk diameter (DBH) of the plant(s). For hedges six feet or less in height use 4.0 lb Aloft GC G per 100 linear feet of hedge. For hedges greater than six feet in height use an additional 1.0 lb per foot of hedge height over six feet. Before application remove any plastic or other barriers that may prevent the insecticide from reaching the root zone of the plants.

For Bronze Birch Borer, Flat Headed Borers (excluding Emerald Ash Borers) and other borers: Make an application around the trunk of the trees when the trees are actively growing. Use 0.2-0.4

Ib Aloft GC G per inch DBH in a 1-2 ft area around trunks of the trees.

RESTRICTIONS

- Do not apply more than a total of 160 lbs of **ALOFT GC G** (0.4 lb of clothianidin active ingredient, 0.2 lb bifenthrin) as a broadcast application per acre per year.
- If the maximum season limit of ALOFT GC G has been applied and pest populations require additional treatments, use another pesticide labeled for control of the targeted insects that does not contain clothianidin. You may use additional products containing bifenthrin as long as the maximum amount of bifenthrin does not exceed 0.4 lbs active ingredient per acre per year for use in lawns and perimeters.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in a cool, dry place away from excessive heat. Do not store near food, feed, fertilizers, seeds, or other pesticides. Keep this product out of reach of children and animals at all times. Store in original container only. Open the product container carefully and after use, seal tightly.

To contain a spill: Cover the spilled product to prevent the material from spreading. If the product container is damaged, place the entire container along with the product in a secure receptacle and clearly label the contents.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL:

Paper and plastic bags: Completely empty bag into application equipment. Then dispose of in a sanitary landfill, or (if allowed by State and local authorities) by burning. If burned, stay out of smoke. FOR 24-HOUR CHEMICAL EMERGENCY (Spill, Leak, Fire or Accident) ASSISTANCE: Call CHEMTREC at 1-800-424-9300.

Warranty and Disclaimer Statement

The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Such risks may arise from weather conditions, soil factors, off-target movement, unconventional farming techniques, the presence of other materials, the manner of use or application, or other unknown factors, all of which are beyond the control of Arysta LifeScience North America Corporation ("Arysta"), and can cause crop injury, injury to non-target crops or plants, ineffectiveness of the product, or other unintended consequences. All such risks shall be assumed by the user or buyer.

Arysta warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks described above, when used in accordance with the Directions for Use under normal conditions. This warranty does not extend to the use of this product contrary to label instructions or under conditions not reasonably foreseeable to Arysta, and is subject to the inherent risks described above.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, ARYSTA DISCLAIMS ALL OTHER WARRANTIES. EXPRESSED IMPLIED. INCLUDING ANY WARRANTY OR OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, ARYSTA, MANUFACTURER, AND SELLER DISCLAIM AND SHALL NOT BE LIABLE FOR ANY SPECIAL, INCIDENTAL, INDIRECT, OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE, HANDLING, APPLICATION, STORAGE, OR DISPOSAL OF THIS PRODUCT OR FOR DAMAGES IN THE NATURE OF PENALTIES, AND THE USER AND BUYER WAIVE ANY RIGHT THAT THEY MAY HAVE TO SUCH DAMAGES. NO AGENT, REPRESENTATIVE OR EMPLOYEE OF ARYSTA IS AUTHORIZED TO MAKE ANY WARRANTY, GUARANTEE OR REPRESENTATION BEYOND THOSE CONTAINED HEREIN OR TO MODIFY THE WARRANTIES CONTAINED HEREIN. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE TOTAL LIABILITY OF ARYSTA, MANUFACTURER, AND SELLER, SHALL BE LIMITED TO THE PURCHASE PRICE PAID, OR AT ARYSTA'S ELECTION, THE **REPLACEMENT OF THE PRODUCT.**

ALOFT is a trademark of Arysta LifeScience North America, LLC.