

PM 1466550-1 p71/2

Bird Shield®

Bird Repellent Concentrate

ACCEPTED
MAR 10 1997**ACTIVE INGREDIENT**

Methyl Anthranilate (MA) methyl 2-aminobenzoate

By Weight

26.4%

INERT INGREDIENTS

73.6%

Total

100.0%

This product contains 2.29 lbs. of Methyl Anthranilate (MA) per gallon.
 This concentrate has been formulated from food grade ingredients that meet or exceed U.S. standards.

BIODEGRADABLE MATERIAL - USE BY [Insert date before shipping product]**WARNING****KEEP OUT OF REACH OF CHILDREN****Hazards to Humans and Domestic Animals****Warning**

Causes substantial but temporary eye injury. Harmful if swallowed. Do not get in eyes or on clothing. Avoid contact with skin. Wear goggles or face shield. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash clothing before reuse.

FIRST AID

IF IN EYES: Hold eyelids open and flush with a steady stream of water for 15 minutes. Get medical attention.

IF SWALLOWED: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting. Do not induce vomiting or give anything by mouth to an unconscious person.

IF ON SKIN: Wash with plenty of soap and water. Get medical attention if irritation persists.

ENVIRONMENTAL HAZARDS

Do not apply directly to fish-bearing bodies of water, to areas where surface water may contaminate streams or lakes or to intertidal areas below the mean high water mark.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Use Restrictions. This product may be used

1. to limit feeding of robins (*Turdus migratorius*), starling (*Sturnus vulgaris*), Cedar waxwings (*Bombicilla cedrorum*), jays, magpies and crows (*Corvidae*), ravens (*Corvus spp.*), finches and sparrows (*Fringilidae*) on ripening cherries, blueberries and grapes.
2. to discourage ducks and geese (Family *Anatidae*) from using non-fishbearing bodies of water (tailing ponds, commercial or industrial water impoundment's) and temporary pools of standing water located at or near airports; and
3. to reduce use of turf areas by Canada geese (*Branta canadensis*).

Preparation and Mixing Directions. Warm, in the sun or a water bath to 75°F if any part of the formulation has solidified. Mix liquid concentrated solution with a blade or paddle until ingredients are

282
evenly distributed before adding to water. The product may be mixed with *Bacillus thuringensis* formulations. Do not mix with other pesticides or fertilizers.

Application Directions -

Blueberries, cherries and grapes. Mix 1 part Bird Shield Repellent Concentrate with 99 parts of water. Agitate well before application.

Apply mixture with a commercial or back-pack sprayer, hand-held hose or pressurized applicators. Begin applications when fruit begins to ripen or birds begin feeding on crop. Thoroughly wet all fruit and foliage until solution runs off surfaces. Apply every 6 to 8 days or when odor can not be detected. Repeat as necessary to maintain repellency. Harvest 6 to 8 days after last treatment or after all odor or taste of the product has dissipated (whichever occurs last).

Do not apply this product to wet surfaces. Repeat treatment if heavy rains occur within 24 hour of application.

Non-fishbearing bodies of water and temporary pools associated with airports. Add 1-2 parts of Bird Shield concentrate per 400 parts of water in the body of water (1-2 gal/53 cubic feet of water). Estimate the volume of standing water by considering the depth, length, width and shape of pool and using appropriate mathematical formulas. Calculate the amount of Bird shield concentrate needed to treat the pool within the prescribed rate range. Add undiluted concentrate directly to entire pool surface until the appropriate amount has been applied. Reapply at 3-4 week interval if necessary to maintain repellency.

Turf. Mix 2-8 gallons of Bird Shield concentrate with sufficient water to make 60 gallons of mix. Apply 60 gallons of mix per acre, making sure that turf is thoroughly saturated. Reapply product at intervals of 7-14 days, or after each mowing, if necessary to maintain repellency. Attempt to prohibit or discourage humans from feeding geese on or near areas from which the birds are to be excluded. Do not apply this product to wet surfaces. Repeat treatment if heavy rains occur within 2 hours of application.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal

STORAGE: Store only in original container apart from pesticides, fertilizers, food or feed that may cause cross-contamination from odor and inaccessible to children, pets and domestic animals. Avoid freezing or placing containers in direct sunlight. If frozen slowly thaw and remix contents. Keep tightly closed when not in use.

REPELLENT DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at a waste disposal facility by exposing to direct sunlight until all odor or the product has dissipated or in accordance to Federal or State Procedures.

CONTAINER DISPOSAL: Triple rinse. Offer for recycling or reconditioning or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Bird Shield Repellent Corporation
P.O. Box 785
Pullman, Washington 99163-0785

509/332-1989 Fax 332-0190 e-mail askham@BSRC.com

EPA EST. No. 66550-WA-1
EPA REG. No. 66550-1
2/18/97

Net one U.S. Gallon (3.78 litres) 128 fl. oz.