Reg # 66665-8	PM-18	1
U.S. ENVIRONMENTAL PROTECTION AGENCY Office of Pesticide Programs Registration Division (H7505C) 401 "M" St., S.W. Washington, D.C. 20460	EPA Reg. Number: 66665-8	Date of Issuance: 06/24/94
NOTICE OF PESTICIDE: <u>x</u> Registration Condition		
(under FIFRA, as amended)	Name of Pesticide Product: Ford's Organicide Plant & Garden Spray	
Name and Address of Registrant (include ZIP Code): Innova Corporation, Suite 403, 3550 Hiway 77478	6, Sugarland	, Texas
Note: Changes in labeling differing in substance from that accepted in be submitted to and accepted by the Registration Division prior to use correspondence on this product always refer to the above EPA registrat	of the label in com	
On the basis of information furnished by the registrant, the above nam registered/reregistered under the Federal Insecticide, Fungicide and R		ъу
Registration is in no way to be construed as an endorsement or recomme In order to protect health and the environment, the Administrator, on cancel the registration of a pesticide in accordance with the Act. The with the registration of a product under this Act is not to be constru- exclusive use of the name or to its use if it has been covered by othe	his motion, may at a me acceptance of any med as giving the reg	ny time suspend or name in connection
This product is conditionally registe FIFRA sec. 3(c)(7)(A) provided that you:	red in accor	dance with
1. Submit and/or cite all data requireregistration of your product under FIFRA Agency requires all registrants of similar data; and submit acceptable responses requires of your product under FIFRA section 4.	sec. 3(c)(5 products to) when the submit such
2. Make the following label changes: Revise the EPA Registration N No. 66665-8".		d, "EPA Reg.
3. Submit two copies of the revised label for the record.	final printe	d
If these conditions are not com registration will be subject to cancellati FIFRA sec. 6(e). Your release for shipmen constitutes acceptance of these conditions	on in accord it of the pro	lance with
A stamped copy of the label is records.	enclosed for	your
BES		BLE COPY
Signature of Approving Official:	Date:	J
		<u>Z</u>

3

)

FORD'S ORGANICIDE PLANT & GARDEN SPRAY Contains Rotenone & Pyrethrins for Dilution with Water

FOR INSECT CONTROL ON ORNAMENTALS - FRUITS - VEGETABLES FOR FLEA CONTROL ON DOGS AND IN AND AROUND HOMES One eight ounce bottle makes up to 48 gallons of finished spray.

This product will kill and control 43 common insect pests that attack ORNAMENTALS, SHRUBS & FLOWERS - VEGETABLES - FRUITS & BERRIES

ACTIVE INGREDIENTS:
Rotenone
Other Associated Resins
Pyrethrins0.8%
INERT INGREDIENTS
TOTAL

*This product contains an aromatic petroleum solvent

"PRECAUCION AL USUARIO: Si usted no lee ingles, no use este producto hasta que la etiqueta haya sido explicado ampliamente"

KEEP OUT OF REACH OF CHILDREN W A R N I N G STATEMENT OF PRACTICAL TREATMENT

Call a physician or Poison Control Center TF SWALLOWED: immediately. Promptly drink large quantities of milk, egg whites, gelatin solution, or, if these are not available, drink large quantities of water. Avoid alcohol. Do not induce vomiting. *Aromatic solvent content may present aspiration hazard. IF INHALED: Remove victim to fresh air. Apply artificial respiration if needed. IF IN EYES: Flush eyes with plenty of water. Get medical attention if irritation persists. IF ON SKIN: Remove contaminated clothing and wash affected area with soap and water. Get medical attention if irritation persists.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

WARNING: May be fatal if swallowed. Harmful if absorbed through the skin. Causes skin irritation. May cause substantial but temporary eye injury. Wear safety glasses. Do not get in eyes, on skin, or on clothing. Avoid breathing spray mist. Provide adequate ventilation in treatment area. Wash thoroughly with soap and water after handling and before eating or smoking. Remove contaminated clothing and wash before reuse. Avoid contamination of food, utensils and food preparation areas.

> ACCEPTED with COMMENTS in EPA Letter Dated

EPA REG. NO. EPA EST. NO.

3

JUN 24 1994 Under the Federel Insecticide. Fursicide, and Potenticide Art as zoended, for the pesticide registered under EPA Reg. No. 66665-5 PAGE 1

NOTE TO VETERINARIANS: If animal swallows significant amount of dip, administer an emetic, or gastric lavage if not convulsive, assist respiration as necessary. Deep IM of slow IV drip of diazepam to control convulsions. Calcium gluconate IV and Vitamin B should be administered to preserve liver functions.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to fish. Do not apply directly to water or wetlands ,swamps. bogs, marshes, potholes). Runoff and drift from treated areas may be hazardous to aquatic organisms in adjacent sites. Do not clean equipment or dispose of equipment washwaters in a manner that will contaminate water resources.

PHYSICAL AND CHEMICAL HAZARDS

Do not use, pour, spill or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal isw to use this product in a manner irconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. STORAGE: Store product in its original labeled container in a cool, dry, locked place out or reach of children. PESTICIDE DISPOSAL: Pestizide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance. CONTAINER DISPOSAL: Triple rinse or equivalent. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

FOR USE ON ORNAMENTAL TREES, SHRUBS & FLOWERING PLANTS; FOR VEGETABLES, FRUITS AND BERRIES

Mix this product with water as described in the following charts. Use a hand or automatic sprayer as a coarse spray. Use sufficient spray mix to give complete coverage of the plant. Repeat every 5 to 10 days or as needed. Mix only the amount of spray needed for one complete spray coverage. Do not feed treated foliage to livestock. Do not apply rotenone in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons. Do not enter treated areas without protective clothing until sprays have dried or dusts have settled.

EPA REG. NO. EPA EST. NO.

١

)

PAGE 2

ORNAMENTALS

For use on ornamental trees, shrubs and flowering plants such as Poplar, Linden, Dogwood, Willow, Birch, Boxelder, Maple, Oak, Elm, Pine, Spruce, Cedar, Yew, Spirea, Rose, Lilac, Zinnia, Aster, Petunia, Geranium, Dahlia, Gladiolus, Phlox, Sweet Pea, Marigold, Carnation and Chrysanthemum. Do not apply this product within one day of collecting flowers or cuttings.

AMOUNT TO USE/1 GALLON WATER PESTS OF ORNAMENTALS Aphids, Leafhoppers, Exposed Thrips l teaspoonful Oblique-Banded Leafrollers, Sawfly 2 teaspoonsful Larvae, Boxelder Bugs, Japanese Beetles, Elm Leaf Beetle Larvae, Mites, Lygus Bugs, Lace Bugs, Greenhouse Whiteflies, Stink Bugs, Garden Fleahoppers, Greenhouse Leaftiers, Cucumber Beetles, Rose Chafer, Ants, 4 teaspoonsful Aster Beetles, Flea Beetles

VEGETABLES For use on vegetable plants such as Peas, Beans, Beets, Cucumbers, Squash, Muskmelons, Potatoes, Cabbage, Spinach, Lettuce, Celery, Asparagus, Peppers, Eggplant, Onions, Carrots, and Corn. Do not apply within one day of harvest.

VEGETABLE PESTS

AMOUNT TO USE/1 GALLON WATER

)

)

Aphids, Leafhoppers, Exposed Thrips

Cabbage Worms, Bean Beetles, Beet Webwcrms, Sod Webworms, Asparagus Beetles, Japanese Beetle, Lygus Bugs, Whiteflies, Garden Fleahoppers, Harlequin Bugs, Oblique-Banded Leafrollers, Stink Bugs, Celery Leaftiers, Melon Worms, Tortoise Beetles, Garden Webworms, Alfalfa Webworms, and Mites.

EPA REG. NO. EPA EST. NO.

PAGE 3

BEST AVAILABLE COPY

1 teaspoonful

2 teaspoonsful



(Vegetable pests cont.)

Cucumber Beetles, Flea Beetles, Vegetable 4 teaspoonsful Weevils, Squash Bugs (nymphs), Squash Vine Borer, (European Corn Borers; 3 to 4 complete stalk & leaf applications or more, as eggs start to hatch. Repeat at 5 to 7 day intervals). Consult your Agricultural Extension Service Specialist for details.

FRUITS & BERRIES

For use on apples, pears, peaches, plums, cherries, grapes, currants, gooseberry, raspberry, blackberry, strawberry and blueberry, consult the following chart. Do not apply within one day of harvest.

PESTS ON FRUITS & BERRIES

}

)

Aphids, Leafhoppers, Exposed Thrips

Imported Currant Worms, Gooseberry Fruit Worms, Raspberry Fruit Worms, Blueberry Maggots, Cherry Fruit Flies, Strawberry Root Worms (adults), Red-necked Borers, Japanese Beetles and Mites.

Strawberry Leafrollers and Rose Chafers 2 teaspoonsful

4 teaspoonsful

l teaspoonful

AMOUNT TO USE/1 GALLON WATER

· · · ·

EPA REG. NO. EPA EST. NO.



PAGE 4

KILLS FLEAS, TICKS, AND LICE ON DOGS & CATS, AND IN THEIR LIVING QUARTERS

HOW TO APPLY TO YOUR DOG

While wearing neoprene or rubber household gloves, <u>spray</u>, or <u>sponge</u> in against the lay of pet hair until liquid penetrates to skin; (or) <u>dip</u> the entire pet's body making sure not to get head or face into dipping solution. Make sure to treat animal's back, head to tail, around the neck and ears, the underbelly, the legs, the base of the tail and where fleas, ticks or lice are seen. Do not get solution into pet's eyes, nose or mouth. Afterwards, <u>do not</u> <u>rinse pet since treatment will be ineffective</u>.

FOR PUPPIES

<u>Dilute 1/2 fluid ounce (1 tbsp.) of concentrate in 1 gallon of</u> <u>water</u> for moderate flea or tick infestations (or) <u>1 fluid ounce (2</u> <u>tbsp.) in 1 gallon of water</u> for heavy infestations. Do not exceed the 1 ounce per gallon rate for puppies. See HOW TO APPLY TO YOUR PET. Do not use on puppies less than 4 weeks old.

FOR ADULT DOGS

FLEAS & LICE: <u>Dilute 1 fluid ounce (2 tbsp.) per 1 gallon of</u> water. For young adults, use <u>1/2 fluid ounce per 1 gallon of</u> water. Apply directly to animal as a spray, or as a sponge-on wash. Dip animal until animal's skin is wet. See HOW TO APPLY.

TICKS: <u>Dilute at the rate of 2 fluid ounces (4 tbsp.) per 1 gallon</u> of water. For young animals, use <u>1 fluid ounce per 1 gallon of</u> water. Apply directly to animal as a spray, or as a sponge-on wash. Dip until animal's skin is wet. For ticks, examine pet's ears and treat as needed. See HOW TO APPLY.

NOTE: Avoid spray contact with pet's eyes, nose or mouth. Also to help prevent re-infestation, thoroughly spray animal's living and sleeping quarters (when pet is treated), and place freshly laundered bedding in pet's sleeping quarters.

IMPORTANT

Avoid spray or dip contact with pet's eyes, nose or mouth. Treat pet's ears for ticks. To prevent re-infestation, thoroughly spray the animal's living quarters and bedding with <u>2 fluid cunces per l</u> <u>gallon of water</u>. After treatment, place freshly laundered bedding in sleeping quarters. Repeat as needed for continued control through "flea" season. DO NOT USE THIS MATERIAL ON CATS.

EPA REG. NO. EPA EST. NO.

)

PAGE 5

1

Liquid Measuring Conversions

1. 1/2 oz. = 1 tablespoonful per gallon = 15 milliliters per 3.79 Liters.

- 2. 1 oz. = 2 tablespoonsful per gallon = 30 milliliters per 3.79
- Liters.
- 3. 2 oz. = 4 tablespoonsful per gallon = 60 milliliters per 3.79 Liters.

NOTICE: Buyer and user assume all risk and liability of use, storage and/or handling of this product not in accordance with the terms of this label.

Manufactured by

INNOVA CORPORATION Sugarland, TX 77478

EPA REG. NO. EPA EST. NO.

Ì

}

PAGE 6