

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

PRECAUTIONARY STATEMENTS

CAUTION

Harmful if swallowed. Wash hands thoroughly after handling. May cause eye irritation. Avoid eye contact.

STATEMENT OF PRACTICAL TREATMENT

Immediately contact physician or poison control center if swallowed. If eye contact occurs, flush eyes with plenty of water. Get medical attention if irritation persists.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

STORAGE: Store in a dry place. Do not store where children or animals may gain access.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of onsite or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

See side panel for Application Instructions.



FOR THE CONTROL OF DARKLING BEETLES, HIDE BEETLES, AND FLIES

| | |
|------------------------|--------|
| Active Ingredient: | |
| Orthoboric Acid..... | 99.0% |
| Inert Ingredients..... | 1.0% |
| Total | 100.0% |

- CAUTION -

KEEP OUT OF REACH OF CHILDREN

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONS

EPA REG. NO. 66986-1 EPA EST. NO. 64745-CA-001

A Product of:
PARASITIX CORPORATION
4980 Carroll Canyon Road
San Diego, California 92121
(619) 453-8030

NET WT: 50 pounds

PM-14 171
APPLICATION INSTRUCTIONS

FOR DARKLING AND LESSER MEALWORM BEETLE, HIDE BEETLE, AND FLY CONTROL IN POULTRY HOUSES. (Adults and Larvae)

For Poultry houses in which birds are grown on litter:
DRY APPLICATION

Prior to application of SafeCide Brand IC, remove birds. Apply dust uniformly to floor of poultry house or to old litter with a fertilizer or seed spreader, at the rate of 1-2 pounds for each 100 square feet, in bands along feeder lines. Wear a dust mask or a respirator while applying. Spread fresh litter of at least four inches uniformly over all treated area (floor or old litter). Then introduce birds. Reapply after each grow-out, if needed, in the sequence outlined above.

For poultry houses in which birds are grown in cages (layer or high rise 'plt type' houses):
DRY APPLICATION

Prior to application of SafeCide Brand IC, remove birds. Apply at the rate of 1-2 pounds for each 100 square feet of treated surface. Using appropriate equipment, thoroughly dust side walls, top plates, posts, and framing. Dust into cracks and crevices around insulation. Then introduce birds. Reapply after each grow-out, if needed, in the sequence outlined above. However, if facility is washed down and sanitized using standard cleaning and disinfecting procedures, SafeCide Brand IC must be reapplied to continue beetle control. SafeCide Brand IC can be applied directly to manure for fly control.

WET APPLICATION

Prior to application of SafeCide Brand IC to top levels of layer house, remove birds. Mix dust at the rate of 1-2 pounds per three gallons of water. Apply at the rate of three gallons of mix for each 100 square feet of treated surface. Using appropriate pressure spray equipment, saturate side walls, top plates, posts, and framing. Spray mix into cracks and crevices around insulation. Do not expose birds to spray. Reapply after each grow-out, if needed, in the sequence outlined above. However, if facility is washed down and sanitized using standard cleaning and disinfecting procedures, SafeCide Brand IC must be reapplied to continue beetle control. SafeCide Brand IC can be applied directly to manure for fly control.

P. 2/6
MAR 30 '94 10:46AM MYCOGEN LEGAL/MR