

HAZARDS TO HUMANS AND DOMESTIC ANIMALS PM-10

PRECAUTIONARY STATEMENTS CAUTION

Harmful if swallowed or absorbed through the skin. Causes eye irritation. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling.

Applicators and other handlers of this product must wear a dust mask or a dust/mist filtering respirator, and protective gloves for all application methods described on this label.

STATEMENT OF PRACTICAL TREATMENT

If Swallowed: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. If person is unconscious, do not give anything by mouth and do not induce vomiting.

If in Eyes: Flush eyes with plenty of water. Call a physician if irritation persists.

If on Skin: Wash with plenty of soap and water. Get medical attention.

For emergency medical information, call toll-free 1-800-228-5635 ext. 87

ENVIRONMENTAL HAZARDS

For terrestrial uses, do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

STORAGE: Store in a dry place. Do not store where children or animals may gain access.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of onsite or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. See side panel for Application Instructions.

Reg # 66986-1



IC

FOR THE CONTROL OF DARKLING BEETLES, HIDE BEETLES, AND FLIES IN POULTRY HOUSES

FOR THE CONTROL OF FIRE ANTS\*

Table with 2 columns: Active Ingredient, Inert Ingredients, Total. Values: Orthoboric Acid 99.0%, Inert Ingredients 1.0%, Total 100.0%


- CAUTION - KEEP OUT OF REACH OF CHILDREN

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONS

EPA REG. NO. 66986-1 EPA EST. NO. 64745-CA-001

A Product of: PARASITIX CORPORATION 5501 Oberlin Drive San Diego, California 92121 (619) 453-8030

NET WT: 50 pounds



191 APPLICATION INSTRUCTIONS

FOR DARKLING AND LESSER MEALWORM BEETLE, HIDE BEETLE, AND FLY CONTROL IN POULTRY HOUSES. (Adults and Larvae)

For poultry houses in which birds are grown on litter: DRY APPLICATION

Prior to application of SafeCide Brand IC, remove birds. Apply dust uniformly to floor of poultry house or to old litter with a fertilizer or seed spreader, at the rate of 1-2 pounds for each 100 square feet, in bands along feeder lines. Then introduce birds. Reapply after each grow-out, if needed, in the sequence outlined above.

For poultry houses in which birds are grown in cages (layer or high rise 'pl't type' houses): DRY APPLICATION

Prior to application of SafeCide Brand IC, remove birds. Apply at the rate of 1-2 pounds for each 100 square feet of treated surface. Using appropriate equipment, thoroughly dust side walls, top plates, posts, and framing. Dust into cracks and crevices around insulation. Then introduce birds. Reapply after each grow-out, if needed, in the sequence outlined above. However, if facility is washed down and sanitized using standard cleaning and disinfecting procedures, SafeCide Brand IC must be reapplied to continue beetle control. SafeCide Brand IC can be applied directly to manure for fly control.

WET APPLICATION

Prior to application of SafeCide Brand IC to top levels of layer house, remove birds. Mix dust at the rate of 1-2 pounds per three gallons of water. Apply at the rate of three gallons of mix for each 100 square feet of treated surface. Using appropriate pressure spray equipment, saturate side walls, top plates, posts, and framing. Spray mix into cracks and crevices around insulation. Do not expose birds to spray. Reapply after each grow-out, if needed, in the sequence outlined above. However, if facility is washed down and sanitized using standard cleaning and disinfecting procedures, SafeCide Brand IC must be reapplied to continue beetle control. SafeCide Brand IC can be applied directly to manure for fly control.

FOR THE CONTROL OF FIRE ANTS\*

METHODS OF APPLICATION

Apply SafeCide Brand IC when ants are actively foraging. This is usually when soil temperatures are above 60°F. Avoid application when the soil is wet. Heavy rainfall within 2 to 3 hours of application may reduce effectiveness. In cases where reinfestation occurs or when very large mounds remain active, retreatment may be desirable after 2 to 3 weeks. Do not apply to pasture, rangeland or other areas which may be grazed by cattle, sheep or other domestic animal.

MOUND APPLICATION

Apply 1 to 2 cups of SafeCide Brand IC per mound uniformly distributing material 3 to 4 feet around the base of the mound. Do not contaminate kitchen utensils by use or storage of product

\* Not for use in California