

Reg # 66986-2

PM-10

19/

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

PRECAUTIONARY STATEMENTS

CAUTION

Harmful if swallowed. Wash hands thoroughly after handling. May cause eye irritation. Avoid eye contact.

STATEMENT OF PRACTICAL TREATMENT

Immediately contact physician or poison control center if swallowed. If eye contact occurs, flush eyes with plenty of water. Get medical attention if irritation persists.

For emergency medical information, call toll-free
1-800-228-5635


STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

STORAGE: Store in a dry place. Do not store where children or animals may gain access.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of onsite or at an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.



BAIT

FOR THE CONTROL OF DARKLING BEETLES,
HIDE BEETLES, AND FLIES

Active Ingredient:

Orthoboric Acid.....30.0%

Inert Ingredients.....70.0%

Total 100.0%

- CAUTION -

KEEP OUT OF REACH OF CHILDREN

SEE SIDE PANEL FOR
ADDITIONAL PRECAUTIONS

EPA REG. NO. 66986-2

EPA EST. NO. 64222-GA-1 39573-TX-1 759-GA-1

Superscript corresponds to first number in lot number stamped on container

A Product of:
PARASITIX CORPORATION
4980 Carroll Canyon Road
San Diego, California 92121
(619) 463-8030

NET WT: 50 pounds

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

APPLICATION INSTRUCTIONS

FOR DARKLING AND LESSER MEALWORM BEETLE, HIDE BEETLE, AND FLY CONTROL IN POULTRY HOUSES. (Adults and Larvae)

For poultry houses in which birds are grown on litter:

Prior to application of SafeCide Brand Bait, remove birds. Apply bait uniformly to floor of poultry house or to old litter with a fertilizer or seed spreader, at the rate of 1-2 pounds per 100 square feet, in bands along feeder lines. Spread fresh litter of at least four inches uniformly over all treated area (floor or old litter). Then introduce birds. Reapply after each growout, if needed, in the sequence outlined above.

For poultry houses in which birds are grown in cages (layer or high rise 'pit type' houses):

Birds do not have to be removed prior to application of SafeCide Brand Bait. Apply bait uniformly to floor of poultry house with a fertilizer or seed spreader, at the rate of 1-2 pounds per 100 square feet, as needed, to maintain control. If facility is washed down and sanitized using standard cleaning and disinfecting procedures, SafeCide Brand Bait must be reapplied to continue beetle control. SafeCide Brand Bait can be applied directly to manure for fly control.

BEST AVAILABLE COPY

P.3/6

MAR 30 '94 10:48AM MYCOGEN LEGAL/HR