

PM 03

67505-3

8/13/97

1/6

spot-on label (full)
version of: 06/12/97

full label - FRONT PANEL

ECTO FLEA & TICK INSECTICIDE WITH IGR

PLEASANT CITRUS SCENT

FLEA & TICK CONTROL FOR DOGS AND PUPPIES OVER 12 WEEKS OF AGE

- BREAKS FLEA LIFE CYCLE!
- KILLS & REPELS FLEAS UP TO 3 TO 4 WEEKS!
- KILLS FLEA EGGS AND LARVAE FOR UP TO 123 DAYS!
- PREVENTS FLEA EGGS FROM DEVELOPING INTO BITING ADULTS FOR UP TO 123 DAYS!
- KILLS AND REPELS MOSQUITOES!
- PROTECTS AGAINST BLOOD FEEDING BY MOSQUITOES (vector of HEARTWORM) UP TO 4 WEEKS!
- KILLS & REPELS TICKS FOR UP TO 4 WEEKS!
- KILLS & REPELS DEER TICKS (vector of LYME DISEASE) FOR UP TO 4 WEEKS!
- KILLS & REPELS BROWN DOG TICKS (Rhipicephalus sanguineus) FOR UP TO 3 TO 4 WEEKS!
- KILLS & REPELS AMERICAN DOG TICKS (Dermacentor variabilis) FOR UP TO 3 TO 4 WEEKS!

ACTIVE INGREDIENTS


* Permethrin:	45.0%
**Pyriproxyfen:	5.0%
INERT INGREDIENTS.....	50.0%
TOTAL	100.0%

* (3-phenoxyphenyl) methyl(+)-cis, trans-3-(2,2-dichloroethenyl)-
2,2-dimethylcyclopropanecarboxylate
cis/trans ratio: Max 55% (+) cis and min 45% (+) trans

** 2-[1-methyl-2-(4-phenoxyphenoxy) ethoxy] pyridine
Nylar Registered Trademark of McLaughlin Gormley King and Company

Patent Pending

KEEP OUT OF THE REACH OF CHILDREN
CAUTION
DO NOT USE ON CATS
SEE BACK PANEL FOR ADDITIONAL
PRECAUTIONARY STATEMENTS



Net Contents: Two 1.5mL Applicators

EPA Reg. No.: 67505-3

EPA Est. No.:

Manufactured For: Ecto Development Corporation
1229 N. Jesse James Road
Excelsior Springs, MO 64024

IN CASE OF EMERGENCY CALL: 816-637-2015

(lot number statements as needed)

Mfg. & Printed in U.S.A.

(label revision number)

(product number)

spot-on label (full)
version of: 06/12/97

2/6
READ ENTIRE LABEL BEFORE EACH USE
USE ONLY ON DOGS OVER 12 WEEKS OF AGE
DO NOT USE ON CATS

ECTO FLEA & TICK INSECTICIDE WITH IGR

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Hazards to Humans: Harmful if swallowed. Avoid contact with eyes or clothing. Causes moderate eye irritation. Wash thoroughly with soap and water after handling.

Hazards to Domestic Animals: FOR EXTERNAL USE ON DOGS ONLY. Do not use on puppies under 12 weeks of age. Consult a veterinarian before using this product on debilitated, aged, medicated, pregnant, or nursing dogs. Consult a veterinarian before using on dogs with known organ dysfunction. DO NOT USE ON CATS or animals other than dogs. Cats which actively groom or engage in close physical contact with recently treated dogs may be at risk of toxic exposure. Certain medications can interact with pesticides. It is advisable to consult a veterinarian before using this product with any other pesticide or drug.

FIRST AID

If Swallowed: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger or, if available, by administration of syrup of ipecac. DO NOT INDUCE VOMITING OR GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON.

If In Eyes: Flush eyes with plenty of water. Call a physician if irritation persists.

Adverse Reactions: Some animals may be sensitive to ingredients in this product. Reactions in dogs may include skin sensitivity. Dogs may show lethargy, increased pruritis (itchiness), erythema (redness), rash and hair discoloration or hair loss at the application site. Observe the dog following treatment. Sensitivity may occur after using ANY pesticide product on pets. If signs of sensitivity occur bathe your dog with a mild, non-insecticidal shampoo and rinse with large amounts of water. If signs continue, consult a veterinarian immediately.

ENVIRONMENTAL HAZARDS

This product is extremely toxic to fish. Do not add directly to water. Do not contaminate water when disposing of product or packaging.

spot-on label (full)
version of: 06/12/97

3/6

ECTO FLEA & TICK INSECTICIDE WITH IGR

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DO NOT USE ON CATS or animals other than dogs. Do not get this product in your dog's eyes or mouth. Repeat applications may be made if necessary, but do not apply more often than once every 3 weeks, except to reapply after shampooing the dog.

For Dogs Weighing Less Than 33 pounds: Apply one tube (1.5 mL) of ECTO FLEA & TICK INSECTICIDE WITH IGR solution as a spot or stripe to the dog's back between the shoulder blades.

For Dogs Weighing More Than 33 pounds: Apply one tube (1.5 mL) of ECTO FLEA & TICK INSECTICIDE WITH IGR solution as a spot or stripe to the dog's back between the shoulder blades and apply a second tube (1.5 mL) as a spot or strip to the dog's back directly in front of the base of the tail.

HOW TO APPLY:

FOR FOLDING CARD OR TABBED CARD UNIT-DOSE PACKAGING: Remove product tab by tearing along perforation. Holding tab with tube end pointed up and away from face and body, cut along dotted line. Invert tab over dog and use narrow end of card to part dog's hair. Squeeze tube firmly to apply all of the solution to the dog's skin. Wrap tab and put in trash.

FOR FLAT CARD (doubled) UNIT-DOSE PACKAGING: Separate cards along the perforated line. Holding separated card by its wide end with narrow end pointed up and away from face and body, cut along dotted line. Invert card over dog and use narrow end of the card to part dog's hair. Squeeze tube firmly to apply all of the solution to the dog's skin. Wrap card and put in trash.

**** FOR MULTI-DOSE PACKAGING:** Withdraw 1.5 mL of solution at a time. Apply entire given dose directly to the skin at the indicated locations.

**** FOR DROPPER BOTTLE MULTI-DOSE PACKAGING:** Insert dropper tube into insecticide solution, squeeze bulb and then release to fill tube barrel to 1.5 mL line. Position dropper tube end in dog's hair next to skin and squeeze bulb to apply all of the solution to the dog's skin.

**** FOR SALE TO VETERINARIANS ONLY**

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Storage: Store in cool, dry place. Protect from freezing.

Pesticide Disposal: Securely wrap original container in several layers of newspaper and discard in trash.

Container Disposal: Do not reuse empty container. Wrap container and put in trash.

spot-on label (full)
version of: 06/12/97

4/6

ECTO FLEA & TICK INSECTICIDE WITH IGR

ECTO FLEA & TICK INSECTICIDE WITH IGR is an effective and easy to use product. ECTO FLEA & TICK INSECTICIDE WITH IGR has demonstrated greater than 92% control of fleas within one day of application. ECTO FLEA & TICK INSECTICIDE WITH IGR prevents eggs from fleas on treated dogs from developing into biting adults for up to 123 days. As with all flea and tick control products, ECTO FLEA & TICK INSECTICIDE WITH IGR should be used as part of a program aimed at reducing flea populations in the dog's environment (bedding, carpets, kennel, yard). Consult your veterinarian, entomologist, or retailer for program recommendations.

NOTICE OF WARRANTY

ECTO DEVELOPMENT CORPORATION MAKES NO WARRANTY OF MERCHANTABILITY, FITNESS FOR ANY PARTICULAR PURPOSE, OR OTHERWISE, EXPRESSED OR IMPLIED CONCERNING THIS PRODUCT OR ITS USES WHICH EXTEND BEYOND THE USE OF THE PRODUCT UNDER NORMAL CONDITIONS IN ACCORDANCE WITH THE STATEMENTS MADE ON THIS LABEL.

spot-on tube label
version of: 06/12/97

TUBE LABEL (folding card and flat card packaging designs)

ECTO FLEA & TICK INSECTICIDE WITH IGR

PERMETHRIN 45% W/W
PYRIPROXYFEN 5% W/W

KEEP OUT OF THE REACH OF CHILDREN
READ DIRECTIONS AND
PRECAUTIONS BEFORE USING

EPA Reg. No. 67505-3

6/6

spot-on tab label
version of: 06/12/97

TAB LABEL (folding card and tabbed card packaging designs)

ECTO FLEA & TICK INSECTICIDE WITH IGR

ACTIVE INGREDIENTS

Permethrin: (3-phenoxyphenyl) methyl(±)-cis, trans-3-(2,2-dichloroethenyl)-
2,2-dimethylcyclopropanecarboxylate* 45.0%
Pyriproxyfen: 2-[1-methyl-2-(4-phenoxyphenoxy) ethoxy] pyridine..... 5.0%

* cis/trans ratio: Max 55% (±) cis and min 45% (±) trans

KEEP OUT OF THE REACH OF CHILDREN
SEE MAIN LABEL FOR ADDITIONAL
PRECAUTIONS AND INSTRUCTIONS FOR USE

EPA Reg. No. 67505-3