67505-3

11-29-2010



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, D.C. 20460

OFFICE OF CHEMICAL SAFETY ND POLLUTION PREVENTION NOV 2 9 2010

Mr. Joseph E. Dyer ECTO Development Corporation 1229 Jesse James Road Excelsior Springs, MO 64024-1186

Dear Mr. Dyer:

Subject: Amendment- based on Agency letter dated November 1, 2010 Ecto Flea & Tick Insecticide with IGR EPA Registration Number 67505-3 Your submission dated November 10, 2010

The application referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is

A copy of a stamped label is enclosed for your records.

If you have any questions regarding this action, please contact Linda A. DeLuise of my team at (703) 305-5428.

Sincerely yours,

Richard Gebken Product Manager (13) Insecticide Branch Registration Division (7504P)

Enclosure

FRONT PANEL (ECIO FLEA & TICK INSECTICIDE WITH IGR

· ~ •

¢.

[CAT ICON IN CIRCLE WITH SLASH THROUGH IT] 2

٩

		PLEASANT CIT K CONTROL FOR DOGS AND BIO SPOT FLEA & TIC] [Between 15 & 33 lbs	PUPPIES OVER 12 W K CONTROL FOR DOGS	5							
		DO NOT USE	ON CATS								
	BREAKS FLEA LIFE CY										
		S UP TO 3 TO 4 WEEKS! LARVAE FOR UP TO 123	חאעכו								
		FROM DEVELOPING INTO B		P TO 123 DAYSI							
	KILLS AND REPELS MO			10 120 14110.							
0	PROTECTS AGAINST BLOOD FEEDING BY MOSQUITOES (vector of HEARTWORM) UP TO 4 WEEKS!										
		S FOR UP TO 4 WEEKS!									
0	KILLS & REPELS DEER	TICKS (which may carr	y Lyme disease) FC	R UP TO 4 WEEKS!							
0 0	KILLS & REPELS BROWN KILLS & REPELS AMER WEEKS!	N DOG TICKS (<u>Rhipiceph</u> ICAN DOG TICKS (<u>Dermac</u>	<u>alus sanguineus);</u> entor variabilis) //	FOR US HOP TO A D							
0	MONTHLY APPLICATION	<u>I</u>									
		AND MOSQUITO TREATMENT		NOV 2 9 2010							
		EAS AND TICKS FOR UP T JLT FLEAS FOR UP TO 3		The Rederal Insection							
0	KILLS FLEA EGGS AND	JLT FLEAS FOR UP TO 3 LARVAE [FOR UP TO 4 M CKS FOR UP TO [4 WEEKS	ONTHS] [FOR 123 JA	angleide, and Rodenticide Act.							
•	KILLS AND REPELS TIC	CKS FOR UP TO [4 WEEKS] [3 TO 4 WEEKS]]	NETERING DEER TICKS							
	(which may carry Ly	me disease), BROWN DOG	TICKS & AMERICAN	DOG TICKS!							
0	KILLS AND REPELS [A]	DULT FLEAS] FOR UP TO	3 TO 4 WEEKS!								
E	yriproxyfen: (CAS # ER INGREDIENTS cis/trans ratio: I U.S. Patent No. 5		TC n 45% (<u>+</u>) trans	5.0% <u>50.0%</u> TAL 100.0%							
	(Nylar**) is a Registered Trademark of McLaughlin Gormley King and Company KEEP OUT OF THE REACH OF CHILDREN DO NOT USE ON CATS										
CAUTION SEE SIDE/BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS											
Net		l. oz. (1 mL)][0.05 fl] Applicator	. oz.(1.5mL)][1 fl	.oz (3.0 mL)]							
EPA	Reg. No.: 67505-3		EPA Est. No.:	ι ττίτι ι							
	ufactured For:	Ecto Development Corp Excelsior Springs, MC									
Pac Nov	•	(lot number statement label revision number)		Printed in U.S.A.							
				() () () () () () () () () () () () () (

SIDE/BACK PANEL FLEA & TICK INSECTICIDE WITH IGR

READ ENTIRE LABEL BEFORE EACH USE USE ONLY ON DOGS OVER 12 WEEKS OF AGE DO NOT USE ON CATS

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Hazards to Humans: Harmful if swallowed. Avoid contact with eyes or clothing. Causes moderate eye irritation. Wash thoroughly with soap and water after handling.

Hazards to Domestic Animals: FOR EXTERNAL USE ON DOGS ONLY. Do not use on puppies under 12 weeks of age. Consult a veterinarian before using this product on debilitated, aged, medicated, pregnant, or nursing dogs. Consult a veterinarian before using on dogs with known organ dysfunction. DO NOT USE ON CATS or animals other than dogs. Cats which actively groom or engage in close physical contact with treated dogs may be at risk of serious harmful effect. Certain medications can interact with pesticides. It is advisable to consult a veterinarian before using this product with any other pesticide or drug.

FIRST AID							
 If Swallowed ° Call a poison control center or doctor immediately for treatment advice. ° Have person sip a glass of water if able to swallow. ° Do not induce vomiting unless told to do so by a poison control center or doctor. ° Do not give anything by mouth to an unconscious person. 							
If In Eyes	 Hold eyes open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eyes. Call a poison control center or doctor for treatment advice. 						
HOT LINE NUMBER							
Have the product container or label with you when calling a poison control center							

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-858-7378 for emergency medical treatment information.

Adverse Reactions: Some animals may be sensitive to ingredients in this product. Reactions in dogs may include skin sensitivity. Dogs may show lethargy, increased pruritis (itchiness), erythema (redness), rash and hair discoloration or hair loss at the application site. Observe the dog following treatment. Sensitivity may occur after using ANY pesticide product on pets. If signs of sensitivity occur bathe your dog with a mild, non-insecticidal shampoo and rinse with large amounts of water. If signs continue, consult a veterinarian immediately.

67505-3 Page 2 of 8 November 10, 2010 E03DL01D.DOC-2 <u>ン</u> 9

SIDE/BACK PANEL ECIO FLEA & TICK INSECTICIDE WITH IGR

ENVIRONMENTAL HAZARDS

This product is extremely toxic to fish. Do not add directly to water. Do not contaminate water when disposing of product or packaging.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DO NOT USE ON CATS or animals other than dogs. May be toxic and potentially fatal if applied to or ingested by cats. Do not get this product in your dog's eyes or mouth. Repeat applications may be made if necessary, but do not apply more often than once every 3 weeks, except to reapply after shampooing the dog.

[For Dogs Weighing 15 lbs and Under: (1.0 mL applicator tubes) Apply one tube (1.0 mL) of product solution as a spot or stripe to the dog's back between the shoulder blades.]

[For Dogs Weighing Between 15 and 33 lbs: (1.5 mL applicator tubes) Apply one tube (1.5 mL) of product solution as a spot or stripe to the dog's back between the shoulder blades.]

[For Dogs Weighing Between 33 to 66 lbs: (1.5 mL) applicator tubes) Apply one tube (1.5 mL) of product solution as a spot or stripe to the dog's back between the shoulder blades <u>and</u> apply a second tube (1.5 mL) as a spot or strip to the dog's back directly in front of the base of the tail.] OR

[(3 mL applicator tubes) Apply approximately half of the tube (1.5 mL) of product solution as a spot or stripe to the dog's back between the shoulder blades <u>and</u> apply the rest of the tube's contents (1.5 mL) as a spot or strip to the dog's back directly in front of the base of the tail.] OR

[(3 mL applicator tube) Apply the entire tube (3.0 mL) as a continuous stripe on the dog's back starting between the shoulder blades and ending directly in front of the base of the tail.]

[For Dogs Weighing 66 lbs and Over: (1.5 mL applicator tubes) Apply one tube (1.5 mL) of product solution as a stripe to the dog's back between the shoulder blades <u>and</u> apply a second tube (1.5 mL) continuing the stripe down the center of the dog's back <u>and</u> apply a third tube (1.5 mL) to the dog's back directly in front of the base of the tail.] OR

[(4.5 mL applicator tubes) Apply approximately half of the tube (2.25mL) of product solution as a spot or stripe to the dog's back between the shoulder blades <u>and</u> apply the rest of the tube's contents (2.25 mL) as a spot or strip to the dog's back directly in front of the base of the tail.] OR

[(4.5 mL applicator tube) Apply the entire tube (4.5 mL) as a continuous stripe on the dog's back starting between the shoulder blades and ending directly in front of the base of the tail.]

67505-3 Page 3 of 8 November 10, 2010 E03DL01D.DOC-3

ECTO FLEA & TICK INSECTICIDE WITH IGR

DIRECTIONS FOR USE (Continued)

HOW TO APPLY:

FOR SARAN TUBES IN BLISTER CARD: Remove product tube from package. Holding tube with notched end pointing up and away from face and body, snap or cut off narrow end at notches. Invert tube over dog and use open end to part dog's hair. Squeeze tube firmly to apply all of the solution to the dog's skin. Wrap tube and put in trash.

FOR FLAT CARD (doubled) UNIT-DOSE PACKAGING: Separate cards along the perforated line. Holding separated card by its wide end with narrow end pointed up and away from face and body, cut along dotted line. Invert card over dog and use narrow end of the card to part dog's hair. Squeeze tube firmly to apply all of the solution to the dog's skin. Wrap card and put in trash.

FOR FLAT CARD UNIT DOSE PACKAGING: [For push pin type applicator] Remove one product tube from card. With cap end pointed up and away from face and body, depress cap to pierce tube end. Carefully remove cap, invert tube over dog, and with tube in close to dogs skin, squeeze tube to apply product. Wrap empty tube and put in trash.

FOR FOLDING CARD OR TABBED CARD UNIT-DOSE PACKAGING: Remove product tab by tearing along perforation. Holding tab with tube end pointed up and away from face and body, cut along dotted line. Invert tab over dog and use narrow end of card to part dog,s hair. Squeeze tube firmly to apply all of the solution to the dog's skin. Wrap tab and put in trash.

BLISTER CARD WITH TRI-LAYER TUBES: Remove product tube(s) from package. Fold along perforation between tubes and tear off one tube. {Return unused tubes to package.} Holding tube with notched end pointing up and away from face and body, snap off narrow end at notches. Invert tube over dog and use open end to part dog's hair. Squeeze tube firmly to apply all of the solution to the dog's skin. Wrap tube and put in trash.

FOR MULTI-DOSE PACKAGING: Withdraw 1.0 mL (1.5 mL)(3.0 mL)(4.5 mL) of solution at a time. Apply entire given dose directly to the skin at the indicated locations.

FOR DROPPER BOTTLE MULTI-DOSE PACKAGING: Insert dropper tube into insecticide solution, squeeze bulb and then release to fill tube barrel to 1.0 mL (1.5 mL) (3.0 mL) (4.5 mL) line. Position dropper tube end in dog's hair next to skin and squeeze bulb to apply all of the solution to the dog's skin as indicated for dog's size.

FOR SALE TO VETERINARIANS ONLY

67505-3 Page 4 of 8 November 10, 2010 E03DL01D.DOC-4

SIDE/BACK PANEL ECTO FLEA & TICK INSECTICIDE WITH IGR

ECIO FLEA & TICK INSECTICIDE WITH IGR is an effective and easy to use product. This product has demonstrated greater than 92% control of fleas within one day of application. This product prevents eggs from fleas on treated dogs from developing into biting adults for up to [4 months] 123 days. As with all flea and tick control products, This product should be used as part of a program aimed at reducing flea populations in the dog's environment (bedding, carpets, kennel, yard). Consult your veterinarian, entomologist, or retailer for program recommendations.

STORAGE AND DISPOSAL

Storage: Store in a cool dry place away from heat and open flame. Protect from freezing.

Disposal: Nonrefillable container. If empty: Do not reuse or refill this container. Place in Trash or offer for recycling if available. If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

NOTICE OF WARRANTY

ECTO DEVELOPMENT CORPORATION MAKES NO WARRANTY OF MERCHANIABILITY, FITNESS FOR ANY PARTICULAR PURPOSE, OR OTHERWISE, EXPRESSED OR IMPLIED CONCERNING THIS PRODUCT OR ITS USES WHICH EXTEND BEYOND THE USE OF THE PRODUCT UNDER NORMAL CONDITIONS IN ACCORDANCE WITH THE STATEMENTS MADE ON THIS LABEL.

<u> OR</u>

Buyer assumes all responsibility for safety and use not in accordance with directions.

The use of any notice of warranty statement is optional.

ALTERNATE MARKETING CLAIMS BASED ON ME-TOO APPLICATION:

E03DL01D.DOC-5

For use on Dogs and puppies over 12 weeks of age [Kills, repells and prevents re-infestation of fleas for up to 5 weeks] [Kills, repells and prevents re-infestation for fleas for up to 4 weeks] [Kills flea eggs and larvae for up to 8 weeks] [Kills flea eggs and larvae for up to 6 weeks] [Kills flea eqgs and larvae for up to 4 weeks] [Breaks the flea life cycle] [Kills, repels and prevents re-infestation of ticks for up to 6 weeks] [Kills, repels and prevents re-infestation of ticks for up to 4 weeks] [Kills, repels mosquitoes, lice and mites on dogs for up to 4 weeks] [Kills five different kinds of pests on dogs] [Kills 5 different kinds of pests on dogs] [Kills five different canine pests] [Kills 5 different canine pests] [Kills canine pests belonging to Five different orders] [Kills canine pests belonging to 5 different orders] [Seven in one Protection] [7 in 1 Protection] [Six in 1 Protection] [6 in 1 Protection] [Five in 1 Protection] [5 in 1 Protection] 67505-3 Page 5 of 8 November 10, 2010

6 9

TUBE LABEL (folding card and flat card packaging designs)

,

. .

ECIO FLEA & TICK INSECTICIDE WITH IGR

PERMETHRIN 45% W/W PYRIPROXYFEN 5% W/W

KEEP OUT OF THE REACH OF CHILDREN

DO NOT USE ON CATS OR [CAT ICON IN CIRCLE WITH SLASH THROUGH IT]

READ DIRECTIONS AND PRECAUTIONS BEFORE USING

> EPA Reg. No. 67505-3 Lot No. 000000

67505-3 Page 6 of 8 November 10, 2010 E03DL01D.DOC-6 TAB LABEL (folding card and tabbed card packaging designs)

ECTO FLEA & TICK INSECTICIDE WITH IGR

8

ACTIVE INGREDIENTS

. .

		#52645-53-1)			
Pyriproxyfen	: (CA	\$ #95737-68-1)	 	 	 5.0%

* cis/trans ratio: Max 55% (+) cis and min 45% (+) trans

KEEP OUT OF THE REACH OF CHILDREN

DO NOT USE ON CAIS OR [CAI ICON IN CIRCLE WITH SLASH THROUGH IT]

SEE MAIN LABEL FOR ADDITIONAL PRECAUTIONS AND INSTRUCTIONS FOR USE

EPA Reg. No. 67505-3 Lot No. 000000

67505-3 Page 7 of 8 November 10, 2010 E03DL01D.DOC-7

ECIO FLEA & TICK INSECTICIDE WITH IGR

BIOLAR (NYLAR) inhibits development of the immature stages of the flea for up to 4 months preventing them from reaching the biting adult stage.

(Nylar) Registered Trademark of McLaughlin Gormley King and Company

PICTURE OF FLEA LIFE CYCLE HERE

As Part of a complete flea control program USE Bio Flea Halt! Outdoor Yard Spray Concentrate, and Bio Flea Halt! Carpet Powder or Carpet Spray with Biolar Insect Growth Regulator, along with on-animal flea & tick products.

4 STEPS FOR COMPLETE FLEA CONTROL

- Step 1 Shampoo or dip your dog
- Step 2 Treat indoor and outdoor premise with Bio Flea Halt! indoor and outdoor sprays, foggers and powders
- Step 3 Apply Bio Spot-On for up tp four weeks adult flea and tick control
- Step 4 Maintenance treatments with on-animal sprays and powders

67505-3 Page 8 of 8 November 10, 2010 E03DL01D.DOC-8