spot-on label (full) version of: 11/04/98

full label - FRONT PANEL

ECTO FLEA AND TICK INSECTICIDE

PLEASANT CITRUS SCENT FLEA & TICK CONTROL FOR DOGS

- KILLS & REPELS FLEAS UP TO 3 TO 4 WEEKS!
- KILLS AND REPELS MOSQUITOES!
- PROTECTS AGAINST BLOOD FEEDING BY MOSQUITOES (vector of HEARTWORM) UP TO 4 WEEKS!
- ° KILLS & REPELS TICKS FOR UP TO 4 WEEKS!
- ° KILLS & REPELS DEER TICKS (vector of LYME DISEASE) FOR UP TO 4 WEEKS!
- ° KILLS & REPELS BROWN DOG TICKS (Rhipicephalus sanguineus) FOR UP TO 3 TO 4 WEEKS!
- KILLS & REPELS AMERICAN DOG TICKS (<u>Dermacentor variabilis</u>) FOR UP TO 3 TO 4 WEEKS!
- ° MONTHLY APPLICATION!
- O MONTHLY FLEA, TICK AND MOSQUITO TREATMENT!
- ° KILLS AND REPELS FLEAS AND TICKS FOR UP TO A MONTH!

ACTIVE INCREDIENTS

TOTAL

(3-phenoxyphenyl) methyl(±)-cis, trans-3-(2,2-dichloroethenyl)-2,2-dimethylcyclopropanecarboxylate cis/trans ratio: Max 55% (\pm) cis and min 45% (\pm) trans

KEEP OUT OF THE REACH OF CHILDREN CAUTION DO NOT USE ON CATS SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

Net Contents: Two 1.5mL Applicators

EPA Req. No.: 67505-4

EPA Est. No.: 67517-MO-1

ACCEPTEE

Manufactured For:

Ecto Development Corporation 1229 N. Jesse James Road Excelsior Spring, MO 64024

IN CASE OF EMERGENCY CALL: 816-637-2015

(lot number statements as needed) Mfg. & Printed in U.S.A. (label revision number)

ECTO4LAA.DOC~1

spot-on label (full) version of: 11/04/98

READ ENTIRE LABEL BEFORE EACH USE DO NOT USE ON DOGS DO NOT USE ON DOGS UNDER 6 MONTHS OF AGE DO NOT USE ON CATS

ECTO FLEA & TICK INSECTICIDE

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Hazards to Humans: Harmful if swallowed. Avoid contact with eyes or clothing. Causes moderate eye irritation. Wash thoroughly with soap and water after handling.

Hazards to Domestic Animals: FOR EXTERNAL USE ON DOGS ONLY. Do not use on dogs under 6 months of age. Consult a veterinarian before using this product on debilitated, aged, medicated, pregnant, or nursing dogs. Consult a veterinarian before using on dogs with known organ dysfunction. DO NOT USE ON CATS or animals other than dogs. Cats which actively groom or engage in close physical contact with recently treated dogs may be at risk of toxic exposure. Certain medications can interact with pesticides. It is advisable to consult a veterinarian before using this product with any other pesticide or drug.

FIRST AID

If Swallowed: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger or, if available, by administration of syrup of ipecac. DO NOT INDUCE VOMITING OR GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON.

If In Eyes: Flush eyes with plenty of water. Call a physician if irritation persists.

Adverse Reactions: Some animals may be sensitive to ingredients in this product. Reactions in dogs may include skin sensitivity. Dogs may show lethargy, increased pruritis (itchiness), erythema (redness), rash and hair discoloration or hair loss at the application site. Observe the dog following treatment. Sensitivity may occur after using ANY pesticide product on pets. If signs of sensitivity occur bathe your dog with a mild, non-insecticidal shampoo and rinse with large amounts of water. If signs continue, consult a veterinarian immediately.

PHYSICAL OR CHEMICAL HAZARDS

DO NOT USE OR STORE NEAR HEAT OR OPEN FLAMES

ENVIRONMENTAL HAZARDS

This product is extremely toxic to fish. Do not add directly to water Do not contaminate water when disposing of product or packaging.

spot-on label (full) version of: 11/04/98

ECTO FLEA & TICK INSECTICIDE

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DO NOT USE ON CATS or animals other than dogs. Do not get this product in your dog's eyes or mouth. Repeat applications may be made if necessary, but do not apply more often than once every 3 weeks, except to reapply after shampooing the dog.

For Dogs Weighing Less Than 33 pounds: [1.5 ml applicator tube] Apply one tube (1.5 mL) of ECTO FLEA & TICK INSECTICIDE solution as a spot between the shoulder blades or as a stripe to the dog's back.

For Dogs Weighing More Than 33 pounds: [1.5 ml applicator tube] Apply one tube (1.5 mL) of ECTO FLEA & TICK INSECTICIDE solution as a spot to the dog's shoulder blades or as a stripe to the dog's back and apply a second tube (1.5 mL) as a spot directly in front of the base of the tail or as a stipe to the dogs back.

For Dogs Weighing More Than 33 pounds: [3.0 ml applicator tube] Apply approximately half of the tube (1.5 mL) of ECTO FIEA & TICK INSECTICIDE solution as a spot or stripe to the dog's back between the shoulder blades and apply the rest of the tube's contents (1.5 mL) as a spot or strip to the dog's back directly in front of the base of the tail. Or, apply the entire tube (3.0 ml) as a continuous stripe on the dog's back starting between the shoulder blades and ending directly in front of the base of the tail.

HOW TO APPLY:

FOR FOLDING CARD OR TABBED CARD UNIT-DOSE PACKAGING: Remove product tab by tearing along perforation. Holding tab with tube end pointed up and away from face and body, cut along dotted line. Invert tab over dog and use narrow end of card to part dog,s hair. Squeeze tube firmly to apply all of the solution to the dog's skin. Wrap tab and put in trash.

FOR FIAT CARD (doubled) UNIT-DOSE PACKAGING: Separate cards along the perforated line. Holding separated card by its wide end with narrow end pointed up and away from face and body, cut along dotted line. Invert card over dog and use narrow end of the card to part dog's hair. Squeeze tube firmly to apply all of the solution to the dog's skin. Wrap card and put in trash.

FOR FLAT CARD UNIT DOSE PACKAGING: [For push pin type applicator] Remove one product tube from card. With cap end pointed up and away from face and body, depress cap to pierce tube end. Carefully remove cap, invert tube over dog, and with tube in close to dogs skin, squeeze tube to apply product. Wrap empty tube and put in trask.

- ** FOR MULTI-DOSE PACKAGING: Withdraw 1.5 mL of solution at a time. Apply entire given dose directly to the skin at the indicated locations.
- ** FOR DROPPER BOTTLE MULTI-DOSE PACKAGING: Insert dropper tube into insecticide; solution, squeeze bulb and then release to fill tube barrel to 1.5 mL line. Position dropper tube end in dog's hair next to skin and squeeze bulb to apply all of the solution to the dog's skin.

^{**} FOR SALE TO VETERINARIANS ONLY

spot-on label (full)
version of: 11/04/98

ECTO FLEA & TICK INSECTICIDE

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Storage: Store in cool, dry place. Protect from freezing.


Pesticide Disposal: Securely wrap original container in several layers of newspaper and discard in trash.

Container Disposal: Do not reuse empty container. Wrap container and put in trash.

ECTO FLEA & TICK INSECTICIDE is an effective and easy to use product. ECTO FLEA & TICK INSECTICIDE has demonstrated greater than 92% control of fleas within one day of application. As with all flea and tick control products, ECTO FLEA & TICK INSECTICIDE should be used as part of a program aimed at reducing flea populations in the dog's environment (bedding, carpets, kennel, yard). Consult your veterinarian, entomologist, or retailer for program recommendations.

NOTICE OF WARRANTY

ECTO DEVELOPMENT CORPORATION MAKES NO WARRANTY OF MERCHANTABILITY, FITNESS FOR ANY PARTICULAR PURPOSE, OR OTHERWISE, EXPRESSED OR IMPLIED CONCERNING THIS PRODUCT OR ITS USES WHICH EXTEND BEYOND THE USE OF THE PRODUCT UNDER NORMAL CONDITIONS IN ACCORDANCE WITH THE STATEMENTS MADE ON THIS LABEL.



spot-on tube label
version of: 11/04/98


TUBE LABEL (folding card and flat card packaging designs)

ECTO FLEA & TICK INSECTICIDE

PERMETHRIN 45% W/W

KEEP OUT OF THE REACH OF CHILDREN
READ DIRECTIONS AND
PRECAUTIONS BEFORE USING

EPA Reg. No. 67505-4



spot-on tab label
version of: 11/04/98

TAB LABEL (folding card and tabbed card packaging designs)

ECTO FLEA & TICK INSECTICIDE

ACTIVE INGREDIENTS
Permethrin: (3-phenoxyphenyl) methyl(±)-cis, trans-3-(2,2-dichloroethenyl)-
2,2-dimethylcyclopropanecarboxylate*

* cis/trans ratio: Max 55% (±) cis and min 45% (±) trans

KEEP OUT OF THE REACH OF CHILDREN SEE MAIN LABEL FOR ADDITIONAL PRECAUTIONS AND INSTRUCTIONS FOR USE

EPA Reg. No. 67505-4

