

11/30/2004

1/4

NOV 30 2004

H.R. McLane
H.R. McLane Inc.
7201 Red Road, Suite 206A
Miami, FL 33143

Dear Mr. McLane:

SUBJECT: Golf Ventures, Inc.
Ronstar 1[®] With Fertilizer
Label Amendment
EPA Registration No. 67508-1
Your Application Dated October 25, 2004

The label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable provided that the following revisions are made:

1. At the beginning of the list of Personal Protective equipment (PPE) within the Precautionary Statements, add the statements "Some of the materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart." In addition, revise the requirement for "waterproof gloves" to a requirement for "chemical-resistant gloves made of any waterproof material such as polyethylene or polyvinyl chloride."
2. Within the list of PPE for early re-entry in the Agricultural Use Requirements box, revise the requirement for "waterproof gloves" to a requirement for "chemical-resistant gloves made of any waterproof material"

The amended label supersede all previously accepted ones. A stamped copy of the label is enclosed for your records. Please submit one copy of the final printed label before you release the product for shipment.

Sincerely yours,

|S|

Joanne I. Miller
Product Manager (23)
Herbicide Branch
Registration Division (7505C)

Enclosure

RONSTAR® 1% with FERTILIZER

Label unit 1 of 2

50 lb. Bag Treats up to 11,000 Sq.Ft. of Ornamental Turfgrass

3/4

FOR SALE TO, USE AND STORAGE BY NURSERY, TURF AND LANDSCAPE PERSONNEL ONLY.

FOR USE ON PARKS, GOLF COURSES, ATHLETIC FIELDS, AND OTHER RECREATIONAL AND ORNAMENTAL TURFGRASS AREAS.

Controls: Crabgrass, Annual Bluegrass, Goosegrass, Florida Pusley, Oxalis, Stinging Nettle, Carpetgrass, Pigweed and Other Weeds as Listed.

ACTIVE INGREDIENT:
Oxadiazon; [2-tert-butyl-4-(2,4-dichloro-5-isopropoxyphenyl) A²-1,3,4-oxadiazolin-5-one] 1.0%
OTHER INGREDIENTS: 99.0%
Total 100.0%
Contains RONSTAR® Herbicide a Registered Trade Mark of Aventis, Corp

KEEP OUT OF REACH OF CHILDREN WARNING - AVISO

Si usted no entiende la etiqueta, busque a alguien para que se le explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID

Have this product container or label with you when calling a poison control center, or doctor, or when going for treatment. **IF SWALLOWED** - Call a Poison Control Center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

IF IN EYES - Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.

IF ON SKIN OR CLOTHING - Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice.

IF INHALED - Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth to mouth if possible. Call a poison control center or doctor for further treatment advice. You may also call 1-800-858-7378 for emergency medical treatment advice. See side panel for additional Precautionary Statements. **SEE SIDE/BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS**

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING: Causes eye and skin irritation. Avoid contact with skin, eyes or clothing. Do not allow domestic animals to graze treated areas. Avoid contamination of feed and foodstuffs.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear coveralls over short-sleeved shirt and short pants; waterproof gloves; chemical resistant footwear plus socks; protective eyewear; chemical-resistant headgear for overhead exposure; chemical resistant apron when cleaning equipment. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation. Do not apply this product through any type of irrigation system.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers, and restricted-entry intervals. The requirements in this box only apply to uses of this product (nursery grown ornamental shrubs, vines, trees and container grown ornamentals) that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls over short sleeved shirt and short pants; waterproof gloves, chemical resistant footwear plus socks; protective eyewear; and chemical resistant headgear for overhead exposure.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from areas treated. Do not apply where runoff is likely to occur. Do not contaminate water when disposing of equipment washwaters. Apply only as specified on this label.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

STORAGE: Store product in its original bag, in a cool, dry, locked place out of reach of children. In case of spill, sweep up, and use according to directions or dispose of as follows.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mix, or empty container is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA regional office for guidance.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke.

GUARANTEED FERTILIZER ANALYSIS

Total Nitrogen (N) - Variable %
Nitrate Nitrogen - Variable %
Ammoniacal Nitrogen/Variable %
Water Soluble Nitrogen
and/or Urea Nitrogen/Variable %
Water Insoluble Nitrogen 3% Variable %
Available Phosphoric Acid (P₂O₅) - Variable %
Soluble Potash (K₂O) - Variable %
Derived from Ammonium Sulfate, Urea,
Muriate of Potash.

When applied according to directions this product can be used safely on the following established turfgrass varieties: Bermudagrass, Perennial Bluegrass, Perennial Ryegrass varieties, St. Augustinegrass varieties, Tall Fescue, and Zoysiagrass lawns and ornamental turf. It may be used for Apartment and Condominium complexes, Parks, Golf Courses, Athletic Fields, Cemeteries and all other ornamental and recreational turfgrass areas.

EPA Reg. No. 67508-1 EPA EST. NO. 67508-FL-1

NON-AGRICULTURAL USES REQUIREMENTS

The requirements of this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The WPS applies when this product is used to produce agricultural plants only on farms, forests, nurseries, sod farms or greenhouses.

A PRE-EMERGENT WEED CONTROL PLUS FERTILIZER FOR PERENNIAL BLUEGRASS, PERENNIAL RYEGRASS, ST. AUGUSTINE GRASS, BERMUDAGRASS, TALL FESCUE and ZOYSIAGRASS.

USE PRECAUTIONS

Do not use on Red Fescue, Bentgrass varieties, Dichondra, or Centipede grass because damage will occur. Do not use on Putting greens or "Tees" due to varying cultural practices used.

For St. Augustinegrass application rates above the 50 lbs. per 7,350 sq. ft. will temporarily discolor and damage the turf. If a higher rate is necessary (such as 50 lbs./5,550 sq. ft. or 50 lbs./6,450 sq. ft.) in order to control Goosegrass, Spurge, Poa Annua, Florida Pusley or Carpetgrass, turf damage can be expected to the St. Augustine, therefore apply only when that damage can be tolerated and when good cultural practices are planned in order to revive the damaged turf.

Avoid product contact with flowers and shrubs except as recommended on the Ornamental Use Supplemental Labeling available for this product. For newly seeded lawns do not apply this product until after the second or third cutting. For best lawn care, place new grass seed in the fall of the year. Spring pre-emergence weed control can then be applied. Reseeding treated Areas: Delay new seeding until 4 months after treatment.

CONTROLS GRASSY WEEDS and BROADLEAF WEEDS

Annual Bluegrass (Poa Annua), Barnyard Grass, Crabgrass, Fall Panicum, Florida Pusley, Goose grass, Green Foxtail, Oxalis, Stinging Nettle, Carpetgrass, Pigweed.

HOW TO APPLY

For best weed control rake the area free of leaves and thatch and mow if necessary so that granules will reach the soil surface. Measure the area to be treated and uniformly apply the required amount using a fertilizer spreader (see Spreader Settings chart). Do not apply to turf when it is wet. Do not over-lap spreader applications, on St. Augustinegrass since discoloration "streaking" may temporarily occur from "double rate". After applications if rain is not expected within 24 hours sprinkle the treated turf with 1/4 to 1/2 inch of water to get maximum weed control.

HOW MUCH TO USE

FOR CRABGRASS

Apply 50 lbs./11,000 Sq.Ft. when turf has been in a yearly maintenance program (equal to 2 lbs. A.I. herbicide per acre.) Apply 50 lbs. to treat 7,350 Sq.Ft. of lawn if it has been heavily infested with Crabgrass; equal to 3 lb. A.I. herbicide per acre. Apply in early spring 1 to 2 weeks prior to weed seed germination. This product is a pre-emergence herbicide. It will not control emerged and growing weeds; only germinating weed seeds. Late applications will control only those weed seeds that germinate after the application. See "When To Apply" Chart.

GOOSEGRASS and PROSTRATE SPURGE

These weeds do not usually germinate until late spring or early summer. Apply 50 lbs. to treat 5,555 Sq.Ft. (4 lbs. A.I./acre) and apply 1 to 2 weeks prior to seed germination. See Chart for When To Apply. See Use Precautions for St. Augustine grass

FOR ANNUAL BLUEGRASS (POA ANNUA)

Apply 50 lbs. per 5,555 to 6,450 Sq.Ft. of Lawn; equal to 3.5 to 4 lbs. active herbicide per acre. Apply in late summer to fall or in early spring 1 to 2 weeks before expected weed seed germination. Will not control annual bluegrass present at time of application; therefore only germinating seeds. Refer to chart for proper timing in your locale. Annual BARNYARDGRASS and PROSTRATE SPURGE may also be controlled by this same rate. See Use Precautions for St. Augustine grass

FOR PIGWEED, FLORIDA PUSLEY, OXALIS, STINGING NETTLE AND CARPETGRASS

Where these weeds are a problem apply 50 lbs. to treat 5,555 to 6,450 Sq.Ft. of Lawn; (equal to 3.5 to 4 lbs. Active Herbicide per acre) in early spring or 1 to 2 weeks before expected weed seed germination. Will not control of emerged and growing weeds; only effective against germinating weed seeds. See Use Precautions for St. Augustine grass

SPREADER SETTINGS

Spreader	Amt. of Product	11,000 Sq.Ft. 2#A.I./Acres	7,350 Sq.Ft. 3#A.I./Acres	5,555 Sq.Ft. 4#A.I./Acres
Cyclone	50 lbs.	4 1/2	5 1/2	6 1/2
Scotts #35	50 lbs.	3 1/2	4	4 1/2

NOTE: Spreader rate varies with walking speed.
Check spreader over known area.

When To Apply

JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEPT.	OCT.	NOV.	DEC.
					1	2	3				
					4	2a	3a	4			

- For CRABGRASS and other Grasses and Broadleaf Weeds germinating in the Spring in Northern turf.
 - For GOOSEGRASS and other Late Germinating Annual Grasses and Broadleaf Weeds such as PROSTRATE SPURGE. 2a. GOOSEGRASS in Southern Turf.
 - For POA ANNUA in Northern Turf. 3a. For POA ANNUA in Southern Turf.
 - For Fall-Winter-Spring control of ANNUAL WEEDS in Southern Turf.
- NOTE: Buyer and user assume all risk and liability of use, storage and/or handling of this product not in accordance with the terms of this label. (67508-1-P65 - 10-26-04)

NETWEIGHT 50 POUNDS

GOLF VENTURES, INC.
2101 E. Edgewood Drive - Lakeland, FL 33803

RONSTAR® 1% with FERTILIZER

50 lb. Bag Treats up to 11,000 sq.ft.
FOR WOODY ORNAMENTAL SHRUBS, VINES, TREES & CONTAINER GROWN ORNAMENTALS

4/4

WEED CONTROL AROUND WOODY ORNAMENTAL SHRUBS, VINES & TREES. CONTROLS THE FOLLOWING GRASSY WEEDS and BROADLEAF WEEDS

Annual Bluegrass, Barnyardgrass, Bittercrass, Carpetweed, Carpetgrass, Common Groundsel, Common Purslane, Crabgrass, *Evening Primrose, Green Foxtail, Fall Panicum, Florida Pusley, *Galinsoga, Golden Ragwort, *Goosegrass, Lambsquarter, Oxalis, Pennsylvania Smartweed, Petty Spurge, *Prostrate Spurge, Redroot Pigweed, Spotted Catear, Stinging Nettle, *Swinecress, Yellow Woodsorrel.

Can be applied to both newly transplanted and established ornamentals plants where both weed control and fertilization is desired. Rainfall or overhead irrigation will improve weed control. Do not disturb or cultivate soil after treatment. Do not apply to wet foliage or under conditions in which granules will collect on leaves.

USE PRECAUTIONS

Avoid contact with flowers and shrubs except as recommended on this label.

HOW MUCH TO USE

For susceptible weeds apply 50 lbs. to treat 7,350 to 11,000 sq.ft. of Ornamental planted area. This supplies plants with approximately 0.68 to 1.02 lbs. actual (N) Nitrogen plant food per 1,000 sq.ft. of plant bed. See Chart WHEN TO APPLY. (50 lbs./7,350 sq.ft. is preferred rate.) *For hard to kill weeds as *indicated use the rate of 50 lbs. for 7,350 sq.ft. or 50 lbs. for 5,555 sq.ft. (3 to 4 lbs. A.I./Acre)=6.8 lbs. to 9 lbs. actual product per 1000 sq. ft. of plant bed.

FOR AREAS WHERE WEEDS ARE GROWING

If treatment is desired in non-turf areas where weeds are presently growing first remove existing weeds. Rake and smooth the soil by "breaking-up" clumps. Apply at rate as directed for specific weeds. (50 lbs. to treat either 7,350 or 11,000 sq.ft.) This treatment will only control weeds germinating from seeds. It will not control weeds sprouting from roots! Irrigate the area after treatment with 1/4 to 1/2 inch water to get maximum weed control.

WEED AND FEED FOR CONTAINER GROWN ORNAMENTAL PLANTS

This product is recommended on a wide variety of Container and Field Grown Woody Ornamental Shrubs, Vines and Trees. It may be used on the following list of Container Grown Plants. Use at the rate of 50 lbs. per 5,555 sq.ft. of pot soil surface area. Plants in 14 inch containers set out in 12 foot wide rows X 100 feet long will require 10 3/4 lbs. product per row (Equivalent to 4 lbs. A.I. per acre). Apply with a "shoulder strap held" grass-seeder or granule applicator that will direct the granules to each pot soil surface.

Agapanthus	Cedar	Heath Heather	Oemanthus	Rose
Andromeda	Cotoneaster	Hedera	Periwinkle	Star Jasmine
Arborvita	Cypress	Holly	Photina	Syzygium
Archtoastaphylos	Dogwood	Honeysuckle	Pine-Varieties	Viburnum
Azalea	Douglas Fir	Jasmine	Pittosporum	Xyocma
Barberry	Eucalyptus	Juniper-Varieties	Podocarpus	
Boxwood	Eucynimus	Justica	Privet	
Campan-Japanese	Evergreen	Leucothoe	Pycnantha	
Callistemon	Forsythia	Magnolia	Quercus	
Camellia	Gardenia	Mahonia	Raphiolepis	
Ceanothus	Ginko	Myrtle	Rhododendron	

FOR FIELD GROWN ORNAMENTALS

*Apply at the rate of 50 lbs. per 7,350 or 11,000 sq.ft. of area. This product may be used around the following Established Plants.

Arborvita	Euonymus	Pittosporum	Ground Covers:
Ash	Forsythia	Privet	Carpobrotus
Azalea	Holly	Rhododendron	Osteospermum
Barberry	Honeysuckle	Rose	Sedum
Japanese	Juniper	Russian Olive	Algerian Ivy
Birch	Lilac	Spruce	Vinca Major
Camellia	Maple	Taxus	English Ivy
Cotoneaster	Oak	Viburnum	Pachсандra
Crabapple	Periwinkle	Xyocma	Ajuga
Dogwood	Pine		Euonymus

*Use higher rate (50 lbs./7,350 sq.ft.) for hard to kill weeds. May be used around the following Newly Transplanted Ornamentals.

Camellia	Holly	Pine	Spruce
Dogwood	Juniper	Privet	Taxus
Forsythia	Oak	Rhododendron	Tsuga
Hedera	Periwinkle	Russian Olive	

To obtain continued season long weed control on the following list of plants additional applications may be made 60 days after the previous application in the following situations.

APPLY 2 REPEAT TREATMENTS

50 lbs / 7,350 SQ.FT ON FOLLOWING PLANTS

Established - Container Grown Plants - Abelia, Common Boxwood, English Ivy, Holly, Juniper, Pine, Rhododendron. Newly Transplanted - Container Grown Plants: Common Boxwood, English Ivy, Juniper, Pine, Rhododendron.

APPLY 3 REPEAT TREATMENTS

50 lbs / 7,350 SQ.FT. ON FOLLOWING PLANTS

Established - Container Grown Plants - Abelia, Holly, Rhododendron. Forest Nurseries - For Pines - Loblolly, Slash, Eastern White.

NOTICE: Buyer and user assume all risk and liability of use, storage and/or handling of this product not in accordance with the terms of this label. (Rev. 2-10-26-04)

ACCEPTED
with COMMENTS
In EPA Letter Dated:
NOV 3 0 2004
Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.

67508-1

EPA Reg. No. 67508-1 EPA EST. NO.67508-FL-1
NET WEIGHT 50 POUNDS

GOLF VENTURES, INC.
2101 E. Edgewood Drive - Lakeland, FL 33803