


PM 10 67517-4

FRONT PANEL



INSECTICIDE MIST

- FOR DAIRY BARNS, MILK ROOMS, FOOD PROCESSING PLANTS, MILLS
- AIDS IN CONTROLLING FLIES, MOTHS, CRAWLING INSECTS

ACTIVE INGREDIENTS:

Pyrethrins.....0.1%
Piperonyl butoxide, Technical*.....1.0%

INERT INGREDIENTS:.....98.9%
TOTAL 100.0%

*Consists of 0.8% (butylcarbityl) (6-propylpiperonyl) ether and 0.2% related compounds.

This product contains petroleum distillates

KEEP OUT OF REACH OF CHILDREN

CAUTION

**SEE SIDE/BACK PANEL FOR
ADDITIONAL PRECAUTIONARY STATEMENTS**

EPA Reg. No.: 67517-4
NET CONTENTS: ONE U.S. GALLON (3.785 L)

EPA Est. No.: 67517-MO-1

Manufactured By
PM RESOURCES, INC.
13001 St. Charles Rock Road
Bridgeton, MO. 63044

SIDE/BACK PANEL**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION**

Harmful if Swallowed. Avoid breathing spray mist. Wash thoroughly with soap and water after handling. Do not apply to uncovered human foodstuffs or food products. Avoid contamination of milk and milk equipment. Do not remain in closed area being treated; ventilate area when treatment is completed. Do not allow spray to remain on plastic materials.

STATEMENT OF PRACTICAL TREATMENT

If Swallowed: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. If person is unconscious, do not give anything by mouth and do not induce vomiting.

If in Eyes: Flush eyes with plenty of water. Call physician if irritation persists.

If on Skin: Wash with plenty of soap and water. Get medical attention if irritation persists.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not add directly to water. Do not contaminate water when disposing of equipment washwaters or rinsate.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Dairy Barns and Milk Rooms: Station cows for milking. Close doors and windows to eliminate drafts. Direct mist from sprayer towards ceiling to contact flying insects. When using fixed fogging devices, disperse **Insecticide Mist** horizontally below level of ceiling rafters. Use at the rate of 1/2 ounce for each 1000 cubic feet of space. Allow mist to settle on animals. Keep building closed for 5 minutes after misting. Repeat spraying daily or as needed. If desired, **Insecticide Mist** can also be applied directly to the animals. To control horn flies and for protection against stable flies, mosquitoes, and gnats - apply 1 to 2 ounces per animal once daily, preferably in the morning.

Animal and Poultry Shelters: Use **Insecticide Mist** for killing house flies, stable flies, lesser house flies, mosquitoes and gnats at the rate of 1/2 ounce for each 1000 cubic feet of space. Direct mist towards roof of animal or poultry shelter to contact flying and resting insects. Mist will settle on floor areas killing flies attracted to droppings, litter or animals. Repeat spraying as necessary. **Barns, Stables, and Kennels-** Keep buildings closed for 5 minutes after misting. **Pig Parlors, Loafing Bars,** **Turkey Confinement Porches-** In and about open-type buildings spray **Insecticide Mist** when air is still. **Cage Laying Houses-** Collect eggs before spraying.

SIDE/BACK PANEL

DIRECTIONS FOR USE

For Mills, Elevators, Granaries, Warehouses, Box Cars, Dairies, Meat Packing Companies, Poultry and Food Processing or Storage Plants, Canneries: Use **Insecticide Mist** when mills or food processing plants are not in operation. Close doors and windows, turn off ventilating systems, and extinguish open flames. In poultry and food processing plants - remove exposed products during spraying. Clean premises thoroughly by washing before operations are resumed. Repeat treatment as necessary.

To kill exposed stages of flying insects such as grain, flour, or meal moths, flies, mosquitoes, gnats and wasps- Direct the spray mist towards the upper portions of the enclosure. Use **Insecticide Mist** at the rate of 1/2 ounce for each 1000 cubic feet of space. Keep doors and windows closed for 10 minutes following spraying.

To kill exposed stages of weevils, bran bugs, cadelles, roaches, ants, grain or cheese mites - Use **Insecticide Mist** as a space spray at the rate of 2 ounces for each 1000 cubic feet of space. Also apply as a coarse wet spray into all places commonly inhabited by these insects. Use 1 gallon of **Insecticide Mist** for each 750 square feet of surface. Spray floors, walls and other surfaces of bins, storage and handling areas thoroughly.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage: Do not store near heat or open flame.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke.

NOTICE OF WARRANTY

PM RESOURCES, INC. MAKES NO WARRANTY OF MERCHANTABILITY, FITNESS FOR ANY PARTICULAR PURPOSE, OR OTHERWISE, EXPRESSED OR IMPLIED CONCERNING THIS PRODUCT OR ITS USES WHICH EXTEND BEYOND THE USE OF THE PRODUCT UNDER NORMAL CONDITIONS IN ACCORD WITH THE STATEMENTS MADE ON THIS LABEL.