

294

Reason For Issue:
PM Resources label

Date Issued: 6/14/95
Prev. Date Issued: None

Front Panel

EPA Reg. No. 67517-8

STARLICIDE™
BRAND

COMPLETE

A SLOW ACTING AVICIDE FOR THE CONTROL OF STARLINGS AND BLACKBIRDS AROUND
LIVESTOCK AND POULTRY OPERATIONS.

ACTIVE DRUG INGREDIENT:

3-chloro p-toluidine hydrochloride..... 0.1%

INERT INGREDIENTS..... 99.9%

100.0%

KEEP OUT OF REACH OF CHILDREN

CAUTION:

See Below For Additional Cautions

NET WT 25 LB

G-

Manufactured By
PM Resources, Inc.
13001 St. Charles Rock Road
Bridgeton, MO 63044



How to Use Starlicide Complete

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

The application of Starlicide Complete should be made at the proper time and place under the direction of personnel trained in bird damage control. Normally, the starling's diet consists of insects and fruits; but in winter months when food is in short supply, they're attracted to feedlot areas. Best results are obtained where starlings have few alternate foods. Careful observations of starling habits should be made to establish preferred feeding sites and to determine that no protected bird species are present that may feed on the bait. Prebaiting may be necessary for successful use of the agent.

This product is not effective against English or house sparrows.

For Livestock Feedlots

Place in feeding stations or scatter Starlicide Complete thinly and uniformly over dry or frozen areas of empty pens, alleyways, and other areas inaccessible to livestock. Application should be made before starlings arrive for their first morning meal. One baiting may last up to three days.

The quantity of Starlicide Complete required should not exceed twenty pounds per acre for large lots (over ten acres) nor exceed fifty pounds per acre for small lots (under ten acres).

For Poultry Operations

Place in feeding stations or scatter Starlicide Complete thinly and uniformly over dry or frozen areas on the outside of pens, buildings, or range areas. Application should be made before starlings arrive for their first morning meal. One baiting may last up to three days. Because Starlicide Complete is toxic to chickens, turkeys or other poultry, it should not come in contact with these species.

The quantity of Starlicide Complete required should not exceed twenty pounds per acre for large lots (over ten acres) nor exceed fifty pounds per acre for small lots (under ten acres).


NOTE: Do not expect to find large numbers of dead starlings and/or blackbirds in the baited area as most die within one to three days after consuming the treated pellets. Most die at their roost and along flight lines to the feeding areas. If starlings or blackbirds continue to be a problem, reassess the problem and consider prebaiting. Do not rebait within three days of initial application as some birds consuming a lethal dose of PURINA Starlicide Complete may require three days to be affected.

CAUTION

Handle with protective gloves. Wash hands and equipment used for bait distribution thoroughly with soap and water after handling. Avoid contact with eyes, skin and clothing. If swallowed, induce vomiting and call a physician.

Do not contaminate streams, lakes or ponds with this material.

Starlicide Complete must not be used in any manner that may endanger desirable and protected bird species and domestic fowl. If there is a question of hazard or legality, consult local, state and Federal game authorities before undertaking starling control with Starlicide.



STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

EPA Est. 67517-MO-1

9506 A

Printed in U.S.A.

