

PM-14

Reg # 67517-8

193

FRONT PANEL

STARLICIDE™ COMPLETE

RESTRICTED USE PESTICIDE

DUE TO ACUTE HAZARDS TO EYES, NONTARGET BIRDS AND AQUATIC INVERTEBRATES, AND TO THE NEED FOR HIGHLY SPECIALIZED APPLICATOR TRAINING.

FOR RETAIL SALE TO AND USE ONLY BY CERTIFIED APPLICATORS OR PERSONS UNDER THEIR DIRECT SUPERVISION AND ONLY FOR THOSE USES COVERED BY THE CERTIFIED APPLICATOR' CERTIFICATION.

FOR USE ONLY BY PERSONNEL, SPECIFICALLY TRAINED IN BIRD CONTROL OR PERSONS UNDER THEIR DIRECT SUPERVISION.

A SLOW ACTING AVICIDE FOR THE CONTROL OF STARLINGS AND BLACKBIRDS AROUND LIVESTOCK AND POULTRY OPERATIONS.

ACTIVE INGREDIENT:

3-chloro p-toluidine hydrochloride.....

0.1%

INERT INGREDIENTS:.....


99.9%

TOTAL 100.0%

KEEP OUT OF REACH OF CHILDREN

CAUTION:

SEE SIDE/BACK PANEL FOR
ADDITIONAL PRECAUTIONARY STATEMENTS



EPA Reg. No.: 67517-8
NET WT. 25 LB

EPA Est. No.: 67517-MO-1

Manufactured By
PM RESOURCES, INC.
13001 St. Charles Rock Road
Bridgeton, MO. 63044

293

SIDE/BACK PANEL

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION**

Harmful if swallowed or absorbed through skin. Avoid contact with eyes, skin and clothing. Wear chemical resistant gloves. Wash thoroughly with soap and water after handling.

STATEMENT OF PRACTICAL TREATMENT

If Swallowed: Call a physician or Poison Control Centers. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. If victim is unconscious, do not give anything by mouth and do not induce vomiting.

If On Skin: Wash with plenty of soap and water. Get medical attention if irritation exists.

If In Eyes: Flush eyes with plenty of water. Call a physician if irritation persists.

ENVIRONMENTAL HAZARDS

Starlicide Complete must not be used in any manner that may endanger desirable and protected bird species and domestic fowl. If there is a question of hazard or legality, consult local, state and Federal game authorities before undertaking starling control with Starlicide Complete.

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

The application of Starlicide Complete should be made at the proper time and place under the direction of personnel trained in bird damage control. Normally, the starling's diet consists of insects and fruits; but in winter months when food is in short supply, they're attracted to feedlot areas. Best results are obtained where starlings have few alternate foods. Careful observations of starling habits should be made to establish preferred feeding sites and to determine that no protected bird species are present that may feed on the bait. Prebaiting may be necessary for successful use of the agent.

This product is not effective against English or house sparrows.

SIDE/BACK PANEL

DIRECTIONS FOR USE (Continued)

For Livestock Feedlots

Place in feeding stations or scatter Starlicide Complete thinly and uniformly over dry or frozen areas of empty pens, alleyways, and other areas inaccessible to livestock. Application should be made before starlings arrive for their first morning meal. One baiting may last up to three days.

The quantity of Starlicide Complete required should not exceed twenty pounds per acre for large lots (over ten acres) nor exceed fifty pounds per acre for small lots (under ten acres).

For Poultry Operations

Place in feeding stations or scatter Starlicide Complete thinly and uniformly over dry or frozen areas on the outside of pens, buildings, or range areas. Application should be made before starlings arrive for their first morning meal. One baiting may last up to three days. Because Starlicide Complete is toxic to chickens, turkeys or other poultry, it should not come in contact with these species.

The quantity of Starlicide Complete required should not exceed twenty pounds per acre for large lots (over ten acres) nor exceed fifty pounds per acre for small lots (under ten acres).

NOTE: Do not expect to find large numbers of dead starlings and/or blackbirds in the baited area as most die within one to three days after consuming the treated pellets. Most die at their roost and along flight lines to the feeding areas. If starlings or blackbirds continue to be a problem, reassess the problem and consider rebaiting. Do not rebait within three days of initial application as some birds consuming a lethal dose of Starlicide Complete may require three days to be affected.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

NOTICE OF WARRANTY

PM RESOURCES, INC. MAKES NO WARRANTY OF MERCHANTABILITY, FITNESS FOR ANY PARTICULAR PURPOSE, OR OTHERWISE, EXPRESSED OR IMPLIED CONCERNING THIS PRODUCT OR ITS USES WHICH EXTEND BEYOND THE USE OF THE PRODUCT UNDER NORMAL CONDITIONS IN ACCORD WITH THE STATEMENTS MADE ON THIS LABEL.