FRONT PANEL

DAIRY SPRAY

KILLS HORN FLIES, LICE, HOUSE FLIES, MOSQUITOES, AND GNATS

* Consists of 0.384% of (Butylcarbityl) (6-Propylpiperonyl) Ether and 0.096% of Related Compounds

This product contains petroleum distillates.

KEEP OUT OF THE REACH OF CHILDREN CAUTION SEE SIDE/BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

EPA Req. No.: 67517-12

EPA Est. No.: 67517-MO-1

NET 5 GAL.

NOV 2 9 2000

Under the Federal Insecticide.
Fungicide, and Rodenticide Actno creended, for the posticide

Manufactured By PM RESOURCES, INC 13001 St. Charles Rock Road Bridgeton, MO 63044

(Lot/Code/Product/Rev./Numbers as needed)

Mfg. & Printed in U.S.A.

Precautionary Labeling Must Meet Type Size and Prominence Requirements.

PAGE 1 OF 4 NOVEMBER 13, 2000 PMR12LAB.DOC-1

SIDE/BACK PANEL

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS DANGER

Harmful if swallowed. Avoid contact with skin eyes or clothing. Avoid breathing spray mist. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash clothing before reuse.

Do not wet the skin of animals. Avoid contamination of feed, foodstuffs, milk and milking utensils. Remove animals from barn when fogging or misting.

FIRST AID

If Inhaled ° Move person to fresh air. ° If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice. If On Skin ° Take off contaminated clothing. or Clothing ° Rinse Skin immediately with plenty of water for 15-20 minutes. ° Call a poison control center or doctor for treatment advice. ° Hold eye open and rinse slowly and gently with water for 15-20 If In Eyes minutes. ° Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. If Swallowed ° Call a poison control center or doctor immediately for treatment ° Have person sip a glass of water if able to swallow. On the induce vomiting unless told to do so by a poison control center or doctor. ° Do not give anything by mouth to an unconscious person. HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor for treatment. You may also contact 1-800-XXX-XXXX for emergency medical treatment information.

NOTE TO PHYSICIAN

This product contains petroleum distillate - vomiting may cause aspiration pneumonia.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters or rinsate.

PAGE 2 OF 4 NOVEMBER 13, 2000 PMR04LAB.DOC-2

SIDE/BACK PANEL

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Dairy Spray is a quick knockdown contact spray for killing and repelling horn flies, house flies, mosquitoes and gnats and also kills lice. For fly control, apply daily; after using 3 days the residual repellent and killing action may be sufficient so that applications may be needed only every other day.

Dairy Cattle:

Fly Control: Apply Dairy Spray as a mist to the hair coat. Hold sprayer about 18 inches from the cow; apply mist to entire body from mid-neck region to tail, including the underline and the legs. The amount of spray used should not exceed 1 to 2 ounces per adult animal. Spray calves more sparingly. Do not wet the skin. Wash udder before milking.

Lice Control: Spray the animals with 2 ounces of Dairy Spray. Immediately rub the spray into the hair, paying particular attention to infested areas. Repeat as necessary, but not more often than once every three weeks.

Dairy Barns and Milk Rooms: When used as a space spray contact spray for flies and other insects, remove animals from barn and close doors and windows tightly. Spray Dairy Spray towards ceiling until a light fog hangs in air. Repeat as necessary.

Beef Cattle, Hogs, and Other Farm Animals (Except Horses): - Apply as a mist to the hair coat of farm animals for temporary repellent and killing action. Aids in controlling horn flies, house flies, mosquitoes and gnats. Do not wet the skin.

PAGE 3 OF 4 NOVEMBER 13, 2000 PMR12LAB.DOC-3

SIDE/BACK PANEL

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Storage: Do not pour, spill, use or store near heat or open flame.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on or at an approved waste disposal facility.

Container Disposal (Plastic): Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities.

NOTICE OF WARRANTY

PM RESOURCES, INC. MAKES NO WARRANTY OF MERCHANTABILITY, FITNESS FOR ANY PARTICULAR PURPOSE, OR OTHERWISE, EXPRESSED OR IMPLIED CONCERNING THIS PRODUCT OR ITS USES WHICH EXTEND BEYOND THE USE OF THE PRODUCT UNDER NORMAL CONDITIONS IN ACCORD WITH THE STATEMENTS MADE ON THIS LABEL.

PAGE 4 OF 4 NOVEMBER 13, 2000 PMR12LAB.DOC-4