



067517-00021-070396

Printed: 12:49:10 Friday, 21 May, 1999 # 14 / 2880

Systems Integration Group, Inc.

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

Mr. Joseph E. Dyer
PM Resources, Inc.
13001 St. Charles Rock Road
Bridgeton MO 63044

03 JUL 1996

Dear Mr. Dyer:

Subject: Amendment - Addition of Swine Claim
Purina Cattle Duster Insecticide
EPA Registration Number 67517-21
Your Application Dated March 4, 1996

310/52795
17/

The amendment referred to above, submitted in connection with registration under FIFRA sec. 3(c)(7)(A), is acceptable provided that you:

1. Submit and/or cite all data or other material required for registration/reregistration of your product under FIFRA sec. 3(c)(5) or FIFRA sec. 4 when the Agency requires all registrants of similar products to submit such data.

2. Make the labeling changes listed below before you release the product for shipment:

- a. Add the new site, swine, to the front label panel.
- b. Under the Restrictions section, add the following statement:

Do not apply to sick, stressed or convalescent animals.
- c. Under the direct treatment and bedding treatment of swine, add the following statement:

Treat no more than 6 times per season.
- d. Under Swine bedding treatment, delete the second "often than" from the statement "Repeat as necessary.... etc."
- e. Under NOTE:, revise the statement "No withdrawal necessary prior to slaughter." to read "No interval is required between treatment and slaughter or between treatment and use of milk as feed."

3. Submit two copies of your final printed labeling before you release the product for shipment. Refer to the A-79 Enclosure for a further description of final printed labeling.

CONCURRENCES							
SYMBOL	UCA						
SURNAME							
DATE							

2

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Sincerely yours,

George T. LaRocca
Product Manager 13
Insecticide-Rodenticide Branch
Registration Division (7505C)

Enclosure

FRONT PANEL

CATTLE DUSTER INSECTICIDE

- For dairy or beef cattle *in stables*
- For control of horn flies and lice
- Aids in reducing face flies on cattle

ACCEPTED
with COMMENTS
in EPA Letter Dated

03 JUL 1996

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.
67517-21

ACTIVE INGREDIENT:

0,0-Diethyl 0-(3-chloro-4-methyl-2-oxo-2H-
1-benzopyran-7-yl) phosphorothioate..... 1.0%

INERT INGREDIENTS 99.0%
TOTAL 100.0%

KEEP OUT OF REACH OF CHILDREN

CAUTION

**SEE SIDE/BACK PANEL FOR
ADDITIONAL PRECAUTIONARY STATEMENT**

EPA Reg. No.: 67517-21
NET WT. 12 ½ LBS.

EPA Est. No.: 67517-MO-1

Manufactured by
PM RESOURCES, INC.
13001 St. Charles Rock road
Bridgeton, MO. 63044

SIDE/BACK PANEL

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION**

Harmful if swallowed, inhaled or absorbed through skin. Avoid breathing dusts. Avoid contact with eyes, skin, and clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash clothing before reuse.

STATEMENT OF PRACTICAL TREATMENT

If Swallowed: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching the back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

If Inhaled: Remove victim to fresh air. If not breathing give artificial respiration preferably mouth-to-mouth. Get medical attention.

If On Skin: Wash with plenty of soap and water. Get medical attention if irritation persists.

If In Eyes: Flush eyes with plenty of water. Call a physician if irritation persists.

NOTE TO PHYSICIAN: Atropine sulfate by injection is antidotal. 2-PAM is also antidotal and may be administered in conjunction with atropine.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to birds, fish, and aquatic invertebrates. Do not apply directly to water. Do not contaminate water when disposing of equipment, washwater or rinsate.

RESTRICTIONS

For external, insecticidal use on above specified animals only. Avoid contamination of feed, troughs, water and water utensils. Provide thorough ventilation while dusting.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling

TO CONTROL HORNFLIES AND LICE ON BEEF AND DAIRY CATTLE:

A ready-to-use insecticide dusting powder for use in the refillable Cattle Duster Bag – a weather-proof bag for use in open or sheltered areas. This product can also be used for hand dusting. Effective in controlling horn flies and lice; aids in controlling face flies. No withdrawal period required; may be used with dairy cattle in production or beef cattle being fed for slaughter.

SIDE/BACK PANEL

DIRECTIONS FOR USE (Continued)

DUST BAG

Empty contents of bag (Cattle Duster Insecticide) into the Cattle Duster Bag through opening along edge of upper side. Close opening. Use 1 lb. of Cattle Duster Insecticide per 8 animals every two weeks. Do not exceed 3 lbs. of Cattle Duster Insecticide per animal per year.

HOW TO HANG:

Hang Cattle Duster Bag by means of rope or chain placed through straps along top edge. For Horn Fly and Lice control adjust height so that bottom of bag will hang a little below the backline of the animals using it. To aid in reducing face flies hang bag so that animals will be forced to dust face daily.

WHERE TO HANG:

Suspend Cattle Duster Bag in locations where animals will frequently dust themselves on a free-choice basis; or in gateways or lanes through which animals are forced to pass daily for water, feed or mineral.

Dairy Cattle: For lactating dairy cows, hang in the exit of the milking barn.

Restrictions: Do not hang Cattle Duster Bag over feed, mineral or water troughs.

HAND DUSTING

Apply not more than 2 oz. (6 level tablespoonfuls) evenly into the hair over the head, neck, shoulders, back and tailhead. Repeat as necessary, but not more often than every 14 days. (Maximum Seasonal Application Rate: 26 applications/year).

TO CONTROL LICE ON SWINE:

Apply not more than 1 oz. (3 level tablespoonfuls) per head as a uniform coat to the head, shoulders, and back by use of a shaker can or suitable mechanical dust applicator. Repeat as necessary but not more often than once every ten days. *Treat no more than 6 times per season*

Bedding Treatment: Apply 2 oz. (6 level tablespoonfuls) uniformly over each 30 square feet of fresh, dry bedding by use of a shaker can or a suitable mechanical dust applicator. Repeat as necessary but not more often than ~~often than~~ once every ten days. *Treat no more than 6 times per season*

NOTE: In severe infestations both individual animals and the bedding may be treated as directed above. No withdrawal necessary prior to slaughter. *Never*

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

SIDE/BACK PANEL

STORAGE AND DISPOSAL (Continued)

Container Disposal: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

NOTICE OF WARRANTY

PM RESOURCES, INC. MAKES NO WARRANTY OF MERCHANTABILITY, FITNESS FOR ANY PARTICULAR PURPOSE, OR OTHERWISE, EXPRESSED OR IMPLIED CONCERNING THIS PRODUCT OR ITS USES WHICH EXTEND BEYOND THE USE OF THE PRODUCT UNDER NORMAL CONDITIONS IN ACCORD WITH THE STATEMENTS MADE ON THIS LABEL.