

**PRECAUTIONARY STATEMENTS
HAZARD TO HUMANS & DOMESTIC ANIMALS**

CAUTION

Keep away from humans, domestic animals and pets. If swallowed this material may reduce the clotting ability of the blood and cause bleeding.

Note To Physician: If ingested, administer Vitamin K, intramuscularly or orally as indicated in bishydroxycoumarin overdoses. Repeat as necessary based on monitoring of prothrombin times.

ENVIRONMENTAL HAZARDS

This product is toxic to mammals and birds. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

READ THIS LABEL: Read this entire label and follow all use directions and use precautions.

IMPORTANT: Do not expose children, pets, or other nontarget animals to rodenticides. To help to prevent accidents:

1. Store product not in use in a location out of reach of children and pets.
2. Apply bait in locations out of reach of children, pets, domestic animals and nontarget wildlife, or in tamper-resistant bait stations. These stations must be resistant to destruction by dogs and by children under six years of age, and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from stations when they are lifted, units must be secured or otherwise immobilized. Even stronger bait stations are needed in areas open to hooded live-stock, raccoons, bears, or other potentially destructive animals, or in areas prone to vandalism.
3. Dispose of product container, and unused, spoiled, and unconsumed bait as specified on this label.

Use Restrictions: For control of Norway rats, Roof rats and House mice in and around homes, industrial buildings, and similar man-made structures. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food. Do not broadcast bait.

Selection of Treatment Areas: Determine areas where rats or mice will most likely find and consume the bait. Generally, these are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents and/or their signs have been seen. Protect bait from rain or snow. Remove as much alternative food as possible.

For Rats (Roof and Norway): Apply one to four 4-oz. placepacks per placement. Maintain an uninterrupted supply of fresh bait for at least 10 days.

For House Mice: Open placepack and apply 1/4 to 1/2 ounce (1-2 level tablespoons) of bait at 8 to 12 foot intervals. Larger placements (up to 2 ounces) may be needed at points of very high mouse activity. Maintain an uninterrupted supply of fresh bait for at least 15 days.

Rats/Mice: Replace contaminated or spoiled bait immediately. Collect and dispose of all dead animals and leftover bait properly. To prevent reinfestation, limit sources of rodent food, water, and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present establish permanent bait stations and replenish bait as needed.

STORAGE AND DISPOSAL

Storage: Store only in original container, in a dry place inaccessible to children and pets.

Disposal: Do not reuse empty container. Securely wrap in newspaper and discard in trash.

EPA Reg. No. 67517-27

PM Resources

PLACE PACK® RAT CONTROL

Kills Norway Rats, Roof Rats, and House Mice

ACCEPTED

DEC 1 6 1995

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 67517-27.

ACTIVE INGREDIENT:

Warfarin, 3-(alpha-Acetylbenzyl)-4-hydroxycoumarin	0.025%
INERT INGREDIENTS	99.975%
TOTAL	100.000%

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE RIGHT PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

Manufactured by
PM RESOURCES, INC.
13001 St. Charles Rock Road
Bridgeton, MO 63044

**NET WT 15 LB.
(60 4-OZ. PACKS)**

9511 K

EPA Est. 67517-MO-1

Printed in U.S.A.

D-8329

Reduced to 77% to fit on
8 1/2" x 11"

PM 14 67517-27
PS 133

PM RESOURCES

PLACE PACK RAT CONTROL



Kills Norway Rats, Roof Rats and House Mice

ACTIVE INGREDIENT:

Warfarin, 3-(alpha-Acetylbenzyl)-4-hydroxycoumarin

0.025%

INERT INGREDIENTS

99.975%
100.000%

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE BELOW FOR PRECAUTIONARY STATEMENTS

PRECAUTIONARY STATEMENTS
HAZARD TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Keep away from humans, domestic animals and pets. If swallowed this material may reduce the clotting ability of the blood and cause bleeding.

NOTE TO PHYSICIAN: If ingested, administer Vitamin K intramuscularly or orally as indicated in bishydroxycoumarin overdoses. Repeat as necessary based on monitoring of prothrombin times.

ENVIRONMENTAL HAZARDS: This product is toxic to mammals and birds. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark.

NET WT 4 OZ

MANUFACTURED BY

PM RESOURCES, INC.

13001 ST. CHARLES ROCK ROAD

BRIDGETON, MO 63044

BEST COPY AVAILABLE

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

READ THIS LABEL: Read this entire label and follow all use directions and use precautions.

IMPORTANT: Do not expose children, pets, or other nontarget animals to rodenticides. To help to prevent accidents:

1. Store product not in use in a location out of reach of children and pets.
2. Apply bait in locations out of reach of children, pets, domestic animals and nontarget wildlife, or in tamper-resistant bait stations. These stations must be resistant to destruction by dogs and by children under six years of age, and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from stations when they are lifted, units must be secured or otherwise immobilized. Even stronger bait stations are needed in areas open to hoofed livestock, raccoons, bears, or other potentially destructive animals, or in areas prone to vandalism.
3. Dispose of product container, and unused, spoiled, and unconsumed bait as specified on this label.

USE RESTRICTIONS: For control of Norway rats, Roof rats and House mice in and around homes, industrial buildings, and similar man-made structures. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food. Do not broadcast bait.

SELECTION OF TREATMENT AREAS: Determine areas where rats or mice will most likely find and consume the bait. Generally, these are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents and/or their signs have been seen. Protect bait from rain or snow. Remove as much alternative food as possible.

FOR RATS (Roof and Norway): Apply one to four 4-oz. placepacks of bait per placement. Maintain an uninterrupted supply of fresh bait for at least 10 days.

FOR HOUSE MICE: Open placepack and apply 1/4 to 1/2 ounce (1-2 level tablespoons) of bait at 8 to 12 foot intervals. Larger placements (up to 2 ounces) may be needed at points of very high mouse activity. Maintain an uninterrupted supply of fresh bait for at least 15 days.

RATS/MICE: Replace contaminated or spoiled bait immediately. Collect and dispose of all dead animals and leftover bait properly. To prevent reinfestation, limit sources of rodent food, water, and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present establish permanent bait stations and replenish bait as needed.

STORAGE AND DISPOSAL

STORAGE: Store only in original container, in a dry place inaccessible to children and pets.

DISPOSAL: Do not reuse empty container. Securely wrap in newspaper and discard in trash.

EPA Est. 67517-MO-1

D-8324

9411 K

E.P.A. Reg. No.
67517-27

Printed in U.S.A.

BEST COPY AVAILABLE