

FRONT PANEL
RABON DUST FOR LIVESTOCK AND POULTRY

Contains 3% Rabon®* Insecticide

For Use On Livestock and Poultry

No Milk or Tissue Withdrawal

A Ready-To-Use Insecticide Dust

On Livestock:

Controls Horn Flies Cattle Grubs, Lice and aids in the control of Face Flies on Beef and Dairy Cattle. Controls Lice on Swine.

On Poultry:

Controls Northern Fowl Mites, Chicken Mites, and Lice on or around Chickens. Controls Darkling Beetle (litter beetle) in litter.

ACTIVE INGREDIENT:

Tetrachlorvinphos (CAS # 961-11-5):3.00%

OTHER INGREDIENTS:.....97.00%

TOTAL 100.00%

* Rabon Insecticide

Rabon is a registered trademark of Shell Oil Company

KEEP OUT OF REACH OF CHILDREN

CAUTION

**SEE SIDE/BACK PANEL FOR
ADDITIONAL PRECAUTIONARY STATEMENTS**

PRECAUCION

Al Usuario: Si usted no lee ingles, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.
(ENGLISH TRANSLATION)

CAUTION

To the User: If you cannot read English, do not use this product until the label has been fully explained to you.

EPA Reg. No.: 67517-40

EPA Est. No.: 67517-MO-1

NET WT. 1 LB (454 g)

Manufactured By
PM Resources, Inc.
13001 St. Charles Rock Road
Bridgeton, MO. 63044

IN CASE OF EMERGENCY CALL: NATIONAL PESTICIDE TELECOMMUNICATIONS NETWORK 1-800-888-7378.

ACCEPTED

SEP 9 1999

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. **67517-40**

SIDE/BACK PANEL

**PRECAUTIONARY STATEMENTS
HAZARD TO HUMANS AND DOMESTIC ANIMALS
CAUTION**

HAZARD TO HUMANS: Harmful if swallowed or absorbed through the skin. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Avoid contamination of food and feedstuffs. This product may cause skin sensitization reactions in certain individuals. Wear long-sleeved shirt and pants, chemical resistant gloves, shoes and socks. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Remove PPE immediately after handling this product. Wash the outside of gloves.

FIRST AID

If Swallowed: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching the back of the throat with finger, or if available by administering syrup of ipecac. If person is unconscious, do not give anything by mouth and do not induce vomiting.

If In Eyes: Flush eyes with plenty of water. Call a physician if irritation persists.

If On Skin: Wash with plenty of soap and water. Get medical attention if irritation persists.

NOTE TO PHYSICIAN/VETERINARIAN: Rabon is an organophosphate insecticide. If symptoms of cholinesterase inhibition are present, atropine sulfate by injection is antidotal. 2-PAM is also antidotal and may be administered, but only in conjunction with atropine.

Systems of Exposure: May produce acute cholinesterase depression. Acute cholinesterase depression may be evidenced by headache, nausea, vomiting, diarrhea, abdominal cramps, excessive sweating, salivation and tearing, constricted pupils, blurred vision, tightness in chest, weakness, muscle twitching, and confusion; in extreme cases, unconsciousness, convulsions, and severe respiratory depression may occur.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Drift and runoff may be hazardous to aquatic organisms in adjacent areas. Do not apply directly to water. Do not contaminate water when disposing of equipment washwater. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

SIDE/BACK PANEL

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL BEFORE EACH USE.

APPLICATION RESTRICTION: Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

FOR USE ON POULTRY

TYPE OF HOUSING	APPLICATION	REMARKS
Wire Cages	1 lb/300 birds Plunger, rotary, type duster or shaker-can duster.	For individual treatment direct dust to vent and fluff area. Group treatment may be preferred. Dust should reach skin. Do not repeat more often than once every 14 days. Wire rungs & corners should also be treated.
NOTE: For northern fowl mites on roosters, thorough individual application of the dust will assure long-lasting control and reduce reinfestation of breeding flocks.		
Floor Management Litter	1 lb/100 sq. ft. Plunger or rotary type duster.	Treat evenly and thoroughly. Also treat roosts, cracks and crevices where pests may hide.
Dust Box	2 lb/100 birds.	Use box about 2 ft. by 2 ft. by 1 ft. deep.
Roost Paint	1 lb/100 sq. ft. Make thick slurry by mixing 1 lb. of dust with 1 pint of water, continually stir.	Treat roosts thoroughly, particularly the cracks and crevices.

SIDE/BACK PANEL**DIRECTIONS FOR USE (Continued)****FOR USE ON LIVESTOCK**

Rabon Dust is a ready-to-use insecticide dust which can be applied by hand or through the use of self-treating dust bags. There is no withholding period from last application to slaughter.

Animal	Insect	Remarks
Beef Cattle & Dairy Cattle	Horn Flies, Face Flies & Lice	<p>HAND DUSTING - Apply approximately 2 oz. of dust by shaking can, rotary duster or by spoon to the upper portions of the back as an aid for the control of face flies. Rub in lightly to carry the dust beneath the hair. Do not treat more often than once every two weeks.</p> <p>SELF-TREATING DUST BAG FORCED USE - Put Rabon Dust in cotton cloth or double burlap bags or use weather-proof cattle dust bags and hang in door exits or alleyways leading from animal buildings, salt or mineral blocks, or watering holes. Protect cloth or burlap bags from weather.</p> <p>FREE-CHOICE USE - Use the same dust bags as above but place in loafing sheds, holding pens, feedlots, near watering holes or other areas where cattle gather. The free choice aids in the control of lice.</p>
Beef Cattle & Dairy Cattle	Cattle Grubs	<p>After the grubs have encysted (formed warbles) apply 3-4 oz. down the backline and rub in thoroughly, taking care to get the dust into the warble if possible.</p>

SIDE/BACK PANEL

DIRECTIONS FOR USE (Continued)

<u>Animal</u>	<u>Insect</u>	<u>Remarks</u>
Swine	Lice	HAND DUSTING - Apply 3-4 oz. of dust by conventional hand or power duster to each animal with special attention given to the neck and around the ears. Do not repeat more often than once every 14 days. In severe infestations, both individual animals and bedding may be treated. One lb. of Rabon Dust should be applied per 50 sq. ft. of bedding.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store in a dry place in original container.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on-site or at an approved waste disposal facility.

Container Disposal: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

NOTICE OF WARRANTY

PM RESOURCES, INC. MAKES NO WARRANTY OF MERCHANTABILITY, FITNESS FOR ANY PARTICULAR PURPOSE, OR OTHERWISE, EXPRESSED OR IMPLIED CONCERNING THIS PRODUCT OR ITS USES WHICH EXTEND BEYOND THE USE OF THE PRODUCT UNDER NORMAL CONDITIONS IN ACCORD WITH THE STATEMENT MADE ON THIS LABEL.