PM 03 67517-43 8/4/49 page 185

FRONT PANEL

FLY-A-REST® AQ

PROFESSIONAL STRENGTH - LOW DOSAGE REQUIRED

For use in Schools, Hospitals, Institutions, Supermarkets, Dairies, Hotels, Restaurants, Drive-Ins, Motels, Drug Stores, Food Plants, Milk Rooms and Animal Quarters.

Provides rapid control of house flies, mosquitoes, small flying moths (millers), gnats, wasps and accessible stages of cockroaches, ants, water bugs, silverfish, flour beetles, weevils, grains mites and spiders.

Kills and repels stable flies, horse flies, face flies, deer flies, house flies, horn flies, mosquitoes and gnats.

ACTIVE INGREDIENTS:

Pyrethrins	0.5%
Piperonyl Butoxide, Technical	•
OTHER INGREDIENTS:	
	TOTAL 100.0%

Equivalent to 4.0% (butylcarbityl) (6-propylpiperonyl) ether and to 1% related compounds.

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE SIDE/BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

ACCEPTED

AUG 4 1989

Under the Federal Invecticide.

Suppliede, and Redepticitée Act.

Augusticide, and redepticitée Act.

Augusticide augusticide

The Bog. No. 67517-43

EPA Reg. No.: 67517-43

NET 24 fl. oz.

EPA Est. No.: 67517-MO-1

Manufactured By
PM RESOURCES, INC.
13001 St. Charles Rock Road
Bridgeton, MO. 63044

Page 1 of 5 June 28, 1999 PMR43LAB.DOC-1

SIDE/BACK PANEL

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if swallowed inhaled or absorbed through skin. Avoid breathing vapors or spray mist. Avoid contact with skin and eyes. Wash thoroughly with soap and water after handling.

Do not apply directly to food. In commercial food handling facilities, cover or remove any food and food processing equipment during application. Do not apply while food processing is under way. After space spraying in food processing plants, wash all equipment, benches, shelving, etc., where exposed food will be handled, with potable water. All food processing surfaces and utensils should be covered during treatment or thoroughly washed before use.

Remove pets, birds and cover fish aquariums before room applications are made.

STATEMENT OF PRACTICAL TREATMENT

If Swallowed: Call a physician or Poison Control Centers. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. If victim is unconscious, do not give anything by mouth and do not induce vomiting.

If In Eyes: Flush eyes with plenty of water. Call a physician if irritation persist.

If On Skin: Wash with plenty of soap and water. Get medical attention if irritation exists.

If Inhaled: Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters or rinsate.

PHYSICAL OR CHEMICAL HAZARDS

Contents under pressure. Do not use or store near heat or open flame. Do not puncture or incinerate container. Exposure to temperatures above 130° may cause bursting.

SIDE/BACK PANEL

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

SHAKE WELL BEFORE USE

Hold container upright with nozzle away from you. Press valve down and spray as directed. To avoid possible staining of fabrics or surfaces, do not spray at close range.

House Flies, Mosquitoes, Gnats, Wasps and Small Flying Moths (Millers): Close all doors and windows and spray room at the rate of 2 seconds per 1000 cubic feet. Direct spray to all parts of the room especially windows and other light sources which attract these insects. Keep nozzle in constant motion. Keep room closed for 10-15 minutes after treatment. Repeat as necessary.

Cockroaches, Waterbugs, Silverfish, Crickets, Spiders, Centipedes and Snow Bugs: Spray into hiding places such as cracks and crevices, behind sinks, cabinets, along baseboards and floors, around drains and plumbing, hitting insects with spray whenever possible. Repeat treatment as necessary.

Ants: Spray trails, nests and points of entry. Spray on ants where possible. Repeat as necessary.

Fleas: Spray sleeping quarters, bedding, floor and floor covering where pets are kept. Repeat as necessary. Use only a recommended pet spray on pets.

Carpet Beetles: Spray directly on beetles, baseboards, floors, shelves and carpets. Repeat treatment as necessary.

Bedbugs: Spray mattresses, particularly around tufts and seams. Take beds apart and spray into all joints. Treat baseboards, molding and floors. Allow treated fabric to dry thoroughly before using. Repeat treatment as necessary.

In Animal Quarter and Mild Rooms: For rapid control of house flies, mosquitoes, gnats, wasps and small flying moths, close windows and doors and spray at the rate of 2 seconds per 1000 cubic feet of space. For best results, keep area closed for 10 minutes after spraying. Do not remain in treated area and ventilate after treatment is completed.

To Treat Animals: To protect beef and dairy cattle from attacks of stable flies, horse flies, deer flies, house flies, horn flies, mosquitoes and gnats, spray bout 3 seconds on each side being careful to spray back, withers and forelegs thoroughly. To protect from face flies, spray the face and head, but do not spray into the eyes. Repeat treatment as necessary when flies are troublesome.

SIDE/BACK PANEL

DIRECTIONS FOR USE (Continued)

Exposed Stages of Indian Moths and Angoumois Grain Moths: Close doors and window. Directly spray into all parts of room, especially around stored food product containers, pallets and darkened areas. Spray 7 to 10 seconds for each 1000 cubic feet of space. Keep area closed for 15 minutes following treatment. Repeat application weekly or as needed. Infested stored products should be fumigated or treated by other effective methods using an approved product intended for this purpose.

Exposed Stages of Flour Beetles, Weevils and Grain Mites: Spray into cracks and crevices, processing and handling equipment, around cartons or containers and other areas where these insects tend to congregate. Spray 30 seconds for each 100 square feet of surface. Keep area closed for 45 minutes to 1 hour after treatment. Repeat applications as necessary. Infested stored food products in the home should be disposed of.

Controls Garden Insects on Ornamental and Vegetable Plants: Spray with slow sweeping motion, not closer that 18 inches from plant. Apply in accordance with the chart below. Avoid wetting plants. Spray plants only when air is still. For single plants, spray in short bursts for 1-2 seconds. Repeat as necessary for control. Be cautious about wetting tender foliage, young plants and new growth. Do not spray plants when temperatures exceed 90° F. Contact insects directly whenever possible. Repeat application of 7 to 10 day intervals or after heavy rains.

Acceptable for use to day of harvest on edible plants.

APPLICATION

USE ON	SECONDS OF SPRAY / FOOT OF ROW OR PER PLANT	TO CONTROL
Roses	2-3	Aphids, Roses Chafers, Japanese Beetles, Rose Slugs, Tent Caterpillars, Scales
Asters	2-3	Leafhoppers, Tarnish Plant Bugs, Lace Bugs, Thrips (Exposed) Mealy Bugs.
Dahlias	2-3	Aphids, Red Spider Mites, Plant Bugs.
African Violets	1	Broad Mites, Citrus Mealy Bugs.
Azaleas	1-2	Azalea Lace Bugs, Whiteflies, Leafminers.
Euonymus	1-2	Euonymus Scales, Aphids, Thrips (Exposed).

Page 4 of 5 June 28, 1999 PMR43LAB.DOC-4

5 45

SIDE/BACK PANEL APPLICATION (Continued)

	SECONDS OF SPRAY / FOOT	
<u>USE ON</u>	OF ROW OR PER PLANT	TO CONTROL
Hydrangeas	1-2	Aphids Red Spider Mites, Hydrangea
		Leaftiers
Junipers	3-5	Juniper Scales, Red Spider Mites,
, ·		Bagworms.
Lilacs	3-5	Leafminers, Leafrollers, Scales.
Tomatoes	2-4	Flea Beetles, Hornworms, Stink Bugs,
		Colorado Potato Beetles, Green Peach Aphids.
Peppers, Eggplants	1-2	Colorado Potato Beetles, Leafhoppers,
		Potatoes Blister Beetles, Flea Beetles, Green Peach Aphids.
Cucumber, Squash,	2-4	Striped and Spotted Cucumber Beetles.
Melons		
Beans	1-2	Mexican Bean Beetles, Whiteflies, 12-
		Spotted Beetles.
Sweet Corn	2-3	Japanese Beetles.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store in a cool, dry area away from heat or open flame.

Pesticide Disposal: Securely wrap container in several layers of newspaper and discard in trash.

Container Disposal: Replace cap and discard container in trash. Do not incinerate or puncture.

NOTICE OF WARRANTY

PM RESOURCES, INC. MAKES NO WARRANTY OF MERCHANTABILITY, FITNESS FOR ANY PARTICULAR PURPOSE, OR OTHERWISE, EXPRESSED OR IMPLIED CONCERNING THIS PRODUCT OR ITS USES WHICH EXTEND BEYOND THE USE OF THE PRODUCT UNDER NORMAL CONDITIONS IN ACCORD WITH THE STATEMENTS MADE ON THIS LABEL.

Page 5 of 5 June 28, 1999 PMR43LAB.DOC-5