

FRONT PANEL

HARD HITTER® CATTLE POUR-ON INSECTICIDE

- Controls Lice and Flies on Dairy and Beef Cattle
- Controls Keds and Lice on Sheep

ACTIVE INGREDIENT:

Permethrin (3-Phenoxyphenyl) methyl (±) cis, trans-3-(2,2-dichloroethenyl)-2,2-dimethylcyclopropanecarboxylate* 1.0%

OTHER INGREDIENTS: 99.0%

TOTAL 100.0%

Contains petroleum distillates

*Cis/trans isomer ratio: Min. 35% (±) cis and max. 65% (±) trans

KEEP OUT OF REACH OF CHILDREN

CAUTION

**SEE SIDE/BACK PANEL FOR
ADDITIONAL PRECAUTIONARY STATEMENTS**

EPA Reg. No.: 67517-44

EPA Est. No.: 67517-MO-1

NET CONTENTS: ONE U.S. GALLON (3.785 L)

IN CASE OF EMERGENCY CALL: CHEMTREC (POISON CONTROL CENTER)
AT 1-800-424-9300

Manufactured By
PM Resources, Inc.
 13001 St. Charles Rock Road
 Bridgeton, MO. 63044

ACCEPTED
 APR 15 1998
 Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 67517-44

SIDE/BACK PANEL

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS
CAUTION**

Avoid contact with eyes. Harmful if swallowed. Wash thoroughly with soap and water after handling.

STATEMENT OF PRACTICAL TREATMENT

If In Eyes: Flush eyes with plenty of water. Call a physician if irritation persists.

If Swallowed: Call a doctor or get medical attention. Do not induce vomiting. Do not give anything by mouth to an unconscious person. Avoid Alcohol.

Note To Physician: Solvent presents aspiration hazard. Gastric lavage is indicated if material was taken internally.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to fish. Use with care when applying to areas adjacent to any body of water. Do not add directly to water. Do not contaminate water by disposal of equipment washwaters. Apply this product only as specified on this label.


PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READY TO USE - No dilution necessary.



SIDE/BACK PANELAPPLICATION

<u>APPLY TO</u>	<u>TARGET INSECTS</u>	<u>APPLICATION INSTRUCTIONS</u>
Lactating and Non-Lactating Dairy Cattle and Beef Cattle and Calves	LICE Horn flies, Face flies AND Aids in Control of: Horse Flies, Stable Flies, Mosquitos, Blackflies and Ticks.	Beginning at the head, pour along the center of the back. Apply ½ fluid ounces per 100 pounds body weight of animal up to a maximum of 5 fluid ounces for any one animal.
Sheep	SHEEP KEDS Lice	Pour along back. Apply ¼ fl. oz (7.5 cc) per 50 lbs body wt of animal, up to a maximum of 3 fl. oz for any one animal.

For cattle and sheep, repeat treatment as needed, but not more than once every 2 weeks. For optimum lice control, two treatments at 14-day intervals is recommended.

Special Note: **HARD HITTER** Cattle Pour-On Insecticide is not effective in controlling cattle grubs.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage: Keep container sealed when not in use. Do not store near food or feed.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Triple rinse (or equivalent). Then offer for recycling or reconditioning or puncture and dispose of in a sanitary landfill or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

NOTICE OF WARRANTY

PM RESOURCES, INC. MAKES NO WARRANTY OF MERCHANTABILITY, FITNESS FOR ANY PARTICULAR PURPOSE, OR OTHERWISE, EXPRESSED OR IMPLIED CONCERNING THIS PRODUCT OR ITS USES WHICH EXTEND BEYOND THE USE OF THE PRODUCT, UNDER NORMAL CONDITIONS IN ACCORD WITH THE STATEMENTS MADE ON THIS LABEL.