

67517-55

12/07/2000  
FRONT PANEL

1/5

**PERMETHRIN WATER-BASE SPRAY**

**A Concentrated Permethrin Stable Emulsion**

To protect cattle (beef and dairy), goats, sheep, hogs, and horses from horn flies, face flies, house flies, mosquitoes and gnats.

For use outdoors and in homes, kennels and non-food/feed areas, warehouses, theaters, office buildings, schools, motels, hotels, and food/feed processing establishments.

For use on dogs to control fleas, ticks and lice.

**ACTIVE INGREDIENT:**

\*Permethrin (CAS#52645-53-1) .....0.50%

**OTHER INGREDIENTS:**.....99.50%

TOTAL 100.00%

\*Cis/trans isomer ratio: Min. 35% (+/-) cis and max. 65% (+/-) trans.

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**


**SEE SIDE/BACK PANEL FOR  
ADDITIONAL PRECAUTIONARY STATEMENTS**

EPA Reg. No.: 67517-55  
NET CONTENTS: 16 fl. oz. (473 ml)

EPA Est. No.: 67517-MO-1

Manufactured By  
**PM Resources, Inc.**  
13001 St. Charles Rock Road  
Bridgeton, MO. 63044

**ACCEPTED**  
DEC 7 2000  
Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
as amended, for the pesticide  
registered under  
EPA Reg. No. 67517-55



SIDE/BACK PANEL

**READ ENTIRE LABEL BEFORE EACH USE**

**DO NOT USE ON CATS**

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS & DOMESTIC ANIMALS  
CAUTION**

**HAZARDS TO HUMANS:**

Harmful if swallowed. May be absorbed through skin. Avoid inhalation of spray mist. Avoid contact with skin, eyes, or clothing. Wash thoroughly after handling and before eating or using tobacco. Avoid contamination of feed and foodstuffs. Do not use on humans. Do not allow children or pets to contact areas until surfaces are dry. Do not apply in classrooms when in use. Do not apply to institutions (including libraries, sports facilities, etc.) when occupants are present in the immediate area. Remove pets and birds and cover fish aquaria before surface applications.


**HAZARDS TO ANIMALS:**

Do not use on dogs under 12 weeks of age. Do not reapply product for 7 days. Consult a veterinarian before using this product on debilitated, aged, pregnant or nursing animals.

Sensitivities may occur after using ANY pesticide product for pets. If signs of sensitivity occur bathe you pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately.

Certain medications can interact with pesticides. Consult a veterinarian before using on medicated animals.

| <b>FIRST AID</b>  |  |
|---|--|
| <b>If inhaled</b>   | <ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>  |
| <b>If on skin or clothing</b>   | <ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>  |
| <b>If in eyes</b>   | <ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>  |
| <b>If swallowed</b>   | <ul style="list-style-type: none"> <li>• Call poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by the poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul> |
| <b>HOT LINE NUMBER</b>  |  |
| Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-xxx-xxxx for emergency medical treatment information. |  |



3/5

**SIDE/BACK PANEL**

**ENVIRONMENTAL HAZARDS**

This product is extremely toxic to fish. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters.

**DIRECTIONS FOR USE**

**It is a violation of Federal law to use this product in a manner inconsistent with its labeling.**

**READ ENTIRE LABEL BEFORE EACH USE.**

Do not use this product in or on electrical equipment due to the possibility of shock hazard.

Do not use in food/feed areas of food/feed handling establishments, restaurants or other areas where food/feed is commercially prepared or processed. Food/feed areas of commercial food/feed handling establishments where the product cannot be used include areas for receiving, serving, storage (dry, cold, frozen, raw), packaging (canning, bottling, wrapping, boxing), preparing (cleaning, slicing, cooking, grinding), edible waste storage, enclosed processing systems (mills, dairies, edible oils, syrups). Do not use in serving areas while food is exposed or facility is in operation. Serving areas are areas where prepared foods are served such as dining rooms but excluding areas where foods may be prepared or held. In the home, all food processing surfaces and utensils should be covered during treatment or thoroughly washed before use. Exposed food should be covered or removed.

Nonfood areas are areas such as garbage rooms, lavatories, floor drains (to sewers), entries and vestibules, offices, locker rooms, machine rooms, boiler rooms, garages, mop closets and storage (after canning or bottling).

Not for use on plants being grown for sale or other commercial use, or for commercial seed production or for research purposes. For use on plants intended for aesthetic purposes and being grown in ornamental gardens or parks or grounds.


Not for Use in Federally inspected Meat and Poultry Plants.

Affix trigger, pump [or other type] sprayer to container for application. Do not transfer to another container.

**SHAKE WELL BEFORE USING**

**ON LIVESTOCK:** To protect cattle (beef and dairy), goats, sheep, hogs and horses from horn flies, face flies, house flies, mosquitoes and gnats: Apply a light mist sufficient to wet the surface of the hair. For face flies, spray face and head, but do not spray into eyes.

**TO CONTROL STABLE FLIES, HORSE FLIES, AND DEER FLIES:** Apply at a rate of 2.0 ounces per adult animal, sufficient to wet the hair thoroughly. Repeat treatment daily or at intervals necessary to give continued protection.



**SIDE/BACK PANEL**

**TO CONTROL BLOOD-SUCKING LICE:** Apply to the infested areas of the animal, using a stiff brush to get the spray to the base of the hair. Repeat every 3 weeks if required.

**TO CONTROL POULTRY LICE:** Spray roosts, walls and nests or cages thoroughly. This should be followed by spraying over the birds with a fine mist.

**TO CONTROL BEDBUGS AND MITES IN POULTRY HOUSES:** Spray crevices of roost poles, cracks in walls, and cracks in nests where the bedbugs and mites hide.

**ON DOGS:** To control fleas, ticks and lice: Start spraying at the tail, moving the dispenser rapidly and making sure that the animal's entire body is covered, including the legs and underbody. While spraying, fluff the hair so that the spray will penetrate to the skin. Make sure spray wets thoroughly. Wear a mitt or glove. Do not spray into eyes and face. Avoid contact with genitalia. Do not reapply product for 7 days. Use only on Dogs. **Do not use on dogs under 12 weeks of age.**

**SURFACE/PREMISE SPRAYING**


To control cockroaches, waterbugs, palmetto bugs, scorpions, millipedes, centipedes, sowbugs, pillbugs, ants, silverfish, firebrats, spiders, crickets, clover mites, cheese mites, granary weevils, rice weevils, confused flour beetles, rust red flour beetles, drugstore beetles, meal worms, grain mites and cadelles: Direct the spray into hiding places, cracks and crevices, behind shelves and drawers. For silverfish, spray bookcases. For ants, spray trails, nests and point of entry.

**TO CONTROL CARPET BEETLES:** Spray edges of carpeting and under carpeting and rugs. Make localized application to floor and baseboards. Spray directly into cracks, crevices and infested area of shelving. Repeat treatment as needed.

**TO CONTROL BEDBUGS:** Spray mattresses lightly, particularly around tufts and seams. Allow mattress to dry thoroughly before use. Take beds apart and spray in all joints. Treat baseboards, moldings and floors. Repeat treatment as needed.

**TO CONTROL FLEAS AND TICKS (ADULTS AND LARVAE):** Thoroughly spray infested areas, pet beds, resting quarters, nearby cracks and crevices, along and behind baseboards, moldings, window and door frames, and entire areas of floor and floor covering. Fresh bedding should be placed in animal quarters following treatment. Repeat treatment as needed.

**TO CONTROL CLOTHES MOTHS:** Remove any infested articles from storage, brush thoroughly, and air for several hours in sunlight, if possible. Apply spray liberally to empty chests, closets, bureaus and other storages directing the spray into cracks, joints and crevices. After airing, infested articles may be treated lightly, keeping nozzle at least three feet from fabric to avoid staining. Allow articles to dry thoroughly before using.



## SIDE/BACK PANEL

5/5

**TO KILL LICE AND LOUSE EGGS:** Spray in an inconspicuous area to test for possible staining or discoloration. Inspect again after drying, then proceed to spray entire area to be treated. Spray from a distance of 8 to 10 inches. Treat only those garments and parts of bedding, including mattresses and furniture that cannot be either laundered or dry cleaned. Allow all treated articles to dry thoroughly before use.

## **OUTDOOR APPLICATION**

**FOR CONTROL OF GARDEN INSECTS:** For use on roses, dahlias, asters and other ornamentals to kill Japanese beetles hit by spray as well as whiteflies, aphids, exposed thrips, spiders, red mites and leafminers.

May be used on plants as a multi-purpose spray in small gardens and for spot treatments of incipient insect infestations to prevent spreading in large garden areas.

May also be used on other plants such as African violets, asters, azaleas, begonias, camellias, carnations, chrysanthemums, dahlias, delphiniums, dogwoods, English ivy, Kentia palms, laurels, marigolds, rhododendrons, roses, rubber plants, snapdragons, stocks, wandering Jews and zinnias.

Effective when used as directed against whiteflies, aphids, spiders and red mites, exposed thrips, lace bugs, Japanese beetles, pavement ants, clover mites and army worms.

When spraying plants, do not operate closer than 18 inches, using sweeping motion. Be cautious about wetting tender foliage, young plants and new growth. Do not spray plants when temperatures exceed 90° F. Contact insects directly whenever possible. Repeat treatment as infestations occur.

## **STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage or disposal.

**Pesticide Storage:** Store in a cool dry place.

**Pesticide Disposal:** Securely wrap original container in several layers of newspaper and discard in trash.

**Container Disposal:** Do not reuse empty containers. Rinse thoroughly before discarding in trash.

## **NOTICE OF WARRANTY**

PM RESOURCES, INC. MAKES NO WARRANTY OF MERCHANTABILITY, FITNESS FOR ANY PARTICULAR PURPOSE, OR OTHERWISE, EXPRESSED OR IMPLIED CONCERNING THIS PRODUCT OR ITS USES WHICH EXTEND BEYOND THE USE OF THE PRODUCT UNDER NORMAL CONDITIONS IN ACCORD WITH THE STATEMENTS MADE ON THIS LABEL.