

PM 04

67517-66

11/19/98

Page 1 of 4
Jacked Copy 8 4

FRONT PANEL

ASSAULT ALL WEATHER BAIT
Contains Bitrex

For Indoor and Outdoor Rodent Control

Kills Roof Rats, Norway Rats and House Mice

Roof Rats, Norway Rats and House Mice usually consume a lethal dose in a single day's feeding, but it may take two or more days from time of bait consumption for these rodents to die. Rodents cease feeding after consuming a lethal dose. ASSAULT ALL WEATHER BAIT is effective against anticoagulant-resistant rats and mice.

ACTIVE INGREDIENT:

Bromethalin: N-methyl-2,4,-dinitro-N-(2,4,6-tribromophenyl)-
6-(trifluoromethyl) benzenamine.....

0.01%


INERT INGREDIENTS..... 99.99%

TOTAL 100.00%

KEEP OUT OF THE REACH OF CHILDREN

CAUTION

SEE SIDE/BACK PANEL FOR ADDITIONAL
PRECAUTIONARY STATEMENTS



EPA Reg. No.: 67517-66

EPA Est. No.: 67517-MO-1

NET WT.: 9 lbs.
(288 x 0.5 Oz.)

Manufactured By
PM RESOURCES, INC.
13001 St. Charles Rock Road
Bridgeton, MD 63044

(Lot/Code/Product/Rev. Numbers as needed)

Mfg. & Printed in U.S.A.

Precautionary Labeling Must Meet Type Size and Prominence Requirements.

PAGE 1 OF 4
AUGUST 11, 1998
PMR66LAC.DOC-1

> delete

3 9 4

SIDE/BACK PANEL

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION

Keep away from humans, domestic animals and pets. May be harmful or fatal if swallowed. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling bait.

NOTE TO PHYSICIAN:

ASSAULT ALL WEATHER BAIT is not an anticoagulant type of rodenticide. If ingested, limit absorption by either emesis or gastric lavage. Sublethal symptoms, if present, would be the result of cerebral edema and should be treated accordingly through administration of an osmotic diuretic and corticosteroid.

ENVIRONMENTAL HAZARDS

This product is toxic to mammals and birds. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

READ THIS LABEL: Read this entire label and follow all use directions and use precautions.

IMPORTANT: Do not expose children, pets or other nontarget animals to rodenticides. To help to prevent accidents:

1. Store product not in use in a location out of reach of children and pets.
2. Apply bait in locations out of the reach of children, pets, domestic animals and nontarget wildlife, or in tamper-resistant bait stations. These stations must be resistant to destruction by dogs and by children under six years of age, and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from stations when they are lifted, units must be secured or otherwise immobilized. Even stronger bait stations are needed in areas open to hoofed livestock, raccoons, bears, or other potentially destructive animals, or in areas prone to vandalism.
3. Dispose of product container and unused, spoiled and unconsumed bait as specified on this label.

SIDE/BACK PANEL

DIRECTIONS FOR USE (CONTINUED)

USE RESTRICTIONS: This product may be used to control Norway rats, roof rats, and house mice, in and around sewers, homes, industrial and agricultural buildings, and similar man-made structures. This product may also be used in alleys located in urban areas, inside transport vehicles (ships, trains, aircraft), and in and around related port or terminal buildings. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food.

SELECTION OF TREATMENT AREAS: Determine areas where rodents will most likely find and consume the bait. Generally, these are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or their signs have been seen. Remove as much alternative food accessible to rodents as possible.


APPLICATION DIRECTIONS:

Rats: Apply 2 to 12 baits per placement. Space placements at intervals of 20 to 30 feet.

Mice: Apply 1 to 2 baits per placement. Space placements at intervals of 8 to 12 feet.

Rats and Mice: Adjust the amount of bait applied to the level of rodent feeding expected at each bait placement location. Highest bait consumption is expected to occur on the first day or two after treatment. After several days, inspect bait placements and replenish bait at sites where there is heavy feeding or where there is continued evidence of rodent activity. Maintain bait for at least one week and continue baiting until all signs of rodent activity have ceased or it becomes clear that the remaining rodents are not attracted to the bait.

Collect and properly dispose of all dead rodents and leftover bait. To discourage reinfestation, limit sources of rodent food, water, and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish as needed.



4 9 4

SIDE/BACK PANEL

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in original container in a cool, dry place inaccessible to children and pets. Keep containers closed and away from other chemicals.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Do not reuse empty containers. Dispose of liner in a sanitary landfill or by incineration, or, if allowed by state or local authorities, by burning. If burned, stay out of smoke.

NOTICE OF WARRANTY

FM RESOURCES, INC. MAKES NO WARRANTY OF MERCHANTABILITY, FITNESS FOR ANY PARTICULAR PURPOSE, OR OTHERWISE, EXPRESSED OR IMPLIED CONCERNING THIS PRODUCT OR ITS USES WHICH EXTEND BEYOND THE USE OF THE PRODUCT UNDER NORMAL CONDITIONS IN ACCORD WITH THE STATEMENTS MADE ON THIS LABEL.