

67517-87

10/7/2010

1/19



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, D.C. 20460

OFFICE OF CHEMICAL SAFETY  
AND POLLUTION PREVENTION

Mr. Joseph E. Dyer  
PM Resources, Inc.  
13001 St. Charles Rock Road  
Bridgeton, MO 63044

OCT - 7 2010

Subject: Label Notification(s) for Pesticide Registration Notice 98-10

Dear Mr. Dyer:

The Agency is in receipt of your Application(s) for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 dated June 23, 2010 for the following product(s):

**Ecto B736 Insecticide**

**Reg. No. 67517-87**

The Registration Division (RD) has conducted a review of this request for applicability under PR Notice 98-10 and finds that the label change(s) requested falls within the scope of PRN 98-10. The label has been date-stamped "Notification" and will be placed in our records.

If you have any questions, please contact Linda A. DeLuise at 703-305-5428.

Sincerely,

A handwritten signature in dark ink, appearing to read "Richard J. Gebken".

Richard J. Gebken  
Product Manager  
Insecticide Branch  
Registration Division (7504P)

Please read instructions on reverse before completing form.

Form Approved OMB No. 2070-0060



United States  
Environmental Protection Agency  
Washington, DC 20460

☐ Registration  
☐ Amendment  
☒ Other

OPP Identifier Number

## Application for Pesticide - Section I

1. Company/Product Number <b>67517-87</b>	2. EPA Product Manager <b>RICHARD GEBKEN</b>	3. Proposed Classification <input type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) <b>ECTO BT36 INSECTICIDE</b>	PM# <b>10</b>	
5. Name and Address of Applicant (Include ZIP Code) <b>PM RESOURCES, INC 13001 ST. CHARLES ROCK ROAD BRIDGETON, MD 63044</b> <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: <b>NOTIFICATION</b> EPA Reg. No. <b>OCT - 7 2010</b> Product Name	

## Section - II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated
<input type="checkbox"/> Resubmission in response to Agency letter dated	<input type="checkbox"/> "Me Too" Application.
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

FOR THE CONTENTS AND ATTACHMENTS TO THIS NOTIFICATION OF LABEL CHANGE REFER TO THE COVER LETTER - NOTIFICATION IN RESPONSE TO R GEBKEN LETTER DATED June 4, 2009

## Section - III

1. Material This Product Will Be Packaged In:			
Child-Resistant Packaging <input type="checkbox"/> Yes* <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	2. Type of Container <input type="checkbox"/> Metal <input type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify)
* Certification must be submitted		If "Yes" Unit Packaging wgt. No. per container	If "Yes" Package wgt No. per container
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container	5. Location of Label Directions <input type="checkbox"/> On Label <input type="checkbox"/> On Labeling accompanying product
6. Manner in Which Label is Affixed to Product <input type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled		<input type="checkbox"/> Other	

## Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name <b>JOSEPH E. DYER</b>	Title <b>REGULATORY AGENT</b>	Telephone No. (Include Area Code) <b>816-637-2015</b>
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.		6. Date Application Received (Stamped)
2. Signature <b>Joseph E. Dyer</b>	3. Title <b>REGULATORY AGENT</b>	
4. Typed Name <b>JOSEPH E. DYER</b>	5. Date <b>06/23/09</b>	

**13001 St. Charles Rock Road  
Bridgeton, MO 63044  
314-291-6720 Fax 314-291-2657**

June 22, 2009

Document Processing Desk (NOTIFICATION PYRETHOIDS)  
Office of Pesticide Programs (7504P)  
U.S. Environmental Protection Agency  
Room S-4900, ONE Potomac Yard  
2777 South Crystal Drive  
Arlington, VA 22202-4501  
ATTENTION: Richard Gebkin

Subject: APPLICATION FOR PESTICIDE NOTIFICATION OF LABEL CHANGE  
Product Name: ECTO B736 INSECTICIDE, EPA Reg. No.: 67517-87  
Your Letter Dated June 4, 2009 - Submission of Revised Label

Dear Mr. Gebken:  
In accordance with 40CFR 152.44, PM Resources, Inc. submits herewith an application for pesticide NOTIFICATION for subject product.

The purpose of this notification is to allow for the update of our label in accordance with subject letter dated June 4, 2009.

The contents and attachments to this amendment include the following:

- ```
-- Cover Letter
-- Response to EPA Letter From R. Gebken Dated June 4, 2009
-- Application For Pesticide NOTIFICATION, EPA Form 8570-1
-- ECTO B736 Insecticide - Revised Label B7367LAE
-- EPA Accepted Label B7367LAD dated June 01, 2004
```

"Notification of label change per June 4, 2009, letter from Environmental Protection Agency. This notification is consistent with the guidance in the June 4, 2009 letter and the requirements of EPA's regulations at 40 CFR part 156. No other changes have been made to the labeling or the Confidential Statement of Formula for this product. I understand that it is a violation of 18 U.S.C. Sec.1001 to willfully make any false statement to EPA. I further understand that if the amended label is not consistent with the requirements of 40 CFR part 156, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA."

If you have any questions concerning this NOTIFICATION please contact me at 816-637-2015 or by fax at 816-637-2116.

Sincerely,

Joseph E. Dyer

Joseph E. Dyer  
Regulatory Agent  
B736NL1A.DOC-1  
Enclosures

SENT RESPONSE BY UPS SECOND DAY AIR TO THE ABOVE ADDRESS:

4/19

DRAFT LABEL/FRONT PANEL  
ECTO B736 INSECTICIDE

A SYNERGIZED, CONCENTRATE INSECTICIDE WITH UP TO 30 DAYS RESIDUAL EFFECTS ON INANIMATE SURFACES

- °Water Based With Pleasant Citrus Scent °Broad Spectrum Insecticide/Acaracide
- °Synergized °Quick Knockdown
- °For Control of Insect Pests on Livestock, Poultry and Farm Premises
- °For Control of Insect Pests on Dogs and Horses
- °For Control of Ticks Including Deer Ticks (carrier of Lyme Disease)
- °Kills troublesome outdoor insects which destroy Lawns, Roses, Flowers, Trees and Shrubs, Fruit and Nut Trees, and Vegetables
- °Provides contact and residual control for up to 4 weeks on many hard-to-kill sucking and chewing insects.
- °Protects Horses by Killing and Repelling Horse Flies, House Flies, Stable Flies, Face Flies, Hornflies, Deer Flies, Gnats, Mosquitoes, Lice, and Ticks (Including Deer Ticks - Vector of Lyme Disease) on Horses.

Protects your lawns, trees, & shrubs, roses, & flowers, fruits & nuts, vegetables from: Alfalfa loopers, Ants, Aphids, Armyworms, Bagworms, Beetles, Borers, Budworms, Cabbageworms, Chinch bugs, Cicadas, Citrus black flies, Cluster flies, Codling moths, Corn earworms, Crickets, Cutworms, Earwigs, Elm spanworms, Fall cankerworms, Fire Ants, Fleas, Green fruitworms, Gnats, Grasshoppers, House flies, Inchworms, Japanese beetles, leafhoppers, Leafminers, Leafrollers, Lone Star ticks, Lygus bugs, Mealybugs, Millipedes, Mole crickets, Mosquitoes, Moths, Navel orangeworms, Scales, Sod webworms, Sowbugs (pillbugs), Spiders mites, Spring cankerworms, Tent caterpillars, Thrips, Tobacco budworms, Weevils, Whiteflies and many others.

**ACTIVE INGREDIENTS:**

|                                                       |        |
|-------------------------------------------------------|--------|
| Permethrin (CAS NO.: 52645-53-1) .....                | 34.00% |
| Piperonyl Butoxide Technical (CAS NO.: 51-03-6) ..... | 8.50%  |

|                          |         |
|--------------------------|---------|
| INERT INGREDIENTS: ..... | 57.50%  |
| TOTAL                    | 100.00% |

Contains 347.5 g permethrin and 86.9 g piperonyl butoxide technical per quart.

KEEP OUT OF THE REACH OF CHILDREN  
CAUTION  
SEE SIDE/BACK PANEL FOR  
ADDITIONAL PRECAUTIONARY STATEMENTS

EPA Reg. No.: 67517-87

EPA Est. No.: 67517-MO-1

Net Contents: 8 fl. oz. (236ml)

Manufactured For  
PM Resources, Inc.  
13001 St. Charles Rock Road  
Bridgeton, MO 63044

PAGE 1 OF 16  
June 22, 2009  
B7367LAE.DOC-1

**NOTIFICATION**  
**OCT - 7 2010**

5/19

DRAFT LABEL  
SIDE/BACK PANEL

=====

ECTO B736 INSECTICIDE

READ ENTIRE LABEL BEFORE USE - FOR USE ON DOGS ONLY - DO NOT USE ON CATS

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMAN & DOMESTIC ANIMALS

CAUTION

**Hazards to Humans:** Causes moderate eye irritation. Harmful if swallowed. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash clothing before reuse.

**Hazards to Domestic Animals:** For external use only. Harmful if swallowed or absorbed through the skin. Causes moderate eye irritation. Avoid contact with eyes. **Do not use on puppies under 12 weeks of age.** Consult a veterinarian before using this product on debilitated, aged, pregnant, or nursing dogs. Groomers or other persons applying the product frequently must wear chemically resistant gloves.

Sensitivities may occur after using ANY pesticide product for pets. If signs of sensitivity occur bathe your pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately.

Certain medications can interact with pesticides. Consult a veterinarian before using on medicated dogs.

FIRST AID

=====

- If Swallowed**
- Call a poison control center or doctor immediately for treatment advice.
  - Have person sip a glass of water if able to swallow.
  - Do not induce vomiting unless told to do so by a poison control center or doctor.
  - Do not give anything by mouth to an unconscious person.
- 

- If Inhaled**
- Move person to fresh air.
  - If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.
  - Call a poison control center or doctor for further treatment advice.
- 

- If On Skin or Clothing**
- Take off contaminated clothing.
  - Rinse skin immediately with plenty of water for 15-20 minutes.
  - Call a poison control center or doctor for treatment advice.
- 

- If In Eyes**
- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
  - Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eyes.
  - Call a poison control center or doctor for treatment advice.
- =====

HOT LINE NUMBER

=====

Have the product container or label with you when calling a poison control center or doctor for treatment. You may also contact 1-800-XXX-XXXX for emergency medical treatment information.

=====

6/19

DRAFT LABEL  
SIDE/BACK PANEL

=====

ECTO B736 INSECTICIDE

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to fish and other aquatic invertebrates. To protect the environment do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area.

Do not water the treated area to the point of run-off.  
Do not make applications during rain.

All outdoor applications must be limited to spot and crack-and-crevice treatments only, except for the following permitted uses.

- (1) Treatment to soil or vegetation around structures
- (2) Applications to lawns, turf, and other vegetation.
- (3) Applications to building foundations, up to a maximum height of 2 feet.

Other than applications to building foundations, all outdoor applications to impervious surfaces (such as windows, doors, and eaves) are limited to spot and crack-and-crevice applications, only.

PHYSICAL OR CHEMICAL HAZARDS

Do not use this product in or on electrical equipment due to the possibility of shock hazard. Do not use or store near heat or open flame.

INDICATIONS

For use on Horses, Beef and Dairy Cattle, Swine, Sheep, Poultry, Dogs, and their premises to Kill Flies, Fleas, Lice, Imported Fire Ants, Mites, and Ticks including Deer Ticks (carrier of Lyme Disease). Aids in Control of Cockroaches, Mosquitoes Spiders and Crickets.

**BUYER GUIDE:** The following statements are made as a general guide to quantity of pesticide required for various applications. See Chart on inside of back panel for complete directions on dilution and use.

**Premise Use as a Spray or Fog:** Mix 1 pt. in 50 gal. of water to make a ready-to-use spray.

**Livestock Use:** Mix 1 pt. in 200 gal. of water to treat up to 800 animals.

**Poultry Use:** Mix 1 pt. in 100 gal. of water to treat up to 10,000 birds.

**Dog Use:** Mix 1/2 fl. oz. (15 mL) in 2 gal. of water for flea and tick spray or dip.

**Horse Use:** Mix 1 fl. oz. (30 mL) in 5-1/4 gal. of water for spray use.

**CONVERSION EQUIVALENTS:** 1 fl. oz. = 30 mL = 2 tablespoons = 6 teaspoons

7/19

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

#### SHAKE WELL BEFORE USE.

Add the required amount of concentrate to water and blend thoroughly. Do not hold dilutions for more than 24 hours. Dilutions of this product should be agitated or mixed at least once every two hours.

Mix ECTO B736 INSECTICIDE and apply the use-diluted material to animals and pest breeding or resting surfaces at the rates shown on accompanying chart. These dilutions and rates will provide most efficient pest control under conditions of heavy pressure when good contact is achieved. Timing and frequency of application should be based on pest populations reaching nuisance levels, but accompanying manure removal and sanitation practices should precede sprays.

Do not spray feed, food or water. Retreat as needed, but not more often than once every two weeks.

=====

ECTO B736 INSECTICIDE - DIRECTIONS FOR USE (CONT'D)

-----

**PESTS ON LIVESTOCK:** BEEF CATTLE, LACTATING AND NON-LACTATING DAIRY CATTLE, SHEEP, GOATS AND SWINE

NOTE: For Swine allow 5 days between last treatment and slaughter. Retreat as needed, but not more often than once every two weeks. Wash udders thoroughly before milking.

-----

| PESTS                                                                                                                                                                                                                                                    | DILUTIONS FOR USE                                                                                                                                                                                                                        | HOW TO APPLY                                                                                                                                                                             |
|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Horn Flies, Face Flies, Stable Flies, House Flies, Lesser House Flies, Horse Flies, Deer Flies, Black Flies, Mosquitoes, Eye Gnats, Blow Flies, Manage Mites, Scabies Mites, Lice, Ticks including Deer Ticks (carrier of Lyme Disease), and Sheep Keds. | Mix 1 part concentrate in 1,600 parts water (1 pt. in 200 gal. <u>or</u> 7 mL in 3 gal.). For severe infestations <u>or</u> for longer residual control, mix 1 part concentrate in 528 parts water [1 pt. in 66 gal. or 7 mL in 1 gal.]. | <b>High Pressure Spray:</b><br>Spray to thoroughly wet entire animal (1-2 qts. per animal). For lice and mites, a second treatment is recommended in 14 days.                            |
| Horn Flies, Face Flies, Ear Ticks and Lice                                                                                                                                                                                                               | Mix 3.5. fl. oz. (100 mL) in 2 gal. water.                                                                                                                                                                                               | <b>Low Pressure Spray:</b><br>For flies and lice, apply 4-5 fl. oz. (120-150 mL) along midline from face to tailhead. For ear ticks, spray directly onto each ear (1/2 fl. oz. per ear). |

CONVERSION EQUIVALENTS: 1 fl. oz. = 30 mL = 2 Tablespoons = 6 teaspoons

-----

**PESTS ON POULTRY:**

| PESTS                                                         | DILUTIONS FOR USE                                                                                                                                                                                                                 | HOW TO APPLY                                                                                                        |
|---------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------|
| Chicken Mite, Northern Fowl Mite, Chiggers, Lice and Bed Bugs | Mix 1 part concentrate in 800 parts water (1 pt. in 100 gal. <u>or</u> 9.5 mL in 2 gal.). For severe infestation or for longer residual control, mix 1 part concentrate in 400 parts water (1 pt. to 50 gal. or 19 mL in 2 gal.). | <b>High Pressure Spray:</b><br>Spray 1 gal. of coarse spray per 100 birds paying particular attention to vent area. |

CONVERSION EQUIVALENTS: 1 fl. oz. = 30 mL = 2 Tablespoons = 6 teaspoons

=====



DRAFT LABEL - SIDE/BACK PANEL

ECTO B736 INSECTICIDE - DIRECTIONS FOR USE (CONT'D)

**PESTS ON FARM PREMISES:** Barns, Loafing Sheds, Milking Parlors, Feed Lots, Stables, Paddocks, Poultry and Livestock Housing, Animal Hospitals, Kennels and Outside Meat Processing Premises. Remove animals prior to treatment for space spraying.

| PESTS                                                                                                                                                                                                                                                                                                    | DILUTIONS FOR USE                                                                                                                                                                                                                     | HOW TO APPLY                                                                                                                                                                                                                                   |
|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| House Flies, Stable Flies, Lesser House Flies, other Manure Breeding Flies, Bed Bugs, Chicken Mites, Fleas, Lesser Mealworms (Darkling Beetles), Lice, Mites, Ticks including Deer Ticks (carrier of Lyme Disease). Aids in control of Cockroaches, Ants, Spiders, Mosquitoes, Crickets, and Face Flies. | Mix 1 part concentrate in 400 parts water (1 pt. in 50 gal. <u>or</u> 19 mL in 2 gal.). For severe infestation or for longer residual control, mix 1 part concentrate in 136 parts water (1 pt. in 17 gal. <u>or</u> 28 mL in 1 gal.) | <b>SURFACE SPRAY:</b> Spray all surfaces thoroughly with diluted emulsion using 1 gal. per 750 - 1000 sq. ft, but do not let run off occur                                                                                                     |
|                                                                                                                                                                                                                                                                                                          | In mist blower or power fogger (non-thermal) or overhead space spray system, mix 1 part concentrate in 80 parts water (1 pt. in 10 gal. or 48 mL in 1 gal.)                                                                           | <b>INDOOR SPACE SPRAY:</b> Apply as fog or fine mist until area is filled with mist, using 2 fl. oz. (60 mL) per 1000 cubic ft. (Can be used in Milk Rooms).<br><b>OUTDOOR SPACE SPRAY:</b> Fill area with mist. Apply while the air is still. |

CONVERSION EQUIVALENTS: 1 fl. oz. = 30 mL = 2 Tablespoons = 6 Teaspoons

DO NOT APPLY DILUTIONS FOR **PREMISE TREATMENT** DIRECTLY TO LIVESTOCK OR POULTRY. ENSURE THAT FEED AND WATER ARE NOT CONTAMINATED BY SPRAY DRIFT. FOR **SPACE SPRAYING** DO NOT REMAIN IN TREATED AREAS AND VENTILATE THE AREA BEFORE REOCCUPYING. THE USE OF ANY RESIDUAL FLY SPRAY SHOULD BE SUPPLEMENTED WITH PROPER MANURE MANAGEMENT AND GENERAL SANITATION TO ELIMINATE FLY BREEDING SITES.

|                    |                                        |                                                                                                                                                                                                                                                                                                                          |
|--------------------|----------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Imported Fire Ants | Mix 7 fl. oz. (210 mL) in 2 gal. water | <b>SPRINKLER CAN:</b> Gently sprinkle one gal. over the surface of each mound. Thoroughly wet mound and surrounding area to a 4 ft. diameter. For best results, apply in cool weather 65-85° (early morning or late evening). Pressurized sprayers may disturb ants and cause migration, reducing product effectiveness. |
|--------------------|----------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|

CONVERSION EQUIVALENTS: 1 fl. oz. = 30 mL = 2 Tablespoons = 6 Teaspoons

9/19  
b1/b

DRAFT LABEL - SIDE/BACK PANEL

ECTO B736 INSECTICIDE - DIRECTIONS FOR USE (CONT'D)

**PESTS ON DOGS AND THEIR PREMISES:**

NOTE: USE ONLY ON DOGS. DO NOT USE ON CATS. Do not treat puppies under 12 weeks of age. Do not get this product in your dog's eyes or mouth. Avoid contact with genitalia. Reapply as necessary.

| PESTS                                                                                                    | DILUTIONS FOR USE                                                                                            | HOW TO APPLY                                                                                                                                                                                                                                                                                                 |
|----------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Fleas, Ticks (including Deer Ticks, carrier of Lyme Disease) Mange Mites, Lice and Stable Flies on Dogs. | Mix 1 part concentrate in 512 parts water (1/2 fl. oz. <u>or</u> 15 mL in 2 gal.).                           | <b><u>SPRAY, DIP OR SPONGE ON:</u></b><br>Spray, dip, sponge or pour diluted product on the dog until its hair and skin are thoroughly wet, making sure that animal's entire body is treated. Do not treat dog's face. Let drip dry on animal and do not rinse off.                                          |
|                                                                                                          | Mix 1 part concentrate in 64 parts water (1/2 fl. oz. <u>or</u> 15 mL in 1 quart).                           | <b><u>LOW VOLUME SPRAY:</u></b><br>Spray 1-2 fl. oz. per animal. Starting at the tail, stroke against the lay of the hair, spraying the parted hair with a fine mist directly behind the hand to insure penetrating the coat. Spray entire animal until hair coat is damp.                                   |
| Fleas and Ticks on Dog Premises.                                                                         | Mix 1 to 2 parts concentrate in 160 parts water (1 pt. in 10 to 20 gal. <u>or</u> 24 mL to 48 mL in 1 gal.). | <b><u>INDOOR PREMISE SPRAY:</u></b><br>Thoroughly spray infested areas, pet beds, resting quarters, nearby cracks and crevices, along baseboards, moldings, windows and door frames, and localized areas of floor and floor covering. Fresh bedding should be placed in animal quarters following treatment. |
|                                                                                                          | Mix 1 fl. oz. (30 mL) concentrate in 10.5 gal. water <u>or</u> 7 mL in 2.5 gal. water.                       | <b><u>OUTDOOR PREMISE SPRAY:</u></b><br>Use enough finished spray to penetrate foliage, usually 50-100 gal. per acre. To prevent infestation of buildings, treat a band of vegetation 6-10 feet adjacent to the structure.                                                                                   |

CONVERSION EQUIVALENTS: 1 fl. oz. = 30 mL = 2 Tablespoons = 6 Teaspoons

10/19

DRAFT LABEL - SIDE/BACK PANEL

ECTO 8736 INSECTICIDE - DIRECTIONS FOR USE (CONT'D)

**PESTS ON EQUINE PREMISES:** Barns, Stables, Paddocks, and Animal Hospitals. Remove animals prior to treatment for space spraying.

| PESTS                                                                                                                                                                                                                                                                          | DILUTIONS FOR USE                                                                                                                                                                                                                            | HOW TO APPLY                                                                                                                                                                                                                    |
|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| House Flies, Stable Flies, Lesser House Flies, other Manure Breeding Flies, Bed Bugs, Chicken Mites, Fleas, Mealworms, Lice, Mites, Ticks including Deer Ticks (carrier of Lyme Disease). Aids in control of Cockroaches, Ants, Spiders, Mosquitoes, Crickets, and Face Flies. | Mix 1 part concentrate in 400 parts water (1 pt. in 50 gal. <u>or</u> 19 mL in 2 gal.). For severe infestation <u>or</u> for longer residual control, mix 1 part concentrate in 136 parts water (1 pt. in 17 gal. <u>or</u> 28 mL in 1 gal.) | <b><u>SURFACE SPRAY:</u></b> Spray all surfaces thoroughly with diluted emulsion using 1 gal. per 750 - 1000 sq. ft. Do not allow run off to occur.                                                                             |
|                                                                                                                                                                                                                                                                                | In mist blower or power fogger (non-thermal) or overhead space spray system mix 1 part concentrate in 80 parts water (1 pt. in 10 gal. <u>or</u> 48 mL in 1 gal.)                                                                            | <b><u>INDOOR SPACE SPRAY:</u></b> Apply as fog or fine mist until area is filled with mist, using 2 fl. oz. (60 mL) per 1000 cubic ft.<br><b><u>OUTDOOR SPACE SPRAY:</u></b> Fill area with mist. Apply while the air is still. |

CONVERSION EQUIVALENTS: 1 fl. oz. = 30 mL = 2 Tablespoons = 6 Teaspoons

DO NOT APPLY DILUTIONS FOR **PREMISE TREATMENT** DIRECTLY TO HORSES. ENSURE THAT FEED AND WATER ARE NOT CONTAMINATED BY SPRAY DRIFT. FOR **SPACE SPRAYING** DO NOT REMAIN IN TREATED AREAS AND VENTILATE THE AREA BEFORE REOCCUPYING. THE USE OF ANY RESIDUAL FLY SPRAY SHOULD BE SUPPLEMENTED WITH PROPER MANURE MANAGEMENT AND GENERAL SANITATION TO ELIMINATE FLY BREEDING SITES.

11/19

DRAFT LABEL - SIDE/BACK PANEL

ECTO B736 INSECTICIDE - DIRECTIONS FOR USE (CONT'D)

PESTS ON HORSES: DO NOT USE ON HORSES INTENDED FOR FOOD

| PESTS                                                                                                                                                                                                | DILUTIONS FOR USE                                                                         | HOW TO APPLY                                                                                                                                        |
|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------|
| Horse Flies, Deer Flies, Stable Flies, House Flies,<br>Horn Flies, Face Flies, Black Flies, Mosquitoes,<br>Eye Gnats, Mange Mites, Lice and Ticks (including<br>Deer Ticks - Vector of Lyme Disease) | Mix 1 fl. oz. (30 mL) concentrate in<br>5-1/4 gal. water <u>or</u> 11 mL in 2 gal. water. | <u>SPRAY OR SPONGE ON:</u><br>Thoroughly wet animal to skin with spray or sponge.                                                                   |
|                                                                                                                                                                                                      | Mix 3.5 fl. oz. (105 mL) concentrate in<br>1.5 gal. water <u>or</u> 35 mL in 2 qt. water. | <u>LOW PRESSURE SPRAY OR WIPE ON :</u><br>Spray or wipe on 1-2 fl. oz. per animal. Pay particular attention<br>to treat back, face, legs, and ears. |

Note: Do not use this method of application on foals under 12 weeks of age.  
Repeat treatment as needed. Do not allow the product to contact horses'  
eyes or mucous membranes.

CONVERSION EQUIVALENTS: 1 fl. oz. = 30 mL = 2 Tablespoons = 6 Teaspoons

12/19

**DRAFT LABEL – SIDE/BACK PANEL  
ECTO B736 INSECTICIDE  
DIRECTIONS FOR USE (CONT'D)**

**PESTS ON HOME LAWNS, HOUSE PLANTS, TREES & SHRUBS, ROSES & FLOWERS, FRUIT & NUT TREES, VEGETABLES, NUISANCE PESTS IN OUTDOOR AREAS, AND WOOD INFESTING INSECTS:**

Not for use on plants being grown for sale or for other commercial use, or for commercial seed production or for research purposes.

**Step #1:** For lawns and vegetables, measure the area to be sprayed. Calculate square feet by multiplying length by width. For all other applications, apply as a thorough cover spray.

**Step #2:** Use a clean sprayer. Carefully measure and mix the amount of concentrate and water as directed.

**Step #3:** Spray as directed. Thorough coverage is important. Flush sprayer with clean water after each use. Make applications when insects first appear. Repeat applications as directed. See directions for complete insect list.

People and pets should not be allowed in treated areas until spray has dried.

Use dilution rates as indicated.

1 fl. oz. = 6 tsp. or 2 Tbsp.

8 fl. oz. = ½ pt. or 1 cup

**DIRECTIONS FOR APPLYING**

**HOME LAWNS**

|                                                                                | <b>Pests:</b>                                                                                                                                                                               | <b>How to Use:</b>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                            |
|--------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Bent, Bermuda, Bluegrass, Dichondra, Fescue, Irish Moss, Merion, St. Augustine | Ants, Armyworms, Brown dog ticks, Chinch bugs, Crickets, Cutworms, Fleas, Grasshoppers, Japanese beetle grubs, Mole crickets, Sod webworms, Ticks that may carry Lyme Disease, Leafhoppers. | Thoroughly wet down grass a few hours before applying. Home lawns should be no taller than 3 inches at time of application. Slightly more water may be used as long as the amount of product per area is as listed below. Do not apply more often than two times per month. Do not treat more often than 10 times per year. For heavy infestations, repeat application after 2 weeks.<br><br>For ants, crickets, and grasshoppers, use 1/6 fl. ozs. in 3 gals. (1 tsp. in 3 gals.) of water to cover 1,000 sq. ft. For other listed insects, use 1/2 fl. oz. in 10 gals. (3 tsp. in 10 gals.) of water to cover 1,000 sq. ft. |

14/19

**DRAFT LABEL – SIDE/BACK PANEL**  
**ECTO B736 INSECTICIDE**  
**DIRECTIONS FOR USE (CONT'D)**

**HOUSEPLANTS**

**INDOORS: TO PROTECT PLANTS IN RESIDENTIAL GREENHOUSES AND LATHHOUSES, AND FOR INDOOR LANDSCAPING:**

|                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                | <b>Pests:</b>                                                                                                                                                                             | <b>How to Use:</b>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                  |
|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Ageratum, Ardisia, Aster, Baby's Breath, Begonia, Birds nest Fern, Bleeding Heart, Carnation, Chrysanthemum, Coleus, Crown of Thorns, Cockscomb, Cyclamen, Dracaena, Dumbcane, English Ivy, Fuchsia, Gladiolus, Grape Ivy, Juniper, Lily, Marigold, Orchid, Palm, Pansy, Peperomia, Petunia, Philodendron, Piggy-Back Plant, Poinsettia, Portulaca, Pothos, Prayer Plant, Purple Passion, Rabbits Foot Fern, Rose, Schefflera, Snapdragon, Snake Plant, Statice, Velvet Plant, Verbena, Zinnia | Aphids, Armyworms, Corn earworms, Exposed thrips, Fall cankerworms, Inchworms, Japanese beetles, Leafminers, Leafrollers, Mealybugs, Scales, Spider mites, Spring cankerworms, Whiteflies | Mix 1/2 fl. oz. in 1 gal. (1 tsp. in 2 gal.) of water. Contact and residual: Use sufficient spray to obtain full coverage of all foliage. If application is to be made indoors, remove infested plants from living areas prior to application. Use in an area not likely to be occupied by people or pets (particularly fish). After spraying, plants can be returned to original location. Do not use more product than recommended, as this may cause some plant injury. Use with care on ferns, palms and fuchsia. As with all plants listed above, especially carnations, chrysanthemums, and roses, there exists a wide variety of species and cultivars, and therefore a wide variety of susceptibility to injury. To ascertain possible phytotoxicity, treat a few plants as described above, wait 24 hours, and check for possible injury. Repeat applications (use interval 4 to 8 days) may be necessary to control citrus, and Adult Japanese beetles on roses. Do not apply more often than 8 times per growing season. |
| Velvet Plants                                                                                                                                                                                                                                                                                                                                                                                                                                                                                  | Fungus Gnats                                                                                                                                                                              |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                     |
| <b>Trees &amp; Shrubs</b>                                                                                                                                                                                                                                                                                                                                                                                                                                                                      |                                                                                                                                                                                           |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                     |
|                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                | <b>Pests:</b>                                                                                                                                                                             | <b>How to Use:</b>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                  |
| Arizona Cypress, Azalea, Birch, Cherry, Non-bearing Citrus, Conifers, Elm, English Ivy, Euonymus, Weeping fig, Fir, Honeysuckle Lilac, Mock Orange, Oak, Palm, Philodendron, Pine, Poinsettia, Tulip Poplar, Rhododendron, Taxus                                                                                                                                                                                                                                                               | Aphids, Bagworms, Cicadas, Exposed thrips, Fall cankerworms, Inchworms, Leafminers, Leafrollers, Mealybugs, Spider mites, Whiteflies, Gypsy Moth, Tent Caterpillar, Japanese Beetles      | Use 1/12 fl. oz. in 1 gal. (or 1 tsp. in 2 gal.) of water. Apply as a thorough spray, wetting leaves and branches to the dripping point. Try to penetrate dense foliage. Spray in the late afternoon or evening, when the temperature ranges from 50° to 75°F and when there is little or no wind. Spray at the first sign of insects. Repeat as necessary; use intervals of 4-8 days. Application can be made up to the day of collection. Do not apply more often than 8 times per year.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                          |
| Cherry, Honeysuckle                                                                                                                                                                                                                                                                                                                                                                                                                                                                            | Fall webworms                                                                                                                                                                             |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                     |
| Conifer                                                                                                                                                                                                                                                                                                                                                                                                                                                                                        | Budworms                                                                                                                                                                                  |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                     |
| Elm                                                                                                                                                                                                                                                                                                                                                                                                                                                                                            | Elm leaf beetles, Elm spanworms, Fall webworms                                                                                                                                            |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                     |
| Douglas and Grand Firs                                                                                                                                                                                                                                                                                                                                                                                                                                                                         | Tussock moths                                                                                                                                                                             |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                     |
| Non-bearing Citrus                                                                                                                                                                                                                                                                                                                                                                                                                                                                             | Citrus black flies                                                                                                                                                                        |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                     |
| Oak                                                                                                                                                                                                                                                                                                                                                                                                                                                                                            | Oak worms                                                                                                                                                                                 |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                     |
| Pine                                                                                                                                                                                                                                                                                                                                                                                                                                                                                           | Pine beetles, Pine moths, Pine needleminers, Needle scales                                                                                                                                |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                     |

On Tent caterpillars, application should be made when caterpillars are young and tents are first noticed. For best results, apply in the late afternoon or evening, when caterpillars have returned to their tents. Wet tents with spray on all sides. A few caterpillars may be seen escaping from the tent; they should die within several hours.

15/19

**DRAFT LABEL – SIDE/BACK PANEL**  
**ECTO B736 INSECTICIDE**  
**DIRECTIONS FOR USE (CONT'D)**  
**ROSES & FLOWERS**

|                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                             | <b>Pests:</b>                                                                                                                                                                                                                                                                                                                 | <b>How to Use:</b>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                              |
|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Ageratum, Ardisia, Aster, Azalea, Baby's Breath, Begonia, Birds-nest Fern, Bleeding Heart, Carnation, Chrysanthemum, Coleus, Common Ninebark and Snowberry, Crown of Thorns, Cockscomb, Cyclamen, Dracaena, Dumbcane, English Ivy, Fuchsia, Gladiolus, Grape Ivy, Gold Bells, Hypoestes, Ivy, Juniper, Lilac, Marigold, Orchid, Palm, Pansy, Pea Shrub, Peperomia, Petunia, Philodendron, Piggy-Back Plant, Poinsettia, Portulaca, Pothos, Prayer Plant, Purple Passion, Rabbits Foot Fern, Rhododendron, Rose, Schefflera, Snapdragon, Snake Plant, Statice, Velvet Plant, Verbena, Zinnia | Aphids, Armyworms, Bagworms, Cabbage worms, Cicadas, Cornworms, Cutworms, Exposed thrips, Fall cankerworms, Fall webworms, Fungus gnat, Inchworms, Japanese beetles, Leafminers, Leafrollers, Mealybugs, Scale crawlers, Spring cankerworms, Spider mites, Whiteflies, Japanese Beetles, Rose Chafer, Psyllids, Scale Insects | Use 1/12 fl. oz. in 1 gal. (or 1 tsp. in 2 gal.) of water. Apply as a thorough spray, wetting the plants to the dripping point. Try to hit underside of leaves and penetrate dense foliage. Spray in the late afternoon or evening, when the temperature ranges from 50°F to 75°F and when there is little or no wind. Spray at the first sign of insects. Repeat as necessary; use intervals of 4-8 days. Application can be made up to the day of collection. Do not treat more often than 12 times per year. |

**FRUIT AND NUT TREES**

**GENERAL DIRECTIONS**

Use 1/6 fl. ozs. in 1 gal. (1 tsp. in 1 gal.) of water. Apply when insects appear. Wet the plants to the dripping point. Try to hit underside of leaves and penetrate dense foliage.

|                                         | <b>Pests:</b>                                                                                                                                                                                       | <b>How to Use:</b>                                                                                                                  |
|-----------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------|
| Almonds, Pecans                         | Navel orangeworms, Peach twig borers                                                                                                                                                                | Do not make more than 2 applications during hull split and 5 applications per season. Can be applied up to 7 days prior to harvest. |
| Apples                                  | Green fruitworms, Aphids, Leafrollers (Oblique-banded, Red banded), Plum curculio, Rosy apple aphids, Spotted tentiform Leafminers, Tarnished plant bugs, White apple leafhoppers, Japanese Beetles | Repeat as required to maintain control. Do not make more than 3 applications per season. Do not apply after petal fall.             |
| Peaches                                 | Borers (Lesser peachtree, Peach twig), Green fruitworms, Plum curculio, Oriental fruit moths, Tarnished plant bugs                                                                                  | Do not apply within 7 days of harvest. Do not make more than 8 applications per season.                                             |
| Pears (dormant through delayed dormant) | Pear psylla                                                                                                                                                                                         | Apply during the dormant through delayed dormant growth period only. Do not make more than 2 applications per season.               |
| Pear (summer)                           | Codling Moths, Green fruitworms, Pear psylla, Aphids                                                                                                                                                | Do not make more than 3 applications per season. Can be applied up to 14 days prior to harvest                                      |

16/19

**DRAFT LABEL – SIDE/BACK PANEL  
ECTO B736 INSECTICIDE  
DIRECTIONS FOR USE (CONT'D)  
VEGETABLES**

|                                                                                                                                                                                                                                                                                                                      | <b>Pests:</b>                                                                                                                          | <b>How to Use:</b>                                                                                                                                                                                                                                                     |
|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Asparagus<br>(pre-harvest)                                                                                                                                                                                                                                                                                           | Asparagus beetles, Cutworms                                                                                                            | Use 1/8 fl. oz. in 1 gal. (3 tsp. in 4 gal.). Apply for uniform coverage as needed. Do not make more than 4 applications per season. Can be applied up to 3 days prior to harvest.                                                                                     |
| Asparagus<br>(post harvest)                                                                                                                                                                                                                                                                                          | Larval stages of asparagus beetle, tarnished plant bug; lygus bugs and adult Japanese beetle.                                          | Treat ferns or bust growth after spear harvest when insects are present.                                                                                                                                                                                               |
| Celery – see Leafy Vegetables                                                                                                                                                                                                                                                                                        |                                                                                                                                        |                                                                                                                                                                                                                                                                        |
| Cole Crops, Broccoli, Brussels Sprouts, Cabbage, Cauliflower                                                                                                                                                                                                                                                         | Beet armyworms, Cabbage aphids and loopers, Diamondback moths, Imported cabbageworms                                                   | Use 1/12 fl. oz. in 1 gal. (1 tsp. in 2 gal). Apply, every 5 days or as needed. Can be applied up to 1 day prior to harvest. Do not make more than 8 applications per season on Cauliflower, Brussels Sprouts, and Broccoli and 10 applications per season on Cabbage. |
| Corn – see Sweet Corn                                                                                                                                                                                                                                                                                                |                                                                                                                                        |                                                                                                                                                                                                                                                                        |
| Cucumber, Melons, Pumpkins, Squash                                                                                                                                                                                                                                                                                   | Flea beetle, Leafhopper                                                                                                                | Use 1/6 fl. oz. in 1 gal. (1 tsp. in 1 gal.). Apply, as needed, but not more than 8 applications per season. Can be applied up to 3 days prior to harvest.                                                                                                             |
| <i>Cucurbits:</i> Balsam pear (bitter melon), Chinese wax gourd (Chinese preserving melon), citron melon, cucumber, gherkin, edible gourds; melons including hybrids such as cantaloupe, casaba, crenshaw, honeydew, honey balls, mango melon, muskmelon, Persian melon, pumpkin, summer & winter squash, watermelon | Aphids, Cabbage looper, Cucumber Beetle (adults), Cutworms, Leafhoppers, Leafminers, Melon worms, Pickleworm, Plant bugs, Squash bugs. | Use 1/6 fl. oz. in 1 gal. (1 tsp. in 1 gal). Apply as needed but not more than 8 applications per season. Can be applied up to day of harvest.                                                                                                                         |
| Eggplant                                                                                                                                                                                                                                                                                                             | Colorado potato beetles, Vegetable Leafminers                                                                                          | Use 1/4 fl. oz. in 1 gal. (3 tsp. in 2 gal.). Apply as needed but not more than 16 applications per season. Can be applied up to 3 days prior to harvest.                                                                                                              |
| Horseradish<br>(Foliar applications)                                                                                                                                                                                                                                                                                 | Imported crucifer weevils                                                                                                              | Use 1/4 fl. oz. in 1 gal. (3 tsp in 2 gal). Apply as needed. Do not exceed 3 foliar applications per season. Can be applied up to 22 days prior to harvest.                                                                                                            |
| Head Lettuce                                                                                                                                                                                                                                                                                                         | Corn earworms, Beet armyworms, Loopers (Alfalfa, Cabbage, Tobacco budworms                                                             | Use 1/6 fl. oz in 1 gal. (1 tsp. in 1 gal.). Apply every 5-10 days as needed. Do not make more than 10 applications per season. Can be applied up to 1 day prior to harvest.                                                                                           |



17/19

**DRAFT LABEL – SIDE/BACK PANEL  
ECTO B736 INSECTICIDE  
DIRECTIONS FOR USE (CONT'D)  
VEGETABLES**

|                                                                                                                                                                                                                                                                                                                                     | <b>Pests:</b>                                                                                                                                                                                                     | <b>How to Use:</b>                                                                                                                                                                     |
|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Leafy Vegetables:</b><br>Amaranth, Arrugula,<br>Celery, Celltuce, Chervil,<br>Corn salad,<br>Chrysanthemum (edible-<br>leaved and garland), cress<br>(garden and upland),<br>Dandelion, Dock, Endive,<br>Fennel, Lettuce (head and<br>leaf), Okra, Parsley,<br>Purslane (garden and<br>winter), Rhubarb, Spinach,<br>Swiss chard | Alfalfa looper, Aphids, Beet<br>armyworm, Cabbage looper, Corn<br>earworm, Cutworms, European<br>corn borer, Fall armyworm, Green<br>clover worm, Southern<br>armyworm, Tobacco budworm,<br>Vegetable leaf miner. | Use 1/6 fl. oz. in 1 gal. (1 tsp. in 1 gal.) Apply every 5 to 10<br>days as needed. Can be applied up to 3 days prior to harvest.<br>Do not make more than 10 applications per season. |
| Melons – see Cucurbit                                                                                                                                                                                                                                                                                                               |                                                                                                                                                                                                                   |                                                                                                                                                                                        |
| Potatoes                                                                                                                                                                                                                                                                                                                            | Beet armyworm, Colorado potato<br>beetle, Potato leafhopper, Potato<br>tuberworm, Cabbage looper,<br>cutworms, European corn borer,<br>Potato psyllid, Tarnis plant bug.                                          | Use 1/8 fl. oz. in 1 gal. (3 tsp. in 4 gal.) Apply as needed. Do<br>not make more than 12 applications per season. Can be applied<br>up to 7 days prior to harvest.                    |
| Peppers                                                                                                                                                                                                                                                                                                                             | Vegetable Leafminers, Cabbage<br>looper, Flea beetle, Corn<br>earworm, Pepper weevil,<br>European corn borer.                                                                                                     | Use 1/6 fl. oz. in 1 gal. (1 tsp. in 1 gal.) Apply as needed but not<br>more than 8 applications per season. Can be applied up to 3<br>days prior to harvest.                          |
| Strawberries, Raspberries,<br>Blueberries                                                                                                                                                                                                                                                                                           | Strawberry weevils, Aphids,<br>Spider mites, Lygus bugs, Flea<br>beetles, Omnivorous leaf-tier,<br>Spittlebugs, Japanese beetles,<br>Field crickets, Red Spider,<br>Leafrollers, weevils                          | Use 1/6 fl. oz. in 1 gal. (1 tsp. in 1 gal.) Allow 14 days between<br>last application and harvest. Make no more than 8 foliar<br>applications per growing season.                     |
| Sweet Corn                                                                                                                                                                                                                                                                                                                          | Armyworms (Fall, Southern),<br>corn earworms, European corn<br>borers                                                                                                                                             | Use 1/8 fl. oz. in 1 gal. (3 tsp. in 4 gal.) Apply for uniform<br>coverage when needed but not more than 6 applications per<br>season. Can be applied up to date of harvest.           |
| Tomatoes                                                                                                                                                                                                                                                                                                                            | Beet and southern armyworm,<br>Tomato pinworm, Tomato<br>fruitworm, Horn worms,<br>Granulate cutworms, Colorado<br>potato beetle, Cabbage looper                                                                  | Use 1/6 fl. oz. in 1 gal. (1 tsp. in 1 gal.) Apply for uniform<br>coverage when needed but not more than 6 applications per<br>season. Can be applied up to date of harvest.           |

18/19

**DRAFT LABEL – SIDE/BACK PANEL  
ECTO B736 INSECTICIDE  
DIRECTIONS FOR USE (CONT'D)**

**NUISANCE PESTS IN OUTDOOR AREAS**

**GENERAL DIRECTIONS**

Use 7/12 fl. oz. in 1 quart (14 tsp. in 1 quart) of water, treating 400 linear feet with a 6-inch spray band.

**OUTSIDE SURFACES OF BUILDING**

Use on Ants, Clover mites, Crickets, Earwigs, Millipedes, Sowbugs (pillbugs) to help prevent infestations of buildings. Treat the building foundation to a height of 2 to 3 feet, where pests are active and may find entrance. Also, apply as a residual spray to outside surfaces of building, including porches, screens, window frames, eaves, patios, garages, refuse dumps and other areas where these pests congregate or have been seen. Repeat treatment as needed to maintain effectiveness. Do not treat more often than one time per week and do not treat more often than 9 times per year.

**OTHER OUTDOOR AREAS**

Use on Ants, Crickets, Earwigs, Fleas, Ticks (including ticks that may carry Lyme Disease) for treatment of localized infestations of these insects in areas where there are weeds or bushy non-crop areas, spray infested areas thoroughly. **For ants**, thoroughly wet hills and runways.

To reduce annoyance from: **Cluster flies, Flying moths, Gnats, House flies, Mosquitoes**. Spray outside surfaces of screens, doors, and window frames or wherever these insects may enter the room. Also treat surfaces around light fixtures on porches, in garages, around shrubbery, and other places where these insects alight or congregate.

Repeat application as reinfestations occur. Do not treat more often than one time per week and do not treat more than 9 times per year.

**WOOD-INFESTING INSECTS (including termites)**

**DO NOT USE INSIDE DWELLINGS. For use around existing structure, not for pre-construction use. For Beetles (Furniture, Deathwatch, False Powderpost, Longhorned, Powderpost), Carpenter Ants (and other wood-infesting ants), Carpenter Bees and termites (including drywood termites).** Treat outside of buildings, porches, wooden decks and patios, wooden fences, window frames, doorways, foundations, eaves, garages, garbage areas and other areas where you may find these pests.

Treat outside surfaces of buildings, porches, wooden decks and patios, wooden fences, window frames, doorways, foundations, eaves, garages, garbage areas, and other areas where you may find these pests. Mix 5/16 fl. oz. with 1 gal. (5 tsp. in 1 gal.) of water to treat 100 sq. ft. Spray infested surfaces until thoroughly wet, avoiding dripping and runoff. Apply evenly on wood surfaces. To treat small areas, apply by brushing or spraying the diluted spray. For large areas, use a coarse, low-pressure spray. Do not spray into electrical outlets or utilities.

Spray infested areas and areas likely to be infested, such as wood surfaces, voids and channels in damaged wood, spaces between wooden parts of structure, and between wood and foundations where infestation is likely to occur. Also spray around fence posts, wood piles (do not use on firewood) and tree stumps.

Suitable protective clothing should be worn in confined or overhead areas to prevent exposure to eyes and skin. The use of eye protection clothing should be worn in confined or overhead areas to prevent exposure to eyes and skin. The use of eye protection, protective head covering, chemical resistant gloves and footwear, a long-sleeved shirt and long pants or coveralls are recommended. Do not permit contact with treated surfaces until spray has dried. Treated confined areas, such as detached garages, under porches and decks, should be ventilated during and immediately following application before the areas are re-occupied.

Repeat treatment as needed to maintain effectiveness. Do not apply more often than one time per month and do not treat more often than 4 times per year.

**For carpenter ants**, spray around doors and windows and other places where ants may enter or crawl and hide. Also treat cracks and crevices or openings into wall voids or other areas where these ants may be present.

**For carpenter bees**, spray or inject into holes and tunnels already made by the bees with a tank sprayer and spray areas where they are beginning to build tunnels into the wood.

**For termites\* (including dry wood termites)**, treat as for other wood-infesting insects. This product will kill workers and winged reproductive forms of termites in the treated areas. Re-treat as needed to maintain effectiveness. Do not treat more often than 8 times per year.

**\*USE OF THIS TYPE OF PRODUCT TO CONTROL TERMITES SHOULD NOT BE CONSIDERED A SUBSTITUTE FOR SOIL TREATMENT OF SUBTERRANEAN TERMITES DONE BY A PROFESSIONAL PEST CONTROL OPERATOR. FOR ACTIVE INFESTATION GET A PROFESSIONAL TERMITE INSPECTION.**

19/19

**DRAFT LABEL – SIDE/BACK PANEL  
ECTO B736 INSECTICIDE  
DIRECTIONS FOR USE (CONT'D)**

**STORAGE AND DISPOSAL**

Do not contaminate water, food, or feed by storage or disposal. **KEEP CONTAINER SEARLED WHEN NOT IN USE. PROTECT FROM FREEZING.**

**Pesticide Storage:** Store upright at room temperature. Avoid exposure to extreme temperatures.

**Pesticide Disposal:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**Container Disposal:** Do not reuse empty container. Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

**NOTICE OF WARRANTY**

PM RESOURCES, INC. MAKES NO WARRANTY OF MERCHANTABILITY, FITNESS FOR ANY PARTICULAR PURPOSE, OR OTHERWISE, EXPRESSED OR IMPLIED, CONCERNING THIS PRODUCT OR ITS USES WHICH EXTEND BEYOND THE USE OF THE PRODUCT UNDER NORMAL CONDITIONS IN ACCORD WITH THE STATEMENTS MADE ON THIS LABEL.