



U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division (H7505C)
401 M St. S.W.
Washington, D.C. 20460

EPA Reg. Number:
67546-2

Date of Issuance:
JAN - 9 2001

**NOTICE OF PESTICIDE
Registration**

Term of Issuance:
Conditional Registration

Name of Pesticide Product:
Darkling Beetle Control

(under FIFRA, as amended)

Name and Address of Registrant (include ZIP Code):
**National Boraxx Corp
3690 Orange Place, Suite 480
CLEVELAND, OH 44122**

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:

1. Submit and/or cite all data required for registration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA section 4.
2. Make the following label changes:
 - a. Revise the EPA Registration Number to read, EPA Reg. No. 67546-2
3. In accordance with PR NOTICE 2000-3, the First Aid Statements, must be place within a box which sets that text apart from the rest of the label. This information must be presented in a narrative format and marked with a symbol such as a hyphen or other type "bullet" along the left margin.
4. The First Aid Statements must be revised to read as follows:

IF SWALLOWED:

- o Call a poison control center or doctor immediately for treatment advice.
- o Have a person sip a glass of water if able to swallow.
- o Do not induce vomiting unless told to do so by a poison control center or doctor.
- o Do not give anything by mouth to an unconscious person.

Signature of Approving Official: *Richard A. ...*

Date:

EPA Registration No. 67546-2

IF IN EYES:


- o Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- o Call a poison control center or doctor for treatment advice.

5. Submit two copies of the revised final printed label for the record.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Sincerely,



Ann Sibold
Acting Product Manager 10
Insecticide Branch
Registration Division (7505C)

Signature of Approving Official:	Date: JAN - 9 2001
----------------------------------	--------------------

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND
DOMESTIC ANIMALS**

CAUTION

Harmful if swallowed. Avoid contact with eyes. Wash thoroughly after handling. Avoid contamination of food and feed.

FIRST AID

IF SWALLOWED: Immediately contact Poison Control Center or physician. If these are unavailable, give patient 1 or 2 glasses of water and induce vomiting by touching the back of throat with fingers. Never give anything by mouth or induce vomiting if patient is unconscious.

IF IN EYES, flush with plenty of water. Get medical attention if irritation persists.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

STORAGE: Keep container tightly closed when not in use. Store in original container, in a dry place inaccessible to children and pets.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: DO NOT reuse container. Rinse 3 times, then offer for recycling or trash collection.

NOTICE

Manufacturer shall not be liable in respect to any injury or consequential damages suffered based on the use of this product contrary to the instructions given on this label.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. See side panel for Application Instructions.

**Darkling Beetle
Control**

**An Insecticide for the Control
of Darkling Beetles, Hide
Beetles, and Flies.**

Active Ingredient:
Boric Acid.....100%

**KEEP OUT OF THE REACH
OF CHILDREN
CAUTION**
See Side Panel for Additional
Precautionary Statements

EPA Reg No. 67546-R
EPA Est. 67542-CIIL-001

Manufactured by:
National BoraXX Corp.
3690 Orange Place, Suite 480
Cleveland, Ohio 44122
Emergency Phone: (800) 292-8680

® Reg. Trademark of National BoraXX Corp.

© 1995 National BoraXX Corp.

ACCEPTED
with 50 lbs. (22.7 Kg.)
Im EPA Letter Dated:

JAN 09 2001

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under EPA Reg. No.

67546-2

APPLICATION INSTRUCTIONS

FOR DARKLING AND LESSER MEALWORM BEETLE, HIDE BEETLE, AND FLY (Adults and Larvae).

For Poultry houses/hogbarns/barns in which birds/hogs are grown:

DRY APPLICATION

Prior to application of the Darkling Beetle Control, remove animals. Apply dust uniformly to floor of poultry house/hogbarns/barns or to old litter with a fertilizer or seed spreader, at the rate of 1-2 pounds for each 100 square feet, in bands along feeder lines. Wear a dust mask or a respirator while applying. Spread fresh litter of at least four inches uniformly over all treated area (floor or old litter). Then introduce animals. Reapply after each grow-out, if needed, in the sequence outlined above.

WET APPLICATION

Prior to application of Darkling Beetle Control, remove animals from poultry house/hogbarns/barns. Mix dust at the rate of 1-2 pounds per 3 gallons of water. Apply at the rate of 3 gallons of mix for each 100 square feet of treated surface. Using appropriate pressure spray equipment, saturate side walls, top plates, posts, and framing. Spray mix into cracks and crevices around insulation. Do not expose animals to spray. Reapply after each grow-out, if needed, in the sequence outlined above. However, if facility is washed down and sanitized using standard cleaning and disinfecting procedures, Darkling Beetle Control must be reapplied. Darkling Beetle Control can be applied directly to manure for fly control.

For poultry houses in which birds are grown in cages (layer or high rise 'pit type' houses):

DRY APPLICATION

Prior to application of the Darkling Beetle Control, remove birds. Apply at the rate of 1-2 pounds for each 100 square feet of treated surface. Using appropriate equipment, thoroughly dust side walls, top plates, posts, and framing. Dust into cracks and crevices around insulation. Then introduce birds. Reapply after each grow-out, if needed, in the sequence outlined above. However, if facility is washed down and sanitized using standard cleaning and disinfecting procedures, Darkling Beetle Control must be reapplied to continue beetle control. Darkling Beetle Control can be applied directly to manure for fly control.