

67546-2

02-24-2004

1/2

National Boraxx Corp.
3690 Orange Place, Suite 480
Cleveland OH 44122

FEB 24 2004


FFB

Dear Mr. Kutcel:

Subject: Label Revisions (2001-1)
Product Name: Darkling Beetle Control
EPA Registration No. 67546-2
Application date: December 16, 2003

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended revising the label in compliance with PR Notice 2001-1 is acceptable, and stamped copy is enclosed for your records.

Sincerely,



Richard Gebken
Acting Product Manager 10
Insecticide Branch
Registration Division 7505

enclosure

**PRECAUTIONARY STATEMENTS
HAZARD TO HUMANS AND
DOMESTIC ANIMALS
CAUTION**

Harmful if swallowed. Causes eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling.

FIRST AID	
If in eyes	-Hold eye open and rinse slowly and gently with water for 15-20 minutes. -Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. -Call a poison control center or doctor for treatment advice.
If swallowed	-Call poison control center or doctor immediately for treatment advice. -Have person sip a glass of water if able to swallow. -Do not induce vomiting unless told to do so by the poison control center or doctor. -Do not give anything by mouth to an unconscious person.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-292-8680 for emergency medical treatment information.	

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

STORAGE: Keep container tightly closed when not in use. Store in original container, in a dry place inaccessible to children and pets.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: DO NOT reuse container. Rinse 3 times, then offer for recycling or trash collection.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. See side panel for Application Instructions.

**Darkling Beetle
Control**

**An Insecticide for the Control
of Darkling Beetles, Hide
Beetles, and Flies.**

Active Ingredient:
Boric Acid.....100%

**KEEP OUT OF THE REACH
OF CHILDREN
CAUTION**
See Side Panel for Additional
Precautionary Statements

EPA Reg No. 67546-2
EPA Est. 67542-CHL-001

Manufactured by:
National BoraXX Corp.
3690 Orange Place, Suite 480
Cleveland, Ohio 44122
Emergency Phone: 1-800-292-8680

®Reg. Trademark of National BoraXX Corp.

©1995 National BoraXX Corp.
ACCEPTED
FEB 24 2004
Net Contents: 50 lbs. (22.7 Kg.)
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticides registered under EPA Reg. No. 67546-2

APPLICATION INSTRUCTIONS
FOR DARKLING AND LESSER MEALWORM BEETLE,
HIDE BEETLE, AND FLY (Adults and Larvae).

For Poultry houses/hogbarns/barns in which birds/hogs are grown:

DRY APPLICATION

Prior to application of the Darkling Beetle Control, remove animals. Apply dust uniformly to floor of poultry house/hogbarns/barns or to old litter with a fertilizer or seed spreader, at the rate of 1-2 pounds for each 100 square feet, in bands along feeder lines. Wear a dust mask or a respirator while applying. Spread fresh litter of at least four inches uniformly over all treated area (floor or old litter). Then introduce animals. Reapply after each grow-out, if needed, in the sequence outlined above.

WET APPLICATION

Prior to application of Darkling Beetle Control, remove animals from poultry house/hogbarns/barns. Mix dust at the rate of 1-2 pounds per 3 gallons of water. Apply at the rate of 3 gallons of mix for each 100 square feet of treated surface. Using appropriate pressure spray equipment, saturate side walls, top plates, posts, and framing. Spray mix into cracks and crevices around insulation. Do not expose animals to spray. Reapply after each grow-out, if needed, in the sequence outlined above. However, if facility is washed down and sanitized using standard cleaning and disinfecting procedures, Darkling Beetle Control must be reapplied. Darkling Beetle Control can be applied directly to manure for fly control.

For poultry houses in which birds are grown in cages (layer or high rise 'pit type' houses):

DRY APPLICATION

Prior to application of the Darkling Beetle Control, remove birds. Apply at the rate of 1-2 pounds for each 100 square feet of treated surface. Using appropriate equipment, thoroughly dust side walls, top plates, posts, and framing. Dust into cracks and crevices around insulation. Then introduce birds. Reapply after each grow-out, if needed, in the sequence outlined above. However, if facility is washed down and sanitized using standard cleaning and disinfecting procedures, Darkling Beetle Control must be reapplied to continue beetle control. Darkling Beetle Control can be applied directly to manure for fly control.

NOTICE

Manufacturer shall not be liable in respect to any injury or consequential damages suffered based on the use of this product contrary to the instructions given on this label.

2/2